

ARCTIC DATA COMPILATION AND APPRAISAL VOLUME 4

Northwest Passage: Chemical Oceanography

D.J. Thomas¹, R.W. Macdonald², A.G. Francis¹,
V. Wood¹, and A.B. Cornford²

¹Arctic Laboratories Limited
Sidney, B.C., V8L 3S1
and

²Institute of Ocean Sciences
Department of Fisheries and Oceans
Sidney, B.C., V8L 4B2

1983

**CANADIAN DATA REPORT OF
HYDROGRAPHY AND OCEAN SCIENCES
NO. 5**



Canada

Canadian Data Report Of Hydrography and Ocean Sciences

These reports provide a medium for the documentation and dissemination of data in a form directly useable by the scientific and engineering communities.

Generally, the reports will contain raw and/or analyzed data but will not contain interpretations of the data. Such compilations will commonly have been prepared in support of work related to the programs and interests of the Ocean Science and Surveys (OSS) sector of the Department of Fisheries and Oceans.

Data Reports are produced regionally but are numbered and indexed nationally. Requests for individual reports will be fulfilled by the issuing establishment listed on the front cover and title page. Out of stock reports will be supplied for a fee by commercial agents.

Regional and headquarters establishments of Ocean Science and Surveys ceased publication of their various report series as of December 1981. A complete listing of these publications and the last number issued under each title are published in the *Canadian Journal of Fisheries and Aquatic Sciences*, Volume 38: Index to Publications 1981. The current series began with Report Number 1 in January 1982.

Rapport statistique canadien sur l'hydrographie et les sciences océaniques

Ces rapports servent de véhicule pour la compilation et la diffusion des données sous une forme directement utilisable par les scientifiques et les techniciens.

En général, les rapports contiennent des données brutes ou analysées mais ne fournissent pas d'interprétations des données. Ces compilations sont préparées le plus souvent à l'appui de travaux reliés aux programmes et intérêts du service des Sciences et Levés océaniques (S.L.O.) du ministère des Pêches et des Océans.

Les rapports statistiques sont produits à l'échelon régional mais sont numérotés et placés dans l'index à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page de titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Les établissements des Sciences et Levés océaniques dans les régions et à l'administration centrale ont cessé de publier leurs diverses séries de rapports depuis décembre 1981. Vous trouverez dans l'index des publications du volume 38 du *Journal canadien des sciences halieutiques et aquatiques*, la liste de ces publications ainsi que le dernier numéro paru dans chaque catégorie. La nouvelle série a commencé avec la publication du Rapport n° 1 en janvier 1982.

**CANADIAN DATA REPORT OF
HYDROGRAPHY AND OCEAN SCIENCES NO. 5**

1983

**ARCTIC DATA COMPILATION AND APPRAISAL
VOLUME 4**

Northwest Passage: Chemical Oceanography

by

**D.J. Thomas¹, R.W. Macdonald², A.G. Francis¹,
V. Wood¹ and A.B. Cornford²**

**1 Arctic Laboratories Limited
Sidney, B.C., V8L 3S1**

**2 Institute of Ocean Sciences
Sidney, B.C., V8L 4B2**

**Institute of Ocean Sciences
Department of Fisheries and Oceans
Sidney, B.C., V8L 4B2**

PREFACE

These catalogues are produced by the Ocean Information Division at the Institute of Ocean Sciences. Joint government and industry contract projects have catalogued marine data sets their focus being primarily upon oceanography and fisheries. Data set quality appraisals are included to assist in establishing the usefulness of certain data for particular kinds of analyses and the confidence to be placed in interpretations. These appraisals will assist in setting priorities for incorporating the most useful data in the national Marine Environmental Data Service (MEDS) archives. Additional uses include research planning (especially for climatological studies), and the provision of the best available resume of marine data sources for environmental assessments.

The accelerating pace of Arctic offshore development activity has emphasized the need to review the sufficiency and suitability of available scientific information for design, regulatory and planning purposes. This review has been divided into three phases: (1) compilation and appraisal of all existing data sets; (2) analysis of the suitability of the historical data for contributing to questions of particular interest; and (3) analysis and interpretation of data and estimation of the scientific confidence in answering particular questions. This report on the chemical oceanographic data of the channels of the Northwest Passage is a contribution to the first phase.

Dr. Alan B. Cornford
Scientific Editor
Arctic Data Compilation
and Appraisal Series

Copyright Minister of Supply and Services Canada - 1983

Cat. No. Fs 97-16/5 ISSN 0711-6721

The correct citation for this publication is:

D.J. Thomas, R.W. Macdonald, A.G. Francis, V. Wood and A.B. Cornford. Arctic Data Compilation and Appraisal. Volume 4. Northwest Passage: Chemical Oceanography.

Can. Data Rep. Hydrogr. Ocean Sci. 5: (Volume 4) 200 pp.

TABLE OF CONTENTS

	Page
PREFACE	ii
TABLE OF CONTENTS	iii
ABSTRACT	v
ACKNOWLEDGEMENTS	vi
SPECIAL CREDITS	vii
VOLUME ABSTRACT	1
1. INTRODUCTION	1
2. STUDY AREA	4
3. CHEMICAL DATA PRESENTATION	4
3.1 Types of data	4
3.2 Concentration units	9
4. OUTLINE OF DATA INVENTORY ORGANIZATION	9
4.1 Description of table headings	14
4.1.1 Table 1	14
4.1.2 Table 2	14
4.1.3 Table 3	15
4.2 Sample use of the inventory	16
5. DATA RATING SYSTEM	16
5.1 Data quality criteria	16
5.2 Definition of the rating system	23
5.3 Significance of the data rating value	25
5.4 Effect of positioning on the data quality	26

TABLE OF CONTENTS (continued)

	Page
5.5 Some important factors relating to the data appraisal process	27
5.5.1 Heavy metals	27
5.5.2 Nutrients	33
5.5.3 Dissolved oxygen	35
5.5.4 Hydrocarbons	35
5.5.5 Chlorinated hydrocarbons	37
6. REFERENCES	38
7. DATA INVENTORY TABLE 1. DATA SET LISTING	47
8. DATA INVENTORY TABLE 2. LISTING OF DATA BY PARAMETER	75
9. DATA INVENTORY TABLE 3. LISTING OF MEASUREMENT LOCATIONS	111
10. MAPS	145
10.1 Geographical occurrence of chemical oceanographic data on a cumulative bi-monthly basis	145
10.2 Geographical occurrence of chemical oceanographic data on a yearly basis	155
11. INDICES	191
11.1 Geographical	193
11.2 References by data set number	194

ABSTRACT

D.J. Thomas, R.W. Macdonald, A.G. Francis, V. Wood and A.B. Cornford. Arctic Data Compilation and Appraisal. Volume 4. Northwest Passage: Chemical Oceanography.

Can. Data Rep. Hydrogr. Ocean Sci. 5: (Volume 4) 200 pp.

This volume is one of a group of catalogues designed to compile and appraise marine data sets for the Canadian Arctic. For ease of reference, the group has been organized with its subject matter divided into three disciplines: physics, chemistry and biology. The Arctic has been arbitrarily divided into seven geographical areas to include, where possible, major oceanographic regions. The format has been structured to facilitate comparison between subjects and regions. With such a large undertaking it is not possible to provide all reports at once. Therefore catalogues which are presently available in the series are indicated on the inside back cover of each volume.

Data collection is a continuing process and further updates of the catalogues are planned. Readers are invited to submit corrections and additions by writing the issuing establishment. These corrections will be incorporated in on-line computerized data set listings; they will be continuously available upon request.

SOMMAIRE

D.J. Thomas, R.W. Macdonald, A.G. Francis, V. Wood and A.B. Cornford. Arctic Data Compilation and Appraisal. Volume 4. Northwest Passage: Chemical Oceanography.

Can. Data Rep. Hydrogr. Ocean Sci. 5: (Volume 4) 200 pp.

Le présent volume fait partie d'un groupe de catalogues destinés à compiler et à évaluer les séries de données marines sur l'Arctique canadien. Pour plus de commodité, la question traitée est structurée en trois grandes disciplines: physique, chimie et biologie. L'Arctique a été divisé arbitrairement en sept régions géographiques qui englobent autant que possible les grandes régions océanographiques. Les catalogues sont présentés de façon à faciliter la comparaison entre les sujets et les régions. Le domaine est si vaste qu'il est impossible de fournir tous les catalogues en une seule fois. Les catalogues de la série actuellement disponibles sont indiqués à la fin de chaque volume à l'intérieur de la couverture.

La collecte des données est un processus permanent et il est prévu de mettre à jour les catalogues par la suite. Les lecteurs sont invités à soumettre par écrit les corrections et les additions à l'établissement auteur. Ces corrections seront traitées en direct sur ordinateur et incorporées aux listes qui pourront être obtenus sur demande.

ACKNOWLEDGEMENTS


The authors wish to acknowledge contributions from many people who assisted with the preparation of this inventory compilation. Brian Smiley of the Institute of Ocean Sciences, Ocean Information Division contributed to format design and provided valuable comment and criticism throughout the project. Ms. T. Yu and Mr. B. Perry of Arctic Laboratories Limited provided dedicated technical support in organizing original data sources. Special thanks are extended to J. Gagnon of the Marine Environmental Data Service (MEDS) in Ottawa and H. Odum of the National Oceanographic Data Center (NODC) in Washington, D.C., U.S.A., for providing access to their data files. Information and assistance were received from Dr. B. Bornhold, Pacific Geoscience Centre, Sidney, B.C., Dr. A. Sekerak, LGL Limited and Dr. M. Dunbar, Marine Sciences Centre, McGill University. P. Rothwell performed the arduous task of preparing the manuscript.

SPECIAL CREDITS


L.S.C. Thomson - Technical editing

Funds for this work were provided by:

Department of Fisheries and Oceans (DFO);
Department of Indian Affairs and Northern Development (DIAND); and
Dome Petroleum Limited



ARCTIC DATA COMPILATION AND APPRAISAL
VOLUME 4
NORTHWEST PASSAGE: CHEMICAL OCEANOGRAPHY



The area (2) covered by this volume is shaded on the map above.

Volume 4: Northwest Passage: Chemical Oceanography

VOLUME ABSTRACT

This inventory contains a catalogue of chemical oceanographic data sets from the Northwest Passage. The inventory includes commonly- measured substances such as dissolved oxygen, major and minor elemental components, nutrients and less frequently measured substances such as trace elements, hydrocarbons and chlorinated hydrocarbons. Suspended particulate matter (although not a truly chemical quantity) is also included. Data sets are included for sea ice, sea water, sediments and biota. Times and locations of measurements are listed and displayed graphically on a yearly and cumulative bi-monthly basis. A geographical index and alphabetical references (by data set number) are also included.

Key Words: Northwest Passage, chemical oceanography, data sets, inventory, dissolved oxygen, nutrients, heavy metals, hydrocarbons, chlorinated hydrocarbons, suspended particulate material, sea ice, sediments, biota.

1. INTRODUCTION

This inventory comprises 75 data sets of chemical oceanographic data collected in the marine channels of the Northwest Passage during the period 1928 - 1981. The quantity and type of data are distributed irregularly over this time period with the bulk of the data collection occurring after 1970 as shown in Figure 1.1.




Figure 1.1 Yearly distribution of chemical oceanographic data sets for the Northwest Passage.

Most of the data reported in this inventory can be classified as baseline data and are statements of observed distributional patterns of the various chemical species. Before 1970, water column measurements such as dissolved oxygen and nutrients were virtually the only data obtained. After 1970, a wider range of data were obtained in response to a need for environmental data by government and industry interested in exploiting regional resources but usually on a very narrow geographical basis only, for example: (1) most of the data for heavy metals in sea water, sediments and biota apply to Strathcona Sound where extensive sampling programs were concerned with potential effects of a lead-zinc mining operation at Nanisivik and (2) most of the hydrocarbon data were collected in the Lancaster Sound area in anticipation of petroleum hydrocarbon resource development. The frequency of occurrence of selected chemical quantities is summarized for each sampling medium (sea water, ice, sediment and biota) in Table 1.1.

There is no evidence in the published literature for any systematic interpretation of the chemical oceanographic data to explain the observed distribution patterns in terms of fundamental oceanographic processes. The main reason for this is the limited data available.

The objective of this inventory was to compile all available Northwest Passage chemical oceanographic data sets into a single data bank and to appraise the quality of the data so as to obtain, for the first time, a broader perspective on what is currently known about the chemical oceanography of the marine channels of the Northwest Passage.

The inventory is ongoing. As new data and previously inaccessible data become available, they will be added to a computerized data base maintained at the Institute of Ocean Sciences, Sidney, B.C. Information about new data sets, older data sets which do not appear in this inventory or errors in this inventory, should be submitted in writing to the Institute of Ocean Sciences.

The following sections contain the rationale for organizing the data as it appears in the tables. Wherever possible, formats were adopted that were consistent with those used in the companion inventory for the physical oceanography of the Northwest Passage (Birch et al., 1983).

TABLE 1.1

FREQUENCY OF OCCURRENCE OF SELECTED CHEMICAL
QUANTITIES FOR EACH SAMPLING MEDIUM

	Sea Water	Ice Cores	Sediments	Biota
Dissolved Oxygen	34	-	-	-
pH	3	-	-	-
Nutrients				
Nitrate	18	2	-	-
Phosphate	23	1	-	-
Silicate	17	-	-	-
Ammonia	2	-	-	-
Chl. a	14	3	-	-
Phaeopigments	2	2	-	-
Suspended Particulates	3	-	-	-
Isotopes	2	-	-	-
Alk _t	1	-	-	-
Carbon	-	-	4	-
Heavy Metals				
Vanadium	-	-	-	1
Chromium	-	-	2	1
Manganese	1	-	4	-
Iron	2	-	6	4
Cobalt	-	-	3	-
Nickel	2	-	4	1
Copper	3	-	5	5
Zinc	2	-	6	7
Arsenic	1	-	5	7
Selenium	-	-	-	4
Silver	-	-	2	-
Cadmium	3	-	7	7
Mercury	-	-	6	7
Lead	2	-	7	6
Hydrocarbons				
Methane	1	-	-	-
Oil	1	-	1	-
Pesticides	-	-	-	4
PCB	-	-	-	4
Tar	6	-	-	-
Hexane/Benzene				
Extractable Compounds	5	-	-	-

2. STUDY AREA

The study area is shown in Figure 2.1; it comprises about 10^5 km^2 , approximately the same area as Baffin Bay or Hudson Bay. The marine channels of the Northwest Passage are usually ice-congested, even in summer; it was not until 1905, two years after entering Lancaster Sound that the Gjoa, a seventy-two foot long converted herring boat commanded by Roald Amundsen, completed the Northwest Passage by sea (Mirsky, 1934).

The islands of the Canadian Archipelago are the exposed parts of the continental shelf which extends north from the mainland. The shores of the southern channels are composed of Precambrian rock of the Canadian Shield which extends up the Boothia Peninsula and Baffin Island. Glaciers covered much of the area during the last ice age. Raised beaches, resulting from isostatic rebound, are widespread today except for the west coast of Banks Island which appears to be sinking. Ice scouring is evident in the surface and submarine topography (Dunbar and Greenaway, 1956).

The marine channels of the Northwest Passage are the submerged portion of the continental shelf. Water depths exceed 500 m in Lancaster Sound and M'Clure Strait; a sill of about 150 m depth in Barrow Strait separates these two regions. Water depths in the southern channels are relatively shallow, about 50 m. The waters of the Northwest Passage are a transition between the Arctic Water Mass to the west and Atlantic and Arctic waters to the east. The sill in Barrow Strait limits the easterly flow of Arctic water to the upper 150 m of the water column. The warmer, more saline Atlantic water is found below the Arctic layer and usually occurs only in the deeper waters of eastern Lancaster Sound.

3. CHEMICAL DATA PRESENTATION

3.1 Types of Data

All chemical data have been grouped according to the environmental medium or compartment in which they are found, as follows:




Figure 2.1 Northwest Passage data compilation study area.

Medium	Constituents Included
Sea Ice	- dissolved or occluded
Water Column (Sea Water)	- dissolved constituents - particulate constituents
Sediments	- surficial sediments - sediment cores (interval sampling) - interstitial pore waters
Biota (flora and fauna)	- seawater dwelling organisms - bottom sediment dwelling organisms - marine mammals

The inventory includes all available data of a "chemical nature". This includes commonly-measured substances such as dissolved oxygen, major and minor elemental components, nutrients and less frequently measured substances such as trace elements, hydrocarbons and chlorinated hydrocarbons. Turbidity and suspended particulate material are not truly chemical quantities in the classical sense, but are included in the inventory because they are important factors in the interpretation of chemical data and because they are more logically included with the chemical inventory rather than the physical or biological inventories.

The largest amount and most diverse data are found for water column constituents. Field-based analyses of samples at the time of collection have been rare because most chemical analyses require specialized or sophisticated equipment. Water samples have been frequently processed in the field to the stage where sample preservation is convenient and then returned to the laboratory for analysis. Samples for dissolved oxygen and pH analysis, on the other hand, have been routinely analysed in the field shortly after collection. Other determinations which can and have been made in the field include methane and the reactive nutrients although the latter have also been preserved and returned to the laboratory. Measurements have been made rarely in situ, and then only for dissolved oxygen.

The other types of samples: biota, sediments and sea ice, have been obtained in decreasing quantities respectively and all have been analysed exclusively in the laboratory after preservation for some extended period of time. A summary of the chemical data types included in the inventory is shown in Table 3.1.

TABLE 3.1
A SUMMARY OF CHEMICAL DATA TYPES INCLUDED
IN THE DATA INVENTORY

Chemical Constituent	Water	Ice Cores	Sediment	Biota
adenosine triphosphate (ATP)	x			
alkalinity (total)	x			
ammonia	x			
arsenic	x		x	x
benzene extractable compounds (BEC)	x			
cadmium	x		x	x
calcium carbonate			x	
carbon	x			
carbonate	x			
carbon dioxide			x	
¹³ C/ ¹² C isotopic ratio	x			
particulate organic carbon	x			
total organic carbon			x	
chlorophyll <u>a</u>	x	x		
chlorophyll <u>b</u>		x		
chlorophyll <u>c</u>		x		
total chlorophyll	x			
chromium				x
cobalt			x	
copper	x		x	x
deoxyribonucleic acid (DNA)	x			
ethane, 2,2-Bis(p-chlorophenyl)-1,1-dichloro- (i.e. p,p'-DDD)				x
ethane, 1-(o-chlorophenyl)-1-(p-chlorophenyl)-2,2,2-trichloro- (i.e. o,p'-DDT)				x
ethane, 1,1-Bis-(p-chlorophenyl)-2,2,2-trichloro (i.e. p,p'-DDT)				x
ethylene, Bis-(chlorophenyl)dichloro- (i.e. DDE)				x
ethylene, 2,2-Bis(p-chlorophenyl)-1,1-dichloro- (i.e. p,p'-DDE)				x
helium				
dissolved	x			
³ He/ ⁴ He isotopic ratio	x			
hexane extractable compounds (HEC)	x			
hexachlorobenzene (HCB)				x
hydrogen				
particulate organic	x			
iron	x		x	x
lead	x		x	x

TABLE 3.1 (continued)
A SUMMARY OF CHEMICAL DATA TYPES INCLUDED
IN THE DATA INVENTORY

Chemical Constituent	Water	Ice Cores	Sediment	Biota
manganese	x		x	
mercury				
elemental			x	x
methyl				x
methane	x			
neon	x			
nickel	x		x	x
nitrogen	x			
nitrate	x	x		
nitrite	x			
organic			x	
total dissolved	x			
particulate organic	x			
oil and grease	x		x	
oxygen				
dissolved	x			
$^{18}\text{O}/^{16}\text{O}$ isotopic ratio	x			
pH	x			
phaeopigments	x	x		
phosphorus				
phosphate	x	x		
soluble reactive	x			
total dissolved	x			
polychlorinated biphenyls (PCB)				x
protein	x			
ribonucleic acid (RNA)	x			
selenium				x
silicate	x			
silicon	x			
silver			x	
starch grains	x			
suspended particulate matter (SPM)	x			
tarballs	x			
tritium	x			
vanadium				x
zinc	x		x	x

3.2 Concentration Units

Several different concentration units have been used over the years to report the chemical results for Northwest Passage data. In order to eliminate confusion and provide for ease of data comparison between data sets, the International System of Units (Système International, SI) has been used wherever possible. Exceptions in the use of SI are classes of compounds comprising assemblages of different molecules such as polycyclic aromatic hydrocarbons and polychlorinated biphenyls. The trend toward worldwide use of SI units as a standard has been established by resolutions of the General Conference of Weights and Measures. In addition, the International Union of Pure and Applied Chemistry (IUPAC) endorses the exclusive use of SI units for chemical quantities. An important advantage of using SI units for chemical oceanographic and geochemical data is the ease by which mass fluxes and chemical budgets can be calculated and compared for various elements and chemical compounds.

Factors used to convert units found in original reports to SI units are listed in Table 3.2.

4. **OUTLINE OF DATA INVENTORY ORGANIZATION**

The data are organized into a chronological series of data sets beginning with the year 1928. No chemical data collected before 1928 could be found. Each data set comprises sampling or chemical measurements taken during a single cruise, or during a sampling excursion usually by a single agency. It is assumed, then, that data within a given data set have been collected uniformly and should be internally consistent insofar as sampling methodology is concerned.

Each data set has been assigned an identification number of the form yy-nnnn, where yy = last 2 digits of the year in which data were collected and nnnn = order of identification for that particular data set for that year. The data set number is a unique identifier which applies throughout this series of inventories; for example, any data set identified as 72-0011 is the same no matter where the reference to it is made. In certain cases, data may have been collected over a period of months or years by a common study team with minor or major differences occurring in the types of data collected at each sampling period. When this occurred, letters were used as a suffix to the data set number to distinguish the various sample

TABLE 3.2

CONVERSION FACTORS AS NUMERICAL MULTIPLES OF SI UNITS

CHEMICAL QUANTITY	TO CONVERT FROM*	TO	MULTIPLY BY
	litres mg L ⁻¹	m ³ g m ⁻³	1,000 1
ammonia	mg L ⁻¹ (NH ₃) µg at L ⁻¹ (NH ₃)	mmol m ⁻³ (NH ₃) mmol m ⁻³ (NH ₃)	58.82 1
arsenic	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	13.35 13.35
cadmium	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	8.90 8.90
calcium	µg L ⁻¹	µmol m ⁻³	24.95
calcium carbonate	% (wt/wt)	mol kg ⁻¹	0.0999
carbon	µg g ⁻¹	µmol kg ⁻¹	83.26
carbon dioxide	% (wt/wt)	mol kg ⁻¹	0.227
chromium	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	19.23 19.23
cobalt	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	16.97 16.97
copper	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	15.74 15.74
p,p'-DDD	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	3.13 3.13
p,p'-DDE	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	3.14 3.14
p,p'-DDT	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	2.82 2.82
o,p'-DDT	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	2.82 2.82
iron	µg L ⁻¹ % Fe (w/w)	µmol m ⁻³ mol kg ⁻¹	17.91 0.179

CONVERSION FACTORS AS NUMERICAL MULTIPLES OF SI UNITS

CHEMICAL QUANTITY	TO CONVERT FROM*	TO	MULTIPLY BY
	litres mg L ⁻¹	m ³ g m ⁻³	1,000 1
lead	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	4.83 4.83
manganese	µg L ⁻¹ % Mn (w/w)	µmol m ⁻³ mol kg ⁻¹	18.20 0.182
mercury	ng L ⁻¹ ng g ⁻¹	nmol m ⁻³ ng kg ⁻¹	4.99 4.99
methane	nL (STP) L ⁻¹	nmol m ⁻³	44.64
nickel	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	17.03 17.03
nitrogen	mg L ⁻¹ mg kg ⁻¹	mmol m ⁻³ mmol kg ⁻¹	71.43 0.0714
nitrate	mg L ⁻¹ (NO ₃) µg at L ⁻¹ (NO ₃ -N)	mmol m ⁻³ (NO ₃ -N) mmol m ⁻³ (NO ₃ -N)	16.13 1
nitrite	mg L ⁻¹ (NO ₂) µg at L ⁻¹ (NO ₂ -N)	mmol m ⁻³ (NO ₂ -N) mmol m ⁻³ (NO ₂ -N)	20.83 1
oxygen	mg L ⁻¹ (O ₂) mL L ⁻¹ (O ₂)	mol m ⁻³ (O ₂) mol m ⁻³ (O ₂)	0.0313 0.0446
phosphate	mg L ⁻¹ (PO ₄) µg at L ⁻¹ (PO ₄ -P)	mmol m ⁻³ (PO ₄ -P) mmol m ⁻³ (PO ₄ -P)	32.29 1
phosphorous	µg L ⁻¹	mmol m ⁻³ (P)	0.0324
potassium	mg L ⁻¹	mmol m ⁻³	25.58
selenium	mg L ⁻¹	mmol kg ⁻¹	0.0127
silicon	mg L ⁻¹ (Si) mg L ⁻¹ (SiO ₂) µg at L ⁻¹ (silicate-Si)	mmol m ⁻³ (Si) mmol m ⁻³ (Si) mmol m ⁻³ (silicate-Si)	35.60 16.64 1
vanadium	µg g ⁻¹	µmol kg ⁻¹	19.63

CONVERSION FACTORS AS NUMERICAL MULTIPLES OF SI UNITS

CHEMICAL QUANTITY	TO CONVERT FROM*	TO	MULTIPLY BY
	litres mg L ⁻¹	m ³ g m ⁻³	1,000 1
zinc	µg L ⁻¹ µg g ⁻¹	µmol m ⁻³ µmol kg ⁻¹	15.30 15.30

* Note:

In Table 2 conversions for dissolved constituents have been made assuming that the density of sea water is 1.00 i.e., that 1 µg L⁻¹ is equivalent to 1 µg kg⁻¹.

collections. For example, data set 73-0013 is divided into three parts in the inventory - 74-0013A, 74-0013B and 74-0013C. While there is insufficient reason to regard the three as separate data sets because all samples were collected during a single cruise, the subdivision is made to emphasize that different parameters were sampled by three separate groups of scientists during the sampling period. Gaps may appear in the sequence of data set numbers in this inventory for a particular year, because each data set will not appear in every discipline and geographical area.

This inventory comprises three main tables followed by supporting figures and tables. Table 1 is a chronological list of data sets by data set number (see above). Table 2 is a summary of specific details for actual chemistry data. Table 3 is a listing of times and locations of individual measurements. Where station coordinates were not specified in reports, approximate station positions were obtained by measuring plotted station locations on figures located in the report. Measurement locations are plotted in a series of maps in Section 10. There are four standard maps; all are Lambert Conformal Conic projection. The map encompassing the study area is drawn to a scale of 1:7 000 000. The other four maps are drawn to a scale of 1:4 000 000; they yield better resolution of data sets containing closely spaced stations. In all cases, the coastlines have been smoothed and small islands removed to avoid clutter. In data sets where stations are very closely spaced a single symbol is used to reduce smudging. In these cases, the number in parentheses to the right of symbol refers to the number of stations at the location represented by the symbol. To maintain uniformity and facilitate comparison, maps for all volumes in the Arctic data compilations have been drawn from common stock.

A listing of the geographical occurrence of data sets is given in the geographic index. Section 11.2 is an index of references ordered by data set number. The first (primary) reference shown for each data set is the original data report or similar document. The secondary references that follow are other reports or refereed papers based on the primary reference. The listing of secondary references should not be considered an exhaustive literature search. Only those secondary references are included as were found while searching for original data set documents.

4.1 Description of Table Headings

4.1.1 Table 1

Table 1 provides general details of sampling excursions and includes:

- (1) identification of the specific region within the study area where sampling was conducted;
- (2) the period of time during which the measurements were made;
- (3) the ship or agency which collected the data;
- (4) a listing of the chemical quantities measured or sampled during the collection period, with those quantities for which data are published and available for inclusion in this inventory in bold type. All entries are not in bold type because
 - (a) all samples collected on original cruises may not have been analysed;
 - (b) samples may have been analysed but results may not be available in the public domain or may be untraceable;
 - (c) samples may be lost or destroyed.
- (5) concurrent physical and biological measurements or samples.

4.1.2 Table 2

In Table 2, specific details including analytical results, are given for each quantity measured in each data set. These include:

- (1) total number of stations sampled;
- (2) total number of samples obtained at all stations;
- (3) the number of samples having analytical results greater than the analytical detection limit or greater than 0 when 0 is used to designate the detection limit;
- (4) methodology information: details of collection, storage (preservation), and analysis of the sample allowing for judging quality and comparability of data;

- (5) the range of reported concentrations. Note that all concentrations are given in Systeme International (SI) units. Conversions between these and previously used units are given in Table 3.1;
- (6) the mean and median of the reported concentrations. The median is included because it is not as easily influenced by extreme values as is the mean. It thus represents a better estimate of the middle of a sample population with skewed distribution. Many environmental parameters fall into a log-normal distribution, so the median is probably a better estimate of central tendency. If the number of observations is odd, the median is the middle one of the observations; if the number is even, the median is the average of the two innermost observations. Where a suite of results includes detection limit values, the detection limit is used for computing the mean;

Note: pH is a logarithmic function. It is not meaningful to express a mean value because the mean can be strongly biased to low pH values; consequently means for pH values are not given.
- (7) a data quality (confidence) rating based on the rules outlined subsequently (by Section 5.2). The data rating scale uses values from 0 to 4 with 4 indicating data judged to be highest quality, and to have the most versatile potential (refer to Section 5.3).

4.1.3. Table 3

Table 3 provides specific spatial and temporal details for collected samples. These include:

- (1) station position (latitude, longitude). For stations where no latitude/longitude are expressed, estimates were made by direct measurement of the plotted station points contained in individual original reports;
- (2) station depth;
- (3) sampling time (year, month, day and hour in GMT or local time);
- (4) number of points (samples in profiles); and
- (5) maximum depth sampled.

4.2 Sample use of the inventory

Example 1 Searching for specific-parameter data: e.g., Copper in sea water

Step 1. Consult Table 1 and scan column labelled "Chemical quantities measured or sampled".
Note data sets listing Copper (in bold print) as a measured parameter

Step 2. Consult data set numbers noted in Step 1, Table 2, to obtain specific details of sample history and reported concentrations.

Step 3. Refer to Table 3 for station positions, depths sampled, etc. If areal coverage of stations is of interest, go to Section 6 and consult maps for the data set(s) of interest. Maps are ordered chronologically.

Step 4. Consult the reference list for reports or publications upon which the data set is based.

Example 2. Searching for data from a specific geographic area:
e.g. Copper in McKinley Bay sea water

Step 1. Consult Geographical Index
Note data set numbers

Step 2. Consult Table 1 for data sets reporting copper in sea water.

Step 3. Continue as in Example 1.

When additional details concerning data are required, the original data sets must be consulted. Access to these documents may be obtained through Ocean Information Division at the Institute of Ocean Sciences, Sidney, British Columbia.

5. DATA RATING SYSTEM

5.1 Data Quality Criteria

The reported chemical data for the Northwest Passage have been appraised using a rating system based on five data quality parameters related to methodology:

- (A) sample collection;
- (B) sample storage/preservation;
- (C) sample analysis;
- (D) analytical precision; and
- (E) analytical accuracy.

These five parameters satisfy a fundamental requirement that the rating criteria allow an assessment to be made of any chemical result at any point in the history of a sample from collection to final analysis. In this way, the five parameters shown above represent a measure of the ultimate believability of results.

A quiet revolution has been taking place in recent years in the approach analytical chemists have for quality control and the establishment of well-defined performance criteria for analytical procedures. The most significant step was the general recognition that analytical quality control has been almost universally inadequate in the areas of environmental chemistry. The residual problem is resistance to change; routine analysis and reporting procedures that have been in use for many years will not change overnight. The new concepts as articulated by Taylor (1983), Kirchmer (1983), Long and Winfordner (1983), Glaser et al., (1981) and Cheeseman and Wilson (1978) will eventually be practised and improved by analytical chemists. This evolutionary process will lead to results for environmental chemical samples far superior to those summarized in this inventory.

A general caution is appropriate at this time. A great deal of chemical data has been collected over the years by biologists. The objective in sampling has often been to merely "get a rough idea of metal contents or hydrocarbon contents etc." rather than produce state-of-the-art results. There is an insidious danger in this approach to chemical oceanography and science in general; namely, that it encourages a lower level of care in all aspects of the methodology chain which, when added together, leads to far greater errors than originally intended or considered to be within acceptable limits. For this reason, all sampling teams should always strive for the best results within their capability rather than those somewhat below.

Some general concepts and comments related to the five data quality criteria introduced earlier which indicate how they can be used in the data evaluation are discussed below. A more detailed discussion for specific chemical constituents is given for Section 5.3.

A. Sample Collection

Sampling has traditionally received little attention and has been the weakest link in marine chemical measurements. The method of sampling is crucial, especially for heavy metals or trace hydrocarbons where baseline values are often at or near the detection limit of many analytical techniques. Very specific steps involving sample preparation and collection methodology must be followed with fanatical attention to detail so as to limit the effects of negative and positive contamination. These details are an integral part of the final reported number and must be specified with the results.

B. Sample Storage and Preservation

Once the sample is collected, it must be preserved in such a way that it remains representative of the water body, ice core, sediment, plant or animal from which it was collected or sampled. Storage containers are very important. For the storage of samples for the determination of heavy metals in sea water, for example, a severe hot acid pretreatment of plastic or Teflon storage bottles must be achieved. Bottles used for storing hydrocarbon samples must be cleaned with solvent and baked to rid the containers of contaminating substances. Procedures specifically applicable to samples of other chemical substances are often necessary. Certain types of samples (e.g., nutrients, chlorophyll a, etc.) which are sensitive to change through biological activity must be analysed or preserved immediately after collection before such activity begins to alter the sample irreversibly. Failure to do so will result in samples becoming unrepresentative of the original water mass. Acid is frequently used as a preservative for seawater samples for heavy metal analysis. Because the concentration of heavy metals is exceedingly low in sea water, only acids with the highest level of purity can be used, lest the acid addition introduce more metals into the samples than those occurring naturally. Consequently, the purity of all preservatives must be carefully tested before use and must be specified in collection details. Length of time between collection and preservation should be reported exactly.

C. Analysis

Assessing the comparability of analytical results depends on a detailed description of the chemical protocol used. The protocol is a set of definitive directions that must be followed, without exception, if the analytical results are to be accepted for a given purpose. Protocol fits into the hierarchy of methodology as outlined by Taylor (1983) in the sequence **technique→method→procedure→protocol**.

The simple statement found in many reports that "sterile procedures were used" or that "standard methods were used" does not provide enough information to form an opinion about the reliability of the technique employed or the validity of results obtained. This follows from the fact that most chemical laboratories dealing with the analysis of marine samples recognize that most procedures are not routine and that operational changes are often made. Thus, the instrumentation and analytical conditions employed; the quality and age of reagent chemicals; the values of reagent and procedural blanks; and the finesse of the analyst with the technique will be critical to the eventual outcome of the analysis. All these details must be specified. Also, quantitative proof that the analytical measurement is in control must be provided. Otherwise, the situation articulated by Eisenhart (quoted in Taylor, 1981) will probably be all too true: "Until a measurement operation has attained a state of statistical control, it cannot be regarded in any logical sense as measuring anything at all".

Detection Limits and Blanks

Important performance criteria of an analytical method are detection limit and blanks. Many definitions for the term "detection limit" can be found in the literature. A consensus appears near. In qualitative terms the limit of detection is the lowest concentration of an analyte that can be reliably distinguished from an analytical blank (Long and Winefordner, 1983). Over the years, the calculated limit of detection for an element has varied as much as an order of magnitude through the use of different statistical approaches and different interpretations of what detection limit really is. This has important implications for the chemical oceanographic data for the Northwest Passage because many data sets contain detection limit values or values reported simply as "less than detection limit" with the detection limit

unspecified. No descriptions are provided in any of the data sets in sufficient detail (in most cases zero detail) so that no data at the lower end of the reported ranges can be used with any degree of certainty.

Blanks are another performance criterion for an analytical process. Highly variable blanks lead to very poor estimates for the detection limit. All data sets are inadequate in describing analytical blank procedures or values for the blanks. The combination of no information for the detection limit and blanks together with many data sets reporting variable detection limits leads to a situation where the data are uninterpretable. The data will probably remain so because there is no known algorithm which can transform incoherent data into coherent data. In addition, the low level of importance placed upon the concepts of detection limit and analytical blank control as recently as 1975 suggests that the chances are small that these performance criteria would have been determined in a satisfactory manner to have any positive impact on the results reported in this inventory.

D. Precision

Precision is a measure of random error. It is essential for defining significant intra-data differences and is usually expressed as the standard deviation of n analytical measurements. It is possible for an analytical result to be very precise but inaccurate. Precision should be determined for each procedure, type of sample and analyst. It is very important that the estimation of precision is made on "real" samples and not on reference materials. A description of how precision was measured must be provided. The value given for this parameter with a group of data must truly apply to the specific analysis used to produce those data and not simply be a statement of that which has been achieved or that which can be achieved or expected by others for the same or a similar analytical technique. Precision may be estimated by numerous replicated determinations on a sample. This, however, probably leads to a biased estimate since the analyst is apt to give greater care with samples known to be used to measure variance. A better estimate can be made by blindfold determination of several replicated samples (covering the range of concentration expected for the samples) run randomly throughout the period of analysis.

E. Accuracy

Accuracy is a measure of systematic error. As pointed out by Kirchmer (1983), there is a problem in defining accuracy because of the discrepancy between the accuracy of individual analytical measurements and the accuracy of average (long-run) values obtained from a number of replicate measurements (n must be specified). Common practice has been to define accuracy as the deviation of the mean of n measurements from the "true" value; this is the definition used here. Another school of thought applies the term "bias" (equivalent to "systematic error") to the definition above and defines accuracy as the combined random and systematic errors of results. Systematic errors can arise from a variety of sources including:

- unrepresentative sampling;
- instability of samples between sampling and analysis;
- interference effects;
- biased calibration;
- a biased blank correction; and
- inability to determine all forms of the determinand.

It is clear that an accurate result can only occur after each of the above factors is defined quantitatively and precisely.

Accuracy can be estimated by comparing analytical results with the certified values for a reference material. Unfortunately, certified reference materials are available for only a limited number of elements or compounds in the various sample matrices or environmental phases. In practice, this means that the matrix of the reference material and that of the sample will rarely, if ever, be chemically equivalent. Consequently, when an analyst uses a reference material to validate an analytical procedure, the user of the data must judge whether the two matrices (sample and reference material) are sufficiently similar to warrant the use of the results for the reference material as evidence for the validity of the analytical results obtained for actual samples. A simple example illustrates this point: In judging the quality of an apple only another apple, which has been judged and certified to be of specified quality, can be used as a reference. An orange, no matter how good, will not do. The apple may, in fact, be of high quality but this cannot be established on the strength of information obtained using an inappropriate reference material (the orange). In the same way, results for heavy metals in seal livers, for example, cannot be validated using National Bureau of Standards Reference Material

1571 (Orchard Leaves). At the very least, the lipid and oil content of the liver tissue would lead to chemical interferences far greater than those likely to occur with orchard leaves. The degree of interference will be specific to the analytical method; therefore, when assessing the quality of a data set, the experience of the scientist is very important for establishing criteria for accepting/rejecting reference materials in a given situation according to the specific details of each analysis (see Note 8 in the preface to Table 2).

When satisfactory reference materials are not available, accuracy of a method is sometimes estimated by measuring the ability of the method to recover a standard spike added to a portion of the sample or added to a material having a matrix similar to that of the sample. This method may fail to provide a reliable estimate of analytical accuracy because the added standard may not have a chemical reactivity which is equivalent to the component in the sample; it may, therefore, respond differently to the chemical steps involved in the determination. While this method of additions cannot prove that an analytical procedure is accurate it can (by demonstrating poor and highly variable recoveries) identify methods that are inherently imprecise. When certified reference materials or standards are used, standardization should be blindfold and occur at random with replication to avoid biased results caused by analysts who pay special attention to standardization samples and who know or are likely to guess the established reference material composition. In addition, more than one reference material should be used to validate a given method whenever possible. This may identify matrix interferences that go unnoticed using a single matrix. When systematic errors are found (under or over-recovery of the determinand) the results should be reported without correction for recovery data because the application of a correction factor may do little beyond replacing one wrong estimate with another.

Alternatives are available for demonstrating analytical accuracy. Agreement of results determined using different analytical methods employed by different laboratories during intercalibration exercises is one way of increasing confidence in the results. Although reference materials and intercalibrated results may generate confidence in an analytical result, they still can not prove whether the value obtained is an accurate representation of the true value in the environment. Satisfactory agreement among triplicate samples, consistently low blanks and results that make sense (and are consistent with other supporting measurements) will increase confidence.

It is important to distinguish between reference materials and primary standards. The chemical composition of a reference material is empirically derived

from the pooled analytical results of several laboratories using different methods and instruments for the assay. Values for reference materials are commonly described as certified values, recommended values or average compositional values. By contrast, primary standard values are "true" values and are independent of the method of analysis. They are inherently more reliable than the values "recommended" for reference materials. A more detailed discussion of "standard" samples can be found in the review by Abbey (1980).

5.2 Definition of the Rating System

All data have been rated by a 5-level rating system, defined as follows.

Rating Score	Data Quality
0	data are found (or judged) to be wrong or unrepresentative example: seawater samples collected in metal Nansen bottles with metal content as the intended analysis
1	data are suspect because of ill-defined doubts example: magnitude of results falls outside previous scientific experience (see Preface notes to Tables 1 and 2 for further details).
2	insufficient information to assess data; data were not investigated (where data no longer exist, an explanation is entered in remarks column).
3	data are internally consistent; patterns or trends within data are probably real but comparison with other data sets may be a problem (because of incomplete or improper standardization);
4	data are internally consistent and are sufficiently standardized or tied to a reference that comparison with other data at this rating score should be possible. (Data may not be accurate in an absolute sense because all data may be defined relative to a reference material having a different matrix).

This rating system can be criticized for not being hierarchical because the 2 rating score is misplaced. An hierarchical rating scheme can be produced, however, by rearranging the rating scores in the order 2, 0, 1, 3, 4. It is suggested that this re-ordering of data quality levels be used by those more comfortable with an hierarchical scheme rather than creating a new rating system. This will avoid the confusion of several rating schemes within the data inventory series and provide continuity with the inventories already published.

This rating system is intended as a guide and not an absolute statement of data quality; it is one of several ways to represent the quality of acquired chemical data. The ideal rating system would use only objective criteria. This is, however, not possible because of the lack of standard analytical procedures in use and because of the significant changes in sample collection and storage techniques and philosophy that have occurred in recent years. Consequently, a certain subjectivity is inherent in the appraisal of data and any given system for objective data quality appraisal is almost doomed to fail as soon as it is chosen because it will rely to some degree on the discretion of the appraiser. This is particularly true in this inventory of retrospective data quality evaluation, because details of sample history are poorly documented and additional clarification cannot be readily obtained.

Ultimately, the quality of the data will reflect the weakest link in the methodology chain (see 5.3 below). Thus, in cases where a deviation from acceptable methodology is considered so serious that the validity of the obtained results is in doubt, a value of 0 is assigned. Consider the collection of seawater samples for heavy metal analyses, for example. Suppose that the samples were stored acidified in unprepared PVC bottles. During the storage period some metals would leach out of the PVC material which forms the walls of the bottle, so introducing positive contamination to the sample. Because all sampling bottles can differ slightly and sometimes greatly in their composition of impurities, the magnitude of the contamination can be random among samples, so that not even comparison of concentration values within the sample set can be justified in a relative sense. The clear lack of confidence in the results together with the strong suspicion that the samples are no longer representative of the original water mass would result in the assignment of a data rating value of 0. It should be noted that the chemical analyses could have been carried out with well-accepted analytical techniques using the finest state-of-the-art instrumentation. In fact, the analyses could be very precise and very accurate based on the analysis of certified reference materials. Unfortunately, despite the excellence of the analysis the results would still be hopelessly wrong. With reference to this inventory of chemical oceanographic data for the Northwest Passage, cases as obvious as the example above were rare. The most common characteristic of the data sets was an insufficient description of how the results were generated. This led to the assignment of a data rating of 2 to the majority of data sets. It must be stressed, however, that a 2 rating is no better than a 0 rating unless missing information is supplied. If that information no longer exists or in fact never

existed, a 0 rating would be warranted. Those readers interested in data sets rated 2 or below are advised to inspect the data very carefully and investigate the data to a level required by the use to which they intend to put the data.

As already noted, merely stating that a given analytical method was used for a chemical analysis is not sufficient information to reach an opinion about the quality or validity of the data. The quality of analysts varies widely. What can be attained by one analyst can be beyond the abilities of others even when "identical" procedures are employed as demonstrated on more than one occasion during inter-laboratory calibration exercises.

5.3 Significance of the Data Rating Value

The usefulness of the data will depend on the use for which the data are intended, i.e., which question or environmental concern is being considered. The data rating value may be seen to separate groups of data; this can lead to different degrees of understanding environmental processes. It should be recognized that the rating system is not specific to chemical data. Rather, the system seeks to distinguish fundamental differences in information such as those data sets which contain invalid data; those which limit comparisons of numbers to within themselves; and those which allow comparison between data sets.

At least three levels of data quality are essential to establish significance of data:

Level I: Identification of Ranges of Values

At this least discriminating level, the data can be assessed for whether or not their reported ranges fall within the general limits expected for coastal or estuarine areas. Gross errors, contamination, or methodology problems would be identified. Even if the ranges of data were physically possible, this level of data scrutiny does not provide any site-specific information, or determine whether the data were representative of a given geographic area, depth or particular time of year when the data were collected. Most data, even data scoring 1 or 2 could be tentatively used in such a way.

Level 2: Comparison of Data within the Data Sets

At this level, comparison of profiles or time series within a given data set could be used to determine whether measurements of water or sediment properties at particular stations were significantly different from each other on the basis of precision of measurement. Data with a rating of 3 could be used in this instance provided the precision was sufficient to resolve differences within the range of measured values.

Level 3: Comparison of Data Between Data Sets

This is the minimum rating level required for studies of long-term variability of chemical components. It is also required for studies describing processes that control lateral and vertical distributions of chemical components. Studies involving the detection of subtle shifts in chemical equilibria that may lead to downstream effects (such as perturbations to biological systems or climatological changes) also require data with a high level of confidence and a measure of absolute accuracy. Only data with a rating of 4 could be considered for such applications, but will be inadequate when the samples are not representative of the environmental medium sampled. Full interpretation of chemical oceanographic data is impossible in isolation from knowledge of water column structure. Thus, when concurrent measurements of temperature, salinity and perhaps nutrients and dissolved oxygen or some other related variable are not available to support the chemical data, even 4-rated data will be of limited value. In such cases conclusions will be tentative, being based primarily on inference and conjecture.

5.4 Effect of Positioning on the Data Quality

Accuracy of the station positions is a factor essential to the proper use of the data inventory. This is particularly relevant when knowledge of spatial distributions of water or sediment properties are essential for the understanding of a particular oceanographic phenomenon. Many station positions were obtained using rudimentary techniques such as dead reckoning, or radar range and bearing at distances far from shore; consequently, there may be considerable uncertainty about the geographical

location at which samples were obtained. This leads to the dilemma that some chemical data which have received a 4-rating may in fact be of little value in defining important spatial distributions.

Data sets for which no details of positioning or position co-ordinates were given are designated by the term NPG (No Position Given) in the remarks column of Table 2. Entries for these positions appearing in Table 3 were estimated graphically from original report figures and are so indicated and are intended as a very rough guide only.

5.5 Some Important Factors Relevant to the Data Appraisal Process.

The sampling and sample processing techniques used in chemical oceanography are not universally applied to all parameters. Reliable results for certain parameters require the successful application of stringent and highly specialized precautions, while reliable results can be obtained for others using standard routine methodologies. Following is a brief discussion of factors that one must consider when evaluating data and examples of difficulties that can occur during processing of samples for the most commonly observed parameters in Table 2 (heavy metals, nutrients, dissolved oxygen, hydrocarbons, chlorinated hydrocarbons).

5.5.1 Heavy Metals

A. Sea Water

Many of the pitfalls associated with obtaining reliable heavy metal data in sea water can be illustrated by following the history of a water sample from the time of collection through to the completion of the analysis.

The first step facing the chemical oceanographer is the collection of a representative sample. Although this is probably the most important link in the chain of analytical operations, historically it has been given far less thought and care than it deserves. The sampling device must be constructed of materials that will not contaminate the sample. Thus, all metallic components of commonly-used samplers must be removed or replaced. In addition, samplers must be thoroughly cleaned and kept clean between sampling casts. Teflon is an excellent construction material for

samplers because it is usually manufactured with only very low trace metal impurities and can be hot acid cleaned. Samplers made from this material must be carefully cleaned, however, because during the fabrication of Teflon into a chemical apparatus, particles of grit, rust and dirt may become embedded in the surface to act as a source of contamination for long periods of time. The standard Niskin-type water sampler (which has been used extensively in Beaufort Sea sampling) usually contains an internal rubber-coated metal spring or rubber shock cord as part of the closing mechanism. Both are unacceptable since metal impurities can be present in these materials (for example zinc oxide at percent levels). Thus, the use of these samplers for the collection of seawater for zinc analysis is not recommended unless the standard internal spring has been replaced by a Teflon-coated spring or similar contamination-reducing component. Because the sea surface microlayer is enriched with heavy metals, it is advisable to obtain subsurface seawater samples by using samplers such as the Niskin GO-FLO sampler that passes closed through the sea surface layer. Peristaltic pumping systems employing acid-cleaned polyethylene or Teflon tubing have also been used to avoid some of the contamination associated with the surface layer and general handling but these are usually practical only for shallow depths.

The necessity for carefully choosing a sampler is clearly apparent from the results of recent sampler intercomparison studies. Spencer et al., (1982) report that surface water samples taken with a Teflon-coated Niskin GO-FLO sampler possessed much higher concentrations of zinc (7-10 fold) and lead (2-3 fold) than those collected directly in Teflon bottles. Bowers and Windom (1982) compared GO-FLO, Niskin and Hydro-Bios samplers. Their results show that the lower concentration seawater samples (Cd, Cu, Ni, Zn, Fe, Mn, Hg, Mo, and V) were collected using GO-FLO bottles in which O-rings and seals were replaced by silicone equivalents; drain cocks were replaced by those made of solid Teflon. Modified Niskin samplers appeared to be only slightly inferior to the modified GO-FLO, but unmodified GO-FLO and Hydro-Bios samplers were generally poorer.

Other precautions that should be observed during the collection of seawater samples for heavy metal analyses include:

- (i) use of plastic-coated steel rope, Kevlar rope or stainless steel wire rope in place of the standard iron hydrographic wire (Bowers and Windom, 1982);
- (ii) stainless steel weights wrapped in plastic to weigh the hydrographic wire;

- (iii) obtaining surface samples from a small boat by heading into the wind against local surface currents and holding the sample bottle so that it precedes the boat through the water.

Once collected, the sample must be transferred to a storage bottle for some period of time. The storage container must be prepared before use in order that the possibility of the bottle contaminating the sample is reduced. This usually involves a multi-stage and multi-day hot acid-cleaning procedure as described by the Participants in the Lead in Seawater Workshop (1976). Cleaned containers are stored wrapped in polyethylene film and handled only with polyethylene-gloved hands. A preservative is usually added to the samples to inhibit biological activity and the absorption of heavy metals onto the walls of the storage containers. The preservative is most often HCl and must be of sufficient purity to ensure that any trace metal impurities associated with the addition of acid are insignificant relative to the quantity of metal present in the sample.

Most heavy metals samples can be stored in Teflon, polyethylene or quartz when properly cleaned. Mercury should not be stored in polyethylene because mercury vapour readily passes into and through the walls of these containers. The question of whether the sample should be filtered before acidification is controversial. The procedures involved during filtration (particularly on a dirty ship or other field environment where laboratory conditions are not easily reproduced) may often result in greater contamination than would otherwise result from the acidification of unfiltered samples. Sometimes filtration cannot be avoided. On these occasions, filtrations should be carried out (a) under reduced pressure in a closed system apparatus which allows the sample to flow directly from the sample bottle through a pre-cleaned filter into a second pre-cleaned storage bottle or (b) using positive pressure (compressed filtered N₂) in a device that allows direct filtration of a seawater sample from the sampler into a storage bottle. Ideally, the elapsed time between sample collection and analysis should also be minimized to reduce the possibility of sample modification during storage. For instance, prolonged storage may favour the formation of very strong or kinetically-hindered metal complexes with naturally-occurring chelating agents which may, in turn, prevent the formation of an extractable complex with a chelating agent, inhibit a colour forming reaction or impede the reduction of an ion at an electrode. The ultimate result may be that normal analytical methods produce low results or miss a component entirely.

The analytical methodology must also be considered in view of current practices. Any analytical method may be internally consistent yet produce vastly different results from another method, making intercomparison difficult. An example is reported by Brewer and Spencer (1970) where results for the determination of cobalt in sea water obtained from the chelation/extraction/atomic absorption method were five to six fold greater than those obtained by neutron activation of the freeze-dried salts of replicate samples. Even primary reference standards may not be able to resolve such a discrepancy because of matrix interferences, for example. Contamination arising during analysis from atmospheric fallout, reagents, sample handling etc. must also be controlled. The recent trend toward performing heavy metal analyses in laminar flow work stations or Clean Rooms is understandable. Even so, it is not unusual to find members of the most advanced and prestigious marine analytical laboratories in the world disagree with each other by factors of 5 or more on the concentrations of metals in standardized seawater samples during international intercalibration exercises (Sugawara, 1978; Bowers *et al.*, 1981; Olafsson, 1982). Deviations from the standardized or accepted values can exceed $\pm 100\%$ in these intercalibration studies.

The facts presented above emphasize several important points. Steps should always be taken to limit handling and the addition of preservatives to samples. It is also very difficult to form an objective opinion about the quality of a trace metal data set when details such as the ones described above are not given. A true perspective of the heavy metal data as a whole can be achieved by considering the scientists' awareness of problems associated with sampling, storage and sample handling at the time the samples were collected. As the importance of controlling contamination became evident over the years, more effort has been made to systematically eliminate or reduce as many sources as possible. The result has been a continued decrease in the reported values for heavy metals in sea water over the past 40 years (see Table 5.1). It is obvious in many cases (and strongly suspected in others) that although most of the heavy metal data obtained for the Northwest Passage were collected during the past eight years, the techniques used and level of awareness are more characteristic of those used 40 or more years ago. Viewed as a whole, therefore, the quality of heavy metal data for Northwest Passage sea water is probably very low and the data little more than collections of random numbers. During the past several years, artfully and carefully analysed samples for heavy metals have generated results conclusively showing that heavy metals are not

TABLE 3.1
BASELINE CONCENTRATIONS OF SELECTED TRACE METALS IN
OPEN-OCEAN WATERS REPORTED DURING THE LAST FORTY YEARS

REFERENCE	UNITS	Cd	Cu	Pb	Zn	Ni
COMPILED DATA						
Sverdrup <i>et al.</i> , 1942	$\mu\text{mol tonne}^{-1}$	present	157	1.9	85.0	---
Goldberg, 1965	$\mu\text{mol m}^{-3}$	0.98	47	0.14	153.0	---
Brewer, 1975	$\mu\text{mol m}^{-3}$	0.89	7.9	0.14	75.	---
RECENT ORIGINAL DATA						
Zirino and Healy, 1971	$\mu\text{mol m}^{-3}$	—	—	—	26.0	---
Chester and Stoner, 1974	$\mu\text{mol m}^{-3}$	0.62	12.6	—	21.0	---
Eaton, 1976	$\mu\text{mol tonne}^{-1}$	0.53	—	—	—	---
Bender and Gagner, 1976	$\mu\text{mol m}^{-3}$	<0.90	1.9	—	—	---
Windom and Smith, 1979	$\mu\text{mol tonne}^{-1}$	—	0.3 - 5.2	—	—	—
Bruland <i>et al.</i> , 1978	$\mu\text{mol m}^{-3}$	—	—	—	0.15 - 9.2	—
Bruland, 1980	$\mu\text{mol tonne}^{-1}$	0.002 - 1.11	0.54 - 5.3	—	0.08 - 9.1	—
Boyle <i>et al.</i> , 1981	$\mu\text{mol tonne}^{-1}$	<0.01 - 0.08	0.05 - 1.5	—	—	1.03 - 3.5
Danielsson, 1980	$\mu\text{mol tonne}^{-1}$	0.07 - 0.84	1.3 - 5.5	0.10 - 0.82	—	3.1 - 16.2
Kremling and Petersen, 1978	$\mu\text{mol m}^{-3}$	0.03 - 0.04	1.6 - 2.7	—	0.76 - 2.0	—
Patterson, 1974	$\mu\text{mol m}^{-3}$	—	—	0.07	—	—
Heggie, 1982	$\mu\text{mol tonne}^{-1}$	—	2 - 25	—	—	—

erratically distributed in the oceans as was the earlier uncomfortable conclusion drawn by many chemical oceanographers based on data dominated by contamination effects. Rather, horizontal and vertical distributions are now known to vary systematically and can be explained by geological, chemical and physical phenomena. Such evidence of physical and geochemical controls on heavy metal distributions leads to further confidence in metal data and is an aid when evaluating data collections.

B. Sediment and Biological Tissues

In general, the pitfalls encountered in the sampling, storage and analysis of marine sediment samples are fewer than those encountered with seawater samples because the concentrations are commonly about three orders of magnitude higher. Obtaining a representative sample still remains a challenge. Many grab samplers such as the Ponar, screen top Van Veen, Petersen and Kahl box sampler collect substrate together with overlying water. Extreme care must be taken that the overlying water does not wash out the fine material within the surface layer during removal of sediment sample from the sampling device. Before sample collection, it is wise to consider the types of analyses that will be performed on the sample so that a suitable sub-sampling strategy can be devised. Walton (1978), for example, suggests that material from the outer portions of the sample can be used for physical analyses, whereas interior material which is more protected from disturbance or contamination can be used for metal or hydrocarbon analyses. Subsampling for one type of compound should not contaminate the remaining sample before removing a subsequent subsample. Some analysts prefer to analyse a subsample from a homogenized whole sediment; others analyse a given grain size fraction. Because trace metal content usually increases with surface area of particles, intercomparison of many data sets is often tenuous when different size fractions have been analysed. Also, errors may result when sieves made from copper or brass cloth are used to segregate grain sizes. For the analysis of heavy metals in benthic biota it is important that the animals be purged of gut contents before analysis, lest the inorganic sediment present in the gut be included in the estimation of biological metal. For small specimens it is often possible to obtain more reliable results by digesting the whole specimen(s) and subsampling the digest as opposed to subsampling the specimens before the digestion step. For larger animals such as fish and marine

mammals specific organs are usually examined due to the vast range of values possible for various body organs. Special precautions must be taken to check for interferences resulting from the presence of natural oils.

For both sediment and biological metals, methods involving conditions which favour the formation of covalent halides (e.g., hot sulphuric, perchloric or phosphoric acid in the presence of halide ions) should be examined carefully given the possible quantitative distillation of chromium, arsenic, antimony, tin, selenium, rhenium and osmium and the vaporization of substantial amounts of germanium, molybdenum and mercury under such conditions, particularly when Teflon digestion bombs are not employed in the procedure.

5.5.2 Nutrients

The measured concentration of a given nutrient in a sea water sample is very dependent on analytical methodology. The term "reactive nutrients" is often used to emphasize that results refer to those quantities of nutrient that react under the conditions of specific analytical methods. Phosphate is usually measured as soluble inorganic orthophosphate ions which react with acidified molybdate reagent to yield a phosphomolybdate complex which is then reduced to a highly coloured blue compound. Currently used methods for ammonia determination (such as the indophenol blue method) usually measure NH_3 plus NH_4^+ . Earlier methods included varying amounts of labile organic nitrogen compounds such as trimethylamine and amino acids in the determination. Nitrite is determined as an azo dye formed by the stepwise stoichiometric reaction of nitrite ion with an aromatic amine. Nitrate is determined by the same method as nitrite after passing the sample through a catalytic reductor column to reduce nitrate to nitrite. Since many procedures do not measure nitrite separately, the results for nitrate would more accurately be stated as nitrate plus nitrite. The error involved is usually not significant, however, because nitrite is present only at about 5% of the concentration of nitrate. Silicon is determined as dissolved inorganic silicate based on the formation of a yellow silicomolybdic acid when an acidic sample is treated with molybdate solution. Colloidal silicic acid in sea water usually reacts, but polymeric chains containing three or more silicic acid units react very slowly.

The procedures used for sampling and storage of nutrients in sea water are often modified for specific applications. Some general points are noteworthy:

- (a) for best analytical results, samples should be analysed within about one hour of collection;
- (b) short-term storage should be in a cool, dark place;
- (c) if long-term storage by freezing is necessary, it should be limited to a maximum of two months to limit variance caused by storage (Macdonald and McLaughlin, 1982);
- (d) quick-freezing is an effective method for long-term storage of all nutrients except for silicate in estuarine water having a salinity of less than 27×10^{-3} . Caution must also be observed for samples containing silicate in excess of ca 70 mmol m^{-3} or stored for longer than five months. In these cases, data may be successfully recovered provided that thaw times are long enough. The addition of preservatives to samples should be avoided to reduce the chance of contamination and possible interference with the analytical technique. In particular, acidification of phosphate samples is not recommended because of the tendency to favour hydrolysis of combined phosphorus. The addition of chloroform to phosphate samples should also be avoided, because this requires a preliminary sample filtration step which should be avoided if possible. Where water is visibly turbid, PO_4 samples must be filtered. This is particularly important in estuarine samples;
- (e) silicate samples must be stored in plastic; phosphate samples keep best in plastic as uptake of phosphate by glass surfaces has been observed; nitrite/nitrate samples can be effectively stored in glass or plastic;
- (f) before analysis, silicate samples should be thawed for a minimum of three hours to allow for depolymerization (exact time depends on length of time stored in the frozen state).

Although nutrient elements have been analysed frequently by many laboratories for many years, they cannot be routinely determined with sufficient confidence to resolve differences in nutrient water column structure. This is illustrated by the results of a 20-nation nutrient method intercalibration exercise conducted by ICES (International Council for the Exploration of the Sea, 1977) which showed that only 9% of the total variance of laboratory values was attributable to the "within laboratory" component of variance. Thus, the ability of each laboratory

to obtain a precise result was much better than the overall accuracy observed during the intercalibration exercise. Coefficients of variation (CV) for the determination of the different nutrients ranged from 4% to 21%. Because the intercalibration was not a blindfold test, these CV's probably overestimate the precision that would be expected under routine laboratory conditions.

5.5.3 Dissolved Oxygen

Almost without exception, the chemical determination of oxygen in sea water is based on the Winkler titration (Carritt and Carpenter, 1966). Samples must be carefully obtained, fixed and stored all the while limiting or eliminating contact with atmospheric oxygen until after formation of the tri-iodide complex. Numerous systematic errors can occur during the determination of dissolved oxygen by the Winkler titration method most of which result in an overestimation of oxygen content. Utmost care must be taken in the preparation of iodate solution for standardization by using only the highest quality primary standards.

In situ oxygen probes have also been used to measure dissolved oxygen. Because sensor calibrations often wander or because compensation for changes in temperature and salinity are insufficient or too slow, the sensitivity and accuracy of the in situ probe may be insufficient to resolve dissolved oxygen concentrations in the water column or during surface tows. Oxygen concentrations obtained from in situ probes should be treated as approximate unless compelling evidence is provided to indicate otherwise.

5.5.4 Hydrocarbons

A. Sea Water

In the past, the methodologies used by various researchers for sampling, sample preservation and storage, and cleaning procedures have generally been used on an ad hoc basis each designed and implemented to suit a particular study or application. The best samplers appear to be made of glass or stainless steel which can be effectively solvent cleaned. Contamination during sampling is a serious and commonly encountered problem. Hydrowire can be easily contaminated by oil and grease found in plentiful quantities on the sampling vessel and particulate fallout

from combustion of the ship's fuel can also be present. Care should be taken to ensure that all sampling equipment is kept clear of ship's exhaust when travelling between stations. Kevlar or stainless steel cable is preferred for sample casts in conjunction with samplers that pass closed through the sea surface to avoid being contaminated by the natural and anthropogenic hydrocarbon compounds which concentrate in slicks at the surface. This is particularly important for ship-based operations where a halo of surface oil can quickly form at a station and extend a considerable distance away from the vessel. Samples should be stored in the dark and in the cool. They should contain added bactericides such as mercuric chloride, sodium azide, chloroform or methylene chloride to limit alteration of the sample through photolysis or bacterial action. Wong *et al.*, (1976) reported that as much as 30% of dissolved PAH was removed from solution by adsorption onto the walls of the storage container during the sample storage period. It is prudent, therefore, to wash sample containers with solvent to recover analyte that would otherwise be lost and lead to an underestimation of the dissolved PAH content of the samples.

B. Sediments and Biota

Hydrocarbons in sediments and biota present fewer contamination problems than seawater samples because concentrations are very much greater than in sea water. Sediments can be successfully collected with grab samplers with the same reservations as outlined above for sampling sediments for heavy metals. No special precautions beyond those already mentioned above need be taken for biological samples.

During the analysis step, however, several pitfalls may occur. Soxhlet extraction of sediments is common. There is some evidence that this may encourage an *in-situ* contamination through the formation of PAH compounds. Results for low boiling compounds (low molecular weight compounds) should be considered highly unreliable if rotary evaporation techniques have been used to reduce extract volumes.

It is very difficult to compare hydrocarbon results among various studies. Results are often presented for classes of compounds such as aliphatics, polyaromatics, chlorins, fatty acids etc. Column chromatography separations vary from one method to another resulting in the aliphatics determined by one procedure not being equivalent to the aliphatics measured in another. Quantification of specific

compounds may be based on the co-injections of standards in a GC or by reference to an internal standard. Total PAH is defined by some authors as the sum of all resolved peaks, a specifically identified assemblage of PAH or the sum of resolved and unresolved PAH. Different methods generate different looking results. For instance, PAH by fluorescence may generate results in units of a standard pure PAH compound such as chrysene while PAH by GC produces results for specific compounds. An example given by Awad (1981) illustrates that it can be impossible to obtain the same results applying two different techniques on the same sample even if they differ in just a single step. In the two techniques by Blumer *et al.*, (1971) and Vandermeulen *et al.*, (1976) applied to sediments, the extraction steps were different but the purification steps were the same. The weights of organic extracts and yield of hydrocarbons produced from the first technique, however, were three to seven times higher than those obtained from the second.

Intercalibrations using different analytical techniques for hydrocarbons are usually unsuccessful. An example is the analysis of sediment samples from an area affected by the ARGO MERCHANT oil spill using u.v. fluorescence spectroscopy and gas chromatography. Although the samples used were considered to be identical, the two research groups carrying out the analyses reported no significant correlation of results for values less than 100 µg/g. In addition, Zsolnay (1978) reported no significant correlation between the results obtained by GC and high performance liquid chromatography (u.v. detector) when analysing organism tissues.

Extreme caution must be exercised when comparing concentrations for various hydrocarbon groupings or compounds among the various data sets.

5.5.5 Chlorinated Hydrocarbons

The analysis of chlorinated hydrocarbons such as pesticides and polychlorinated biphenyls presents great difficulty, because concentrations of these substances in sea water are very low (parts per 10^9 or less) leading to the need for large sample sizes (10 litres or more). Problems arise from the handling of these large volumes during the complicated multi-stage analytical procedures employed in the determination of chlorinated hydrocarbons. Procedures are often more art than science where the skill of the analyst becomes paramount to the end result. Contamination of samples can be a serious hazard because of the previous widespread use of chlorinated hydrocarbons in the manufacture of industrial and commercial

products. In recent years the manufacture of most chlorinated hydrocarbons and their use in other products has been controlled by legislation, so that the chances of contamination during sampling sea water should be decreasing.

Because chlorinated hydrocarbons are hydrophobic and lipophilic, they tend to concentrate at particle surfaces and in the fatty tissues of organisms. Consequently, their determination in biological tissues and sediments is much simpler than for sea water, although procedures remain complicated and non-routine so that results can be erratic if careful attention is not paid to each step of a procedure.

Regardless of the medium sampled, exact details of sampling, storage and analysis must be specified in order that data quality can be assessed. Quantification is very difficult because of the many isomers and interferences normally encountered in chlorinated hydrocarbon assemblages. Demonstrated effective control of low level blanks can increase confidence in the results.

6. REFERENCES

Data set numbers (where applicable) are given in bold type at the end of the reference.

Abbey, S., 1980. Studies in 'Standard Samples' for use in the general analysis of silicate rocks and minerals. Part 6: 1979 Edition of "Useable" Values. Geological Survey of Canada, Paper 80-14, 30 pp.

Apollonio, S., (no date). Phytoplankton production in the Canadian Arctic. Unpubl. manuscript. The Arctic Institute of North America, 38 pp. + figures. **(56-0002, secondary reference)**

Apollonio, S., 1957. Plankton productivity studies in Allen Bay, Cornwallis Island, N.W.T. 1956. Woods Hole Oceanographic Institution and Osborn Zoological Laboratory, Yale University, 8 pp. + tables. **(56-0002)**

Apollonio, S., 1976. Primary production in the Canadian Arctic Archipelago, 1961-1963. Unpubl. manuscript. Bigelow Laboratory for Ocean Sciences Contrib. No. 76-016. 46 pp. **(56-0002, secondary reference)**

Armstrong, F.A.J. and J.F. Uthe, 1971. Semi-automated determination of mercury in animal tissue. Atomic Absorption Newsletter 10(5): 101-103.

- Atlantic Geoscience Center, 1974. Cruise Report, C.S.S. Hudson September 26 to November 1, 1974. Bedford Institute of Oceanography, 7 pp. + appendices. (74-0013B, secondary reference)
- Atlantic Oceanographic Laboratory, Cruise Report No. 77-0024, 22 August to 17 September, 1977. Bedford Institute of Oceanography. (77-0014)
- Awad, H., 1981. Comparative studies on analytical methods for the assessment of petroleum contamination in the marine environment II. Gas chromatographic analysis. Mar. Chem., 10, 417-430.
- Bain, H., D. Thomson, M. Foy and W. Griffiths, 1977. Marine Ecology of Fast-Ice-Edges in Wellington Channel and Resolute Passage, N.W.T. LGL Ltd. Prepared for Polar Gas Project, 215 pp. + appendices. (76-0010)
- Bedford Institute of Oceanography, 1980. Biological Oceanography Report on C.S.S. Hudson, Cruise 80-027, July 24 to August 29, 1980. (80-0007, secondary reference)
- Bedford Institute of Oceanography, 1976. Cruise report no. 76-025. (76-0011B)
- Bender, M.L. and C. Gagner, 1976. Dissolved copper, nickel and cadmium in the Sargasso Sea. J. Mar. Res., 34, 327-339.
- Bewers, J.M., J. Dalziel, P.A. Yeats and J.L. Barron, 1981. An intercalibration for trace metals in Seawater. Mar. Chem., 10, 173-193.
- Bewers, J.M. and H.L. Windom, 1982. Comparison of sampling devices for trace metal determinations in Sea Water. Mar. Chem., 11, 71-86.
- Blumer, M., G. Souza and J. Sass, 1971. Hydrocarbon pollution of edible shellfish by an oil spill. Mar. Biol., 5, 195-202.
- Birch, J.R., D.B. Fissel, D.D. Lemon, A.B. Cornford, R.A. Lake, B.D. Smiley, R.W. Macdonald and R.H. Herlinveaux, 1983. Arctic Data Compilation and Appraisal. Volume 3. Northwest Passage: Physical Oceanography - Temperature, Salinity, Currents and Water Levels. Can. Data Rep. Hydrogr. Ocean Sci. 5: (Volume 3) 262 pp.
- Bohn, A., 1979. Trace metals in fucoid algae and purple sea urchins near a high Arctic lead/zinc ore deposit. Marine Pollution Bulletin 10: 325-327. (75-0031B)
- Bohn, A. and B.W. Fallis, 1978. Metal concentrations (As, Cd, Cu, Pb and Zn) in shorthorn sculpins, Myoxocephalus scorpius (linnaeus), and arctic char, Salvelinus alpinus, from the vicinity of Strathcona Sound, N.W.T. Water Research, 12, No. 9, 1978. pp. 659-663. (74-0015, secondary reference)
- Bohn, A. and R.O. McElroy, 1976. Trace metals (As, Cd, Cu, Fe and Zn) in arctic cod, Boreogadus saida and selected zooplankton from Strathcona Sound, Northern Baffin Island. J. Fish. Res. Bd. Canada, 33: 2836-2840. (75-0031B)
- Bornhold, B., 1973. Unpublished data on Maxwell Bay and Lancaster Sound. Pacific Geoscience Centre, Sidney, B.C. (73-0009B)

- Bornhold, B., 1974. Unpublished sediment data for Lancaster Sound and Barrow Strait. Pacific Geoscience Centre, Sidney, B.C. (74-0013B)
- Bornhold, B., 1975a. Marine Surficial Geology: Central and Eastern Arctic Project 730031. Geological Survey of Canada, Paper 76-1A: 29-31. (75-0029, secondary reference)
- Bornhold, B., 1975b. Unpublished sediment data on Viscount Melville Sound and Strathcona Sound. Pacific Geoscience Centre, Sidney, B.C. (75-0029)
- Bowes, G.W. and C.J. Jonkel, 1975. Presence and distribution of polychlorinated biphenyls (PCB) in arctic and subarctic marine food chains. J. Fish. Res. Bd. Canada, 32: 2111-2123. (70-0010, 71-0009, 72-0016)
- Boyle, E.A., S.S. Husteded and S.P. Jones, 1981. On the distribution of copper, nickel and cadmium in the surface waters of the North Pacific Ocean. J. Geophys. Res. 86, 8048-8066.
- Brewer, P.G. and D.W. Spencer, 1970. Trace Element Intercalibration Study. Reference No. 70-62, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, U.S.A. (unpublished manuscript), 18 pp.
- Brewer, P.G., 1975. Minor elements in seawater. (In) Chemical Oceanography J.P. Riley and G. Skirrow (Eds.), Vol 1 (second edition), pp. 415-496. Academic Press, New York.
- B.C. Research, 1975. Baseline survey of the marine environment of Strathcona Sound, N.W.T. Report to Strathcona Mineral Services, Project 1552, 84 pp. + appendices. (74-0015)
- Brooks, D.J., 1981. Arctic oceanographic survey, Barrow Strait, 1981 Field Report, Canadian Center for Inland Waters. (unpublished manuscript), 66 pp. (81-0007)
- Bruland, K.W., G.A. Knauer and J.H. Martin, 1978. Zinc in north-east Pacific water. Nature 271, 741-743.
- Bruland, K.W., 1980. Oceanographic distributions of cadmium, zinc, nickel and copper in the North Pacific. Earth and Planetary Science Letters, 47, 176-198.
- Buchanan, R.A., W.E. Cross, D.H. Thomson, 1977. Survey of the marine environment of Bridport Inlet, Melville Island. LGL Ltd. Prepared for Petro-Canada, 265 pp. (77-0015)
- Buchanan, R.A., W.E. Renaud, 1978. The marine ecology of Lancaster Sound and adjacent waters: an annotated bibliography. LGL Ltd. Environmental Studies No. 7 Indian and Northern Affairs, Canada, 262 pp.
- Buckley, D.E., 1963. Recent marine sediments of Lancaster Sound, District of Franklin. M.Sc. Thesis. University of Western Ontario, London, Canada, 112 pp. (60-0008)
- Buckley, D.E., 1971. Recent marine sediments of Lancaster Sound, District of Franklin. Maritime Sediments, 7(3): 66-117. (60-0008, secondary reference)

- Bursa, A.S., 1968. Starch in the oceans. J. Fish. Res. Bd. Canada, 25(6): 1269-1284. (62-0009)
- Campbell, J.A. and D.H. Loring, 1980. Baseline levels of heavy metals in the waters and sediments of Baffin Bay. Marine Pollution Bulletin, 11: 257-261. (77-0014, secondary reference)
- Campbell, J.A. and P.A. Yeats, 1982. The distribution of manganese, iron, nickel, copper and cadmium in the waters of Baffin Bay and the Canadian Archipelago. Oceanologica Acta, 5(2): 161-168. (77-0014)
- Carritt, D.E. and J.H. Carpenter, 1966. Comparison and evaluation of currently employed modifications of the Winkler Method for determining dissolved oxygen in sea water; A NASCO Report. J. Mar. Res. 24(3), 186-318.
- Cheeseman, R.V. and A.L. Wilson, 1978. Manual on Analytical Quality Control for Water Industry. Tech. Rpt. TR66, Water Research Centre, 1978.
- Chemical Oceanography Division, Cruise report no. 76-023, July 23 to August 19, 1976. Bedford Institute of Oceanography. (76-0011A)
- Chemical Oceanography Division, Isotope Chemistry Data Listing, Nutrient Chemistry Data Listing, Organic Chemistry Data Listing, Physical Chemistry Data Listing, Cruise 77-024. Atlantic Oceanographic Laboratory, Bedford Institute of Oceanography. (77-0014)
- Chemical Oceanography Division. Location file cruise 78-026, 1978. Bedford Institute of Oceanography. (78-0020)
- Chester R. and J.H. Stoner, 1974. The distribution of zinc, nickel, manganese, cadmium, copper and iron in some surface waters from the World Ocean. Mar. Chem., 2, 17-32.
- Codispoti, L.A., 1979. Arctic Ocean processes in relation to the dissolved silicon content of the Atlantic. Marine Science Communications. 5(6), 361-381 (1979). (73-0005, secondary reference)
- Codispoti, L.A. and T. G. Owens, 1975. Nutrient transports throughout Lancaster Sound in relation to the Arctic Ocean's reactive silicate budget and the outflow of Bering Strait waters. Dept. of Oceanography, University of Washington, contribution no. 783. (73-0005, secondary reference)
- Collin, A.E., 1958. An oceanographic study of Prince Regent Inlet, the Gulf of Boothia and adjacent waters. Atlantic Oceanographic Group, Fisheries Research Board of Canada, Manuscript Report no. 8, 43 pp. + figures. (56-0003)
- Collin, A.E., 1962a. The oceanography of Lancaster Sound. Ph.D. Thesis. McGill University, 204 pp. (56-0003, 57-0003, secondary reference)
- Collin, A.E., 1962b. The waters of the Canadian Arctic Archipelago. Bedford Institute of Oceanography (unpublished manuscript), 9 pp. (56-0003, secondary reference)

- Collin, A.E., 1962c. The waters of the Canadian Arctic Archipelago. Proceedings of the Arctic Basin Symposium. pp. 128-136. (56-0003, 57-0003, secondary reference)
- Danielsson, L.G., 1980. Cadmium, cobalt, copper, iron, lead, nickel and zinc in Indian Ocean water. *Mar. Chem.* 8, 199-215.
- Dunbar, M. and K.R. Greenaway, 1956. Arctic Canada from the Air. Def. Res. Bd. Can., Queen's Printer, Ottawa, 541 pp. + illustrations.
- Dunbar, M.J. and J.C. Acreman, 1980. Standing crops and species composition of diatoms on sea ice from Robeson Channel to the Gulf of St. Lawrence. *Marine Sciences Center, McGill University, Ophelia* 19(1): 61-72. (74-0023)
- Eaton, A.D., 1976. Marine geochemistry of cadmium. *Mar. Chem.*, 4, 141-154.
- Eaton, R.D.P. and J.P. Farant, 1982. The polar bear as a biological indicator of the environmental mercury burden. *Arctic*, 35(3): 422-425. (80-0018)
- Epstein, S. and T. Mayeda, 1953. Variations of ^{18}O content of waters from natural sources, *Geochim. Cosmochim. Acta*, 4: 213-224.
- Fallis, B.W., 1982. Trace metals in sediments and biota from Strathcona Sound, N.W.T., Nanisivik Marine Monitoring Programme, 1974-1979. Canadian Technical Report of Fisheries and Aquatic Sciences No. 1082, 34 pp. (74-0026, 75-0030)
- Fallis, B., 1975. Unpublished sediment data for Strathcona Sound. (75-0030)
- Glaser, J.A., D.L. Foerst, G.D. McKee, S.A. Quave and W.L. Budde, 1981. Trace analyses for wastewaters. *Env. Sci. Tech.* 15(12), 1426-1435.
- Goldberg, E.D., 1965. Minor elements in sea water. (In) Chemical Oceanography. J.P. Riley and G. Skirrow (Eds). Vol 1 (first edition), pp. 163-196. Academic Press, London.
- Heggie, D.T., 1982. Copper in surface waters of the Bering Sea. *Geochim. Cosmochim Acta* 16, 1301-1306.
- Herlinveaux, R.H., 1970. Ice Pack 8/68 - Oceanographic and biological observations. Fisheries Resources Board of Canada, Technical report no. 159, 60 pp. (68-0001)
- Herlinveaux, R.H. and S.E.G. Wilson, 1974. Oceanographic Observations in the Canadian Arctic, September 1972. Marine Sciences Directorate, Pacific Region, (unpublished manuscript) 11 pp. (77-0011)
- Herlinveaux, R.H., D.B. Fissel and S.E.G. Wilson, 1973. Oceanographic observations in Barrow Strait and Wellington Channel, N.W.T., Institute of Ocean Sciences, Sidney, B.C., 170 pp. (73-0006)
- Hoffman, I., R.J. Westerby and M. Hidioglou, 1968. Precise fluorometric microdeterminations of selenium in agricultural materials. *J. Assoc. Off. Anal. Chem.* 51: 1039-1042.

- International Council for the Exploration of the Sea (ICES), 1977. The international intercalibration exercise for nutrient methods. Koroleff, K., K.H. Palmork, O. Ulltang and J.M. Gieskes (Eds.). ICES, Charlottenlund Slot, DK-2920, Charlottenlund, Denmark, 44 pp.
- Irwin, B., W.G. Harrison, C.L. Gallegos and T. Platt, 1980. Phytoplankton productivity experiments and nutrient measurements in the Labrador Sea, Davis Strait, Baffin Bay and Lancaster Sound from 26 August to 14 September, 1978. Canadian Data Report of Fisheries and Aquatic Sciences No. 213. (78-0020)
- Irwin, B., M. Hodgson, P. Dickie and T. Platt, 1978. Phytoplankton productivity experiments in Baffin Bay and adjacent inlets, 22 August to 18 September, 1977. Marine Ecology Laboratory, Bedford Institute of Oceanography, Fisheries and Marine Service Data Report No. 82, 52 pp. (77-0014)
- Jones, E.P. and A.R. Coote, 1980. Nutrient distributions in the Canadian Archipelago: Indicators of summer water mass and flow characteristics. Canadian Journal of Fisheries and Aquatic Sciences, 37(4): 589-599. (77-0014, secondary reference)
- Kiilerich, A.B., 1939. The Godthaab Expedition, 1928. A theoretical treatment of the hydrographical observation material. Meddeleser om Gronland, Bd. 78. Nr. 5, 146 pp. and plates. (28-0001, secondary reference)
- Kirchmer, C.J., 1983. Quality control in water analyses. Envir. Sci. Tech. 17(4), 174A-181A.
- Kremling, K., and H. Petersen, 1978. The distribution of Mn, Fe, Zn, Cd and Cu in Baltic Sea Water; a study on the basis of one anchor station. Mar. Chem. 6, 155-170.
- Lake, R.A., 1978. An oceanographic study of Bridport Inlet, Melville Island, N.W.T. - Part I, August 1978. Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C., 12 pp. (78-0006)
- Lake, R.A., 1979 a. An oceanographic study of Bridport Inlet, Melville Island, N.W.T. -Part II, March 1979. Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C., 288 pp. (79-0014)
- Lake, R.A., 1979 b. An oceanographic study of Bridport Inlet, Melville Island, N.W.T. - Part III, March to August, 1979. Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C. (unpublished manuscript) 59 pp. (79-0014, secondary reference)
- Lake, R., D. Richards and R. Cooke, 1980. Bridport Inlet, (unpublished O₂ data). Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C. (80-0009)
- Levy, E.M., 1979. Concentration of petroleum residues in the waters and sediments of Baffin Bay and the Eastern Canadian Arctic - 1977. Atlantic Oceanographic Laboratory, Bedford Institute of Oceanography, 34 pp. (77-0014)

- Levy, E.M. and J.D. Moffatt, 1975. Floating petroleum residues in the North Atlantic (1971-1974). Bedford Institute of Oceanography, 56 pp. (72-0017, 73-0009A, 73-0011, 74-0013A, 74-0024)
- Levy, E.M. and A. Walton, 1976. High seas oil pollution: particulate petroleum residues in the North Atlantic. J. Fish. Res. Bd. Canada, 33: 2781-2791. (72-0017, secondary reference)
- Levy, E.M., C.C. Cunningham, C.D.W. Conrad and J.D. Moffatt, 1977. The determination of dissolved oxygen in sea water. Bedford Institute of Oceanography, Report Series/BI-R-77-9, 17 pp. (78-0020, secondary reference)
- Lewis, C.F.M., J.R. Horseman and D.I. Ross. Geological, geophysical and hydrographic studies in Lancaster Sound and Maxwell Bay, District of Franklin. Geological Survey of Canada, Paper 74-1A: 251-252. (73-0009B, secondary reference)
- Long, G.L. and J.D. Winefordner, 1983. Limit of Detection: A closer look at the IUPAC definition. Anal. Chem. 55(7), 712A-724A.
- Macdonald, R.W. and F.A. McLaughlin, 1982. The effect of storage by freezing on dissolved inorganic phosphate, nitrate and reactive silicate for sample from coastal and estuarine waters. Water Research 16, 95-104.
- Marine Ecology Laboratory, Location File cruise 80-027. Chemical Oceanography Division, Bedford Institute of Oceanography, 1982. (80-0007)
- McLaren, P., W.B. Barrie, J.M. Semples, R.A. Sieffert, R.B. Taylor and D. Thomson, 1981. Coastal and environmental data from Eastern Lancaster Sound and Northern Baffin Island, N.W.T. Bedford Institute of Oceanography, Data Series/BI-0-81-1, 283 pp. (78-0021)
- Mirsky, J., 1934. To the Arctic: The Story of Northern Exploration from Earliest Times to the Present, pp. 274-278. University of Chicago Press, Chicago U.S.A. 319 pp.
- Morison, S.R. and R.B. Taylor, 1978. Physical characteristics and seasonal changes in an arctic estuarine environment. Geological Survey of Canada, Paper 78-1B: 101-106 (76-0009)
- Olafsson, J., 1982. An international intercalibration for mercury in sea water. Mar. Chem. 11, 129-142.
- Participants of the Lead in Seawater Workshop, 1976. Comparison determinations of lead by investigators analyzing individual samples of sea water in both their home laboratory and in an isotope dilution standardization laboratory. Mar. Chem., 4, 389-392.
- Patterson, C., 1974. Lead in Sea Water. Science, 183, 553-554.
- Patterson, C.D. Settle and B. Glover, 1976. Analysis of lead in polluted coastal sea water. Mar. Chem., 4, 305-319.

- Renewable Resources Consulting Services Ltd., 1977. Analysis of heavy metals and chlorinated hydrocarbon contamination of five seabirds from Lancaster Sound. 55 pp. (76-0021)
- Richards, F.A. and T.G. Thompson, 1952. The estimation and characterization of plankton populations by pigment analysis-II. A spectrophotometric method for the estimation of plankton pigments. J. Mar. Res. 11(2), 156-172.
- Riis-Carstensen, E., 1936. The Godthaab Expedition, 1928. The hydrographic work and material. Meddeleser om Gronland, Bd. 78. Nr. 3, 101 pp. + plates. (28-0001)
- Sadler, H.E., H.V. Serson and R.K. Chow, 1979. The oceanography of Fury and Hecla Strait. Defence Research Establishment Pacific, Victoria, B.C. Technical Memorandum 79-11, 61 pp. + appendices. (76-0007)
- Sekerak, A.D., R.A. Buchanan, W.B. Griffiths and M.G. Foy, 1976. Biological oceanographic studies in Lancaster Sound 1976. LGL Ltd. Prepared for Norlands Petroleum Ltd., 169 pp. + appendices. (76-0008)
- Sekerak, A.D., D. Thomson, H. Bain and J. Acreman, 1976. Summer surveys of the marine environment of Creswell Bay, Somerset Island and Assistance Bay, Cornwallis Island, N.W.T., 1975, LGL Ltd. Report to Polar Gas Project, 215 pp. (75-0013)
- Sekerak, A.D., R.A. Buchanan, M.G. Foy, H. Bain, G.L. Walder and H.E. Stallard, 1979. Studies of plankton in Northwest Baffin Bay and adjacent waters, July - October, 1978. LGL Ltd., Executive Summary, 412 pp. (78-0022)
- Smith, T.G. and F.H.J. Armstrong. Mercury and selenium in ringed and bearded seal tissues from Arctic Canada. Arctic 31(2): 75-84. (75-0033)
- Spencer, M.J., P.R. Betzer and S.R. Piotrowicz, 1982. Concentrations of cadmium, copper, lead and zinc in surface waters of the Northwest Atlantic Ocean. A comparison of GO-FLO and teflon water samplers. Mar. Chem. 11, 403-410.
- Strickland, J.D.H. and T.R. Parsons, 1968. A practical handbook of seawater analysis. Bull. Fish. Res. Bd. Can., 167: 311 p.
- Strickland, J.D.H. and T.R. Parsons, 1972. A practical handbook of seawater analysis, Second Edition. Bull. Fish. Res. Bd. Can., 167: 310 p.
- Sugawara, K., 1978. Interlaboratory comparison of the determination of mercury and cadmium in sea and fresh waters. Deep-Sea Research, 25, 323-332.
- Sverdrup, H.U., M.W. Johnson and R.H. Fleming, 1942. The Oceans: Their Physics, Chemistry and General Biology, p. 220. Prentice Hall, New Jersey.
- Tan, F.C. and P.M. Strain, 1980. The distribution of sea ice meltwater in the Eastern Canadian Arctic. Journal of Geophysical Research, 85(C4): 1925-1932. (76-0011A)
- Taylor, J.K., 1981. Quality assurance in chemical measurements. Anal. Chem. 53, 1588A-1596A.

- Taylor, J.K., 1983. Validation of analytical methods. *Anal. Chem.* 55(6), 600A-608A.
- Thomson, D., S. Woods and J. Acreman, 1975. Marine ecology survey in the central portion of the Canadian Arctic islands, 1974. Marine Sciences Center, McGill University. Report prepared for Polar Gas Project, 77 pp. (77-0014)
- Top, Z., W.B. Clarke, W.C. Eismout and E.P. Jones, 1980. Radiogenic helium in Baffin Bay bottom water. *Journal of Marine Research*, 38(3): 435-452. (77-0014)
- Vandermeulen, J.H., P.D. Keizer and T.P. Ahren, 1976. Compositional changes in beach sediment-bound Arrow Bunker C: 1970-1976. ICES, Fisheries Improvements Committee, C.M./E.S.I., pp. 1-6.
- Walker, E.R., 1977. Aspects of oceanography in the Archipelago. Institute of Ocean Sciences, Sidney, B.C., 186 pp. (75-0031A)
- Walton, A. (Editor), 1978. Methods for sampling and analysis of marine sediments and dredged materials. Ocean Dumping Report No. 1, Department of Fisheries and Oceans, Ottawa, 74 pp.
- Welch, H.E. and J. Kalff, 1975. Marine metabolism at Resolute Bay, N.W.T. Proceedings of the Circumpolar Conference on Northern Ecology, September 15-18, 1975. Part II 69-75. (71-0016)
- Windom, H.L. and R.G. Smith, Jr., 1979. Copper concentrations in surface waters off the southeastern Atlantic Coast, U.S.A. *Mar. Chem.*, 1, 157-163.
- Wong, C.S., W.J. Cretney, R.W. Macdonald and P. Christensen, 1976. Hydrocarbon levels in the marine environment of the Southern Beaufort Sea. Beaufort Sea Technical Report No. 38. Beaufort Sea Project, Institute of Ocean Sciences, Sidney, B.C., 113 pp.
- Yentsch, C.S. and D.W. Menzel, 1963. A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. *Deep-Sea Res.* 10: 221-231.
- Zirino, A.R. and M.L. Healy, 1971. Voltammetric measurement of zinc in the northeastern tropical Pacific Ocean. *Limnol. Oceanogr.*, 16, 773-778.
- Zsolnay, A., 1978. Lack of correlation between gas-liquid chromatograph and U.V. absorption indicators of petroleum pollution in organisms. *Water, Air and Soil Pollution*, 9, 45-51.

7. DATA INVENTORY TABLE 1.

NOTE:

Entries appearing in column 5 in bold type indicate those chemical quantities reported in Table 2. Entries in regular type refer to quantities reported to have been sampled but for which no data were provided by original authors.

LIST OF ABBREVIATIONS USED IN TABLES 1 and 2

Alk _t	-	total alkalinity
AAS	-	atomic absorption spectrophotometry (this general term used when it is not known whether instrument was used in flame (FAAS) or graphite furnace (GFAAS) mode)
Ag	-	silver (AW = 107.88)
AgDDC	-	silver diethyldithiocarbamate
AOACMA	-	Association of Official Analytical Chemistry Methods of Analysis
AOL	-	Atlantic Oceanography Laboratory (of the Bedford Institute of Oceanography)
APDC	-	ammonium pyrrolidine dithiocarbamate
AS	-	Absorption Spectroscopy
As	-	Arsenic (AW = 74.92)
ATP	-	adenosine triphosphate
AuA	-	auto analyser
B	-	boron (AW = 10.81)
BEC	-	benzene extractable compounds
BIO	-	Bedford Institute of Oceanography
BT	-	bathythermograph
C	-	carbon (AW = 12.01) or conductivity when located in concurrent physical measurements column
$\delta^{13}\text{C}$ (‰)	-	$^{13}\text{C}/^{12}\text{C}$ isotopic ratio
CaCO ₃	-	calcium carbonate
CH ₄	-	methane
CO ₂	-	carbon dioxide
CO ₃	-	carbonate
Ca	-	calcium (AW = 40.08)
CCl ₄	-	carbon tetrachloride
Cd	-	cadmium (AW = 112.40)

CHCl ₃	- chloroform
Chl. <u>a</u>	- chlorophyll <u>a</u>
Chl. <u>b</u>	- chlorophyll <u>b</u>
Chl. <u>c</u>	- chlorophyll <u>c</u>
ΣChl.	- total chlorophyll
Cl	- chlorine (AW = 35.45)
CL	- confidence level
Co	- cobalt (AW = 58.94)
Cr	- chromium (AW = 52.00)
CTD	- conductivity-temperature-depth
CVAAS	- cold vapour atomic absorption spectrophotometry
Cu	- copper (AW = 63.55)
DDDC	- diethylammonium diethyldithiocarbamate
DDE	- bis-(chlorophenyl)-1,1-dichloroethene (F.W. = 318)
dl	- detection limit
DNA	- deoxyribonucleic acid
DWB	- dry weight basis
HCB	- Hexachlorobenzene (F.W. = 284.78)
p,p'-DDD	- 2,2-Bis(p-chlorophenyl)-1,1-dichloroethane (F.W. = 320)
p,p'-DDE	- 2,2-Bis (p-chlorophenyl)-1,1-dichloroethylene (F.W. = 318)
o,p'-DDT	- 1-(o-chlorophenyl)-1-(p-chlorophenyl)-2,2,2-trichloroethane (F.W. = 354.5)
p,p'-DDT	- 1,1-Bis-(p-chlorophenyl)-2,2,2-trichloroethane (F.W. = 354.5)
DOC	- dissolved organic carbon
DOE	- Department of the Environment
ECGC	- electron capture gas chromatography
EDTA	- ethylene diamine tetra acetate
EEP	- electric eye photometer
evap	- evaporation
FAAS	- flame atomic absorption spectrophotometry
Fe	- iron (AW = 55.85)
Fe(OH) ₃	- iron (III) hydroxide
FIDGC	- flame ionization detector gas chromatography
FS	- fluorescence spectroscopy

GC	-	Gas chromatography
GCMS	-	gas chromatography - mass spectrometry
GFAAS	-	Graphite furnace atomic absorption spectrophotometry
GLC	-	gas liquid chromatography
^3H	-	tritium
HC	-	hydrocarbons
HCl	-	hydrochloric acid
HCB	-	hexachlorobenzene
HClO_4	-	perchloric acid
He	-	helium
$\delta^3\text{He}$ (‰)	-	$^3\text{H}/^4\text{He}$ isotopic ratio
HF	-	hydrofluoric acid
Hg	-	mercury (AW = 200.59)
HNO_3	-	nitric acid
H_2SO_4	-	sulphuric acid
HPLC	-	high performance liquid chromatography
HTGC	-	high temperature gas chromatography
HEC	-	hexane extractable compounds
K	-	potassium (AW = 39.10)
K.B.	-	Ken Bruland Sediment Sampler
LWB	-	lipid weight basis
Me-Hg	-	methyl mercury
Mg	-	magnesium (AW = 24.31)
MgCO_3	-	magnesium carbonate
Mn	-	manganese (AW = 54.94)
MS	-	mass spectrometry
MIBK	-	methyl isobutyl ketone
N	-	nitrogen (AW = 14.01)
NaBH_4	-	sodium borohydride
N/A	-	not applicable
NC	-	not computable

Ne	-	neon
NH ₃	-	ammonia
NO ₂	-	nitrite nitrogen
NO ₃	-	nitrate nitrogen
Na	-	sodium (AW = 22.99)
NBS	-	National Bureau of Standards (U.S.A.)
Ni	-	nickel (AW = 58.71)
NS	-	not specified
$\delta^{18}\text{O}$ (‰)	-	$^{18}\text{O}/^{16}\text{O}$ isotopic ratio
O ₂	-	dissolved oxygen
OC	-	organic carbon
ON	-	organic nitrogen
PCB	-	polychlorinated biphenyls
POC	-	particulate organic carbon
POH	-	particulate organic hydrogen
PON	-	particulate organic nitrogen
PE	-	polyethylene
PHAEO	-	phaeopigments
PO ₄	-	phosphate - phosphorus
Pb	-	lead (AW = 207.19)
pH	-	$-\log \text{H}^+$
PVC	-	polyvinyl chloride
P	-	total phosphorus
Rb	-	rubidium (AW = 85.47)
RNA	-	ribonucleic acid
S	-	salinity
SCUBA	-	self-contained underwater breathing apparatus
SD	-	standard deviation
Se	-	selenium (AW = 78.96)
S & P	-	Strickland and Parsons
SF	-	spectrofluorometry
SFC	-	surface

S/S	-	stainless steel
SiO ₃	-	silicate
SMOW	-	standard mean ocean water
SPM	-	suspended particulate matter
SRP	-	soluble reactive phosphorous
T	-	temperature
TC	-	total carbon
TDN	-	total dissolved nitrogen
TDP	-	total dissolved phosphate
TDS	-	total dissolved solids
TOC	-	total organic carbon
TIC	-	total inorganic carbon
TLC	-	thin layer chromatography
TSPM	-	total suspended particulate material
TU	-	tritium units (=1 ³ H/10 ¹⁸ H)
UV	-	ultraviolet
V	-	vanadium (AW = 50.94)
V ₂ O ₅	-	vanadium pentoxide
VS	-	visible spectroscopy
WT	-	Winkler titration
WWB	-	wet weight basis
Zn	-	zinc (AW = 65.37)

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
28-0001	<u>GODTHAAB</u>	Aug. 20	Lancaster Sound	Water Column: <u>O₂</u>	T, S, BT		
50-0002	<u>U.S.S. EDISTO</u>	Aug. 17	Barrow Strait, Lancaster Sound	Water Column: <u>O₂</u>	T, S	Secchi depth	
52-0003	<u>U.S.S. EDISTO</u>	Aug. 8-13	Viscount Melville Sound	Water Column: <u>O₂</u>	T, S	Secchi depth	
53-0001	<u>U.S.S. BURTON ISLAND</u>	Aug. 14-Sept. 1	Prince of Wales Strait	Water Column: <u>O₂, PO₄, SiO₃</u> Bottom Sediment: 72 samples collected (snapper - Phleger)	T, S, BT, currents	Plankton hauls	
54-0002	<u>U.S.S. BURTON ISLAND</u>	Aug. 11-Sept. 6	Viscount Melville Sound, Prince of Wales Strait	Water Column: <u>O₂, PO₄</u>	T, S	Secchi depth	
54-0003	<u>U.S.S. NORTHWIND</u>	Aug. 14-Sept. 6	M'Clure Strait, Prince of Wales Strait, Hadley Bay	Water Column: <u>O₂, PO₄, SiO₃</u>	T, S		

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
56-0002	ARCTIC INSTITUTE OF NORTH AMERICA (Montreal)	July 18-Aug. 28	Allen Bay	Water Column: O ₂ , PO ₄ , P, chl. a	T, S, total radiation	Primary productivity, Secchi depth	
56-0003	H.M.C.S. LABRADOR	Sept.	Gulf of Boothia, Prince Regent Inlet, Lancaster Sound	Water Column: O ₂ , PO ₄	T, S, ice distribution and thickness, sediment constituents and distribution		
57-0003	H.M.C.S. LABRADOR	Aug. 20-Sept. 24	Barrow Strait, Prince Regent Inlet, Lancaster Sound	Water Column: O ₂ , PO ₄	T, S, ice distribution	Secchi depth	
60-0008	C.H.S. BAFFIN (University of Western Ontario)	1960	Lancaster Sound	Bottom Sediment: CO ₂	sediment grain size, hydraulic settling velocity of fine sand, heavy mineral distributions		

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
61-0004	C.C.G.S. LABRADOR	Aug. 28- Sept. 14	Lancaster Sound	Water Column: O ₂	T, S		
62-0005A	FISHERIES RESEARCH BOARD OF CANADA (Arctic unit, Montreal)	June 28- Aug. 14	Allen Bay	Water Column: O ₂	T, S	Plankton hauls	
62-0005B	FISHERIES RESEARCH BOARD OF CANADA (Arctic unit, Montreal)	June 23- Aug. 13	Creswell Bay	Water Column: O ₂	T, S	Plankton hauls	
62-0005C	FISHERIES RESEARCH BOARD OF CANADA (Arctic unit, Montreal)	Sept. 7	Cambridge Bay	Water Column: O ₂	T, S	Plankton hauls	
62-0006	C.C.G.S. JOHN A. MACDONALD	Aug. 7- Sept. 6	Barrow Strait, Franklin Strait, M'Clintock Channel	Water Column: O ₂	T, S		
62-0009	FISHERIES RESEARCH BOARD OF CANADA (Arctic Biological Stn., Quebec)	Aug. 27	Cambridge Bay	Water Column: Starch grains			

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
65-0002	M.V. SALVELINUS	July 29- Sept. 5	Dease Strait	Water Column: O ₂	T, S		
66-0004	U.S.S. EDISTO	Aug. 6- Aug. 7	Lancaster Sound	Water Column: O ₂	T, S		
66-0005	M.V. SALVELINUS	Aug. 2- Aug. 24	Dease Strait	Water Column: O ₂	T, S		
67-0001A	M.V. SALVELINUS	Aug. 17- Sept. 7	Dease Strait	Water Column: O ₂ , PO ₄ , NO ₃	T, S		
67-0001B	ICE	July 2- July 29	Dease Strait	Water Column: O ₂ , PO ₄ , NO ₃	T, S		
68-0001	C.C.G.S. LABRADOR (Fisheries Research Board, Biological Stn., B.C.)	Aug. 24- Aug. 31	Lancaster Sound, Viscount Melville Sound, M'Clure Strait	Water Column: O ₂	T, S, BT	Benthos: species composition, biological bottom samples	
70-0010	CANADIAN WILDLIFE SERVICE	May	Lancaster Sound	Biota: (seals) PCB; p,p'-DDE; p,p'-DDT			see 71-0009 and 72-0016

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
71-0009	CANADIAN WILDLIFE SERVICE	Nov., Dec.	Spence Bay	Biota: (polar bears) PCB, P,p'-DDE			see 70-0010 and 72-0016
71-0016	FISHERIES AND MARINE SERVICE BIOLOGICAL STATION, D.O.E. (St. Andrews, N.B.)	Sept. 1971-Aug. 1972	Resolute Bay	Water Column: chl. a, TDN, SRP, P	Light, ice thickness	Phytoplanktonic and benthic photosynthesis, benthic respiration	see 73-0010
72-0011	C.C.G.S. LOUIS ST. LAURENT (Fisheries and Marine Service, Marine Sciences Directorate, Pacific Region)	Sept. 27-Sept. 30	Barrow Strait	Water Column: O ₂ , Si, PO ₄ , NO ₃ Bottom Sediment: Hg	T, S, BT, currents, light transmission	Zooplankton: Species composition	
72-0016	CANADIAN WILDLIFE SERVICE	June	Resolute Bay	Biota: (polar bears and arctic chars) PCB; P,p'-DDE; P,p'-DDT, o, p'DDT			see 70-0010 and 71-0009
72-0017	C.C.G.S. LABRADOR (Atlantic Oceanographic Laboratory, D.O.E., B.I.O.)	Aug. 20	Barrow Strait	Water Column: tar, HEC, BEC	T		see 73-0009A, 73-0011, 74-0013, 74-0024

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
73-0005	C.C.G.S. LOUIS ST. LAURENT	Aug. 6	Lancaster Sound	Water Column: <u>PO₄, NO₂, NO₃, SiO₃</u>	T, S		
73-0006	INSTITUTE OF OCEAN SCIENCES (Patricia Bay)	Apr. 20-21	Barrow Strait	Water Column: <u>SiO₃, NO₃, PO₄</u>	CTD, T, S, currents		
73-0009A	C.S.S. BAFFIN (Atlantic Oceanographic Laboratory, D.O.E., B.I.O.)	Aug. 7, Sept. 19	Lancaster Sound	Water Column: <u>tar, HEC, BEC</u>	T		see 72-0017, 73-0011, 74-0013A, 74-0024
73-0009B	C.S.S. BAFFIN (Geological Survey)	Aug.-Sept.	Maxwell Bay Lancaster Sound	Bottom Sediment: <u>CaCO₃</u>	sediment types and grain size		
73-0010	FISHERIES AND MARINE SERVICE BIOLOGICAL STN., D.O.E. (St. Andrews, N.B.)	May 8, 13	Resolute Bay	Ice Core: <u>chl. a</u>	sub-ice light	epontic photosynthesis	see 71-0016
73-0011	PROTECTEUR (Atlantic Oceanographic Laboratory, D.O.E., B.I.O.)	Aug. 15, 18	Lancaster Sound	Water Column: <u>Tar, HEC, BEC</u>			see 72-0017, 73-009A, 74-0013A, 74-0024

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
74-0013A	C.S.S. HUDSON (Atlantic Oceanographic Laboratory, D.O.E., B.I.O.)	Aug. 15-30	Lancaster Sound	Water Column: Tar, HEC, BEC	T		see 72-0017, 73-0009A, 73-0011, 74-0024
74-0013B	C.S.S. HUDSON (Geological Survey, Atlantic Geoscience Centre, B.I.O.)	Aug. 20-31	Lancaster Sound, Barrow Strait	Bottom Sediment: CaCO ₃	sediment types and grain size		
74-0013C	C.S.S. HUDSON (Marine Sciences Centre, McGill University)	Aug. 27-Sept. 1	Barrow Strait Peel Sound	Water Column: PO ₄ , NO ₃ , SiO ₃ , chl. a	T, S	Plankton hauls Zooplankton and Benthos: Species composition and numerical abundance	see 74-0014, 74-0014B; research continued in 1975, 1977 (see 75-0013, 77-0015)
74-0014A	C.S.S. HUDSON (Marine Sciences Centre, McGill University)	May 7	Barrow Strait	Water Column: chl. a, PO ₄ , NO ₃ Ice Core: chl. a, chl. b, chl. c, phaeo., PO ₄ , NO ₃	T, S	Phytoplankton: Distribution and concentration Zooplankton and Benthos: Species composition and numerical abundance	see 74-0013C and 74-0014B; research continued in 1975, 1977 (see 75-0013, 77-0015)

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
74-0014B	C.S.S. HUDSON (Marine Sciences Centre, McGill University)	Aug. 14-27	Cunningham Inlet	Water Column: chl. a, PO ₄	T, S	Phytoplankton distribution Zooplankton and Benthos: Species composition and numerical abundance	see 74-0013C, 74-0014A; research continued in 1975, 1977 (see 75-0013, 77-0015)
74-0015	B.C. RESEARCH (for Strathcona Mineral Services Ltd.)	Aug. 8-24	Strathcona Sound	Water Column: O ₂ , pH, Zn, Pb, As, Cd, Cu, Fe, Ni Bottom Sediment: Zn, Pb, As, Cd, Fe, Ni	T, S, C, currents, tides, wind, ice conditions	Seabirds, Fish, Benthos and Plankton: Species composition and numerical abundance	Baseline survey before the operation of a zinc-lead mine. Study includes heavy metal levels in streams and lakes connected to the sound.
				Biota: (Sculpins, arctic char, greenland shark, seaweed, shrimps and plankton) Zn, Pb, As, Cd, Cu, Fe, Ni		Secchi depth	see 75-0031B

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
74-0023	MARINE SCIENCES CENTRE (McGill University)	May	Barrow Strait	Ice Core: NO ₃ , chl. a, phaeo., PO ₄ , SiO ₃	S, ice thickness snow cover	Diatoms: Species composition and numerical abundance	
74-0024	C.S.S. BAFFIN (Atlantic Oceanographic Laboratory, D.O.E., B.I.O.)	Aug. 31, Sept. 15	Barrow Strait, Lancaster Sound	Water Column: Tar, HEC, BEC			see 72-0017, 73-0009A, 73-0011, 74-0013A
74-0026	DEPT. OF FISHERIES AND OCEANS, WESTERN REGION	Sept.	Strathcona Sound	Water Column: Pb, Zn, Cd, Cu Bottom Sediment: Pb, Zn, Cd, As, Hg, Cu, Fe			The study includes the water quality of Twin Lakes Creek. See 75-0030, 76-0012, 79-0024
75-0013	LGL Ltd. (for Polar Gas Project)	Aug. 2-25	Creswell Bay (Somerset Island), Assistance Bay (Cornwallis Island)	Water Column: NO ₃ , P, SiO ₃ , chl. a, phaeo.	T, S, bathymetry	Fish, Plankton, Macrophytes and Benthos: Species composition, numerical abundance and distribution Fish: life history	continuation of the Polar Gas marine research in 1974. See 74-0013C, 74-0014A, 74-0014B, 77-0015

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
75-0029	C.C.G.S. LABRADOR C.C.G.S. D'IBERVILLE (Terrain Sciences Division)	Sept. 4-29	Viscount Melville Sound, Strathcona Sound	Bottom Sediment: CaCO ₃ , Pb, Zn, Cu, Ni, Cd, Fe, As, Co, Mn, Cr, Ag, Hg, ON	Sediment grain size and types, mineral content		
75-0030	DEPT. OF FISHERIES AND OCEANS, WESTERN REGION	Aug.-Sept.	Strathcona Sound	Bottom Sediment: Pb, Zn, Cu, Ni, Cd, Fe, As, Co, Mn, Cr, Hg Biota: (Bivalve molluscs and gastropods) Pb, Zn, Cd, As, Hg, Fe, Cu			The study includes the water quality of Twin Lakes Creek. See 74-0026, 76-0012, 79-0024
75-0031A	B.C. RESEARCH (for Strathcona Mineral Services Ltd.)	March	Strathcona Sound	Water Column: pH	T, S, C, currents		See 74-0015

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
75-0031B	B.C. RESEARCH (for Strathcona Mineral Services Ltd.)	June, Aug.	Strathcona Sound	Water Column: O ₂ , pH, SPM	S		See 74-0015 (Walker, E.R., 1977); study includes heavy metal determinations in nearby lakes and streams
				Biota: (Arctic Cod, zooplankton) As, Cd, Cu, Fe, Zn			(Bohn and McElroy, 1976)
				(furoid algae, sea urchins) As, Cd, Cu, Fe, Zn, Pb			(Bohn, A., 1979)
75-0033	ARCTIC BIOLOGICAL STATION, D.O.E.	1975	Aston Bay (Somerset Island)	Biota: (Ringed seals) Hg, Se			See 76-0022
76-0007	DEFENCE RESEARCH ESTABLISHMENT PACIFIC (Victoria)	May 21	Fury and Hecla Strait	Water Column: O ₂	CTD, tide levels, currents, ice thickness, snow cover		

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
76-0008	LGL LTD., M.V. THETA (for Norlands Petroleum Ltd.)	July 22- Sept. 13	Lancaster Sound	Water Column: O ₂ , chl. a, phaeo., tar	T, S, wave height subsurface illumination	Fish, Phyto- plankton and Zooplankton: species composition and numerical abundance	Three stations (CW, EM, NB) were sampled again for chl. a in 1978; See 78-0022
76-0009	TERRAIN SCIENCES DIVISION	Early July -Aug. 29	Cunningham Inlet	Water Column: SPM	Bathymetry, tides, CTD, ice conditions, sediment composition and distribution		
76-0010	LGL LTD. (for Polar Gas Project)	July 5	Resolute Passage	Water Column: NO ₃ , P, SiO ₂ , chl. a	Ice conditions, currents, T, S	Fish, Phyto- and Zooplankton, Benthos: species composition, numerical abun- dances, distri- bution	
76-0011A	C.S.S. HUDSON (Atlantic Geoscience Centre, R.I.O.)	Aug. 16- 18	Lancaster Sound	Water Column: O ₂ , δ ¹⁸ O	T, S		Nutrients were measured but unreliable due to freezer failure

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
76-0011B	C.S.S. HUDSON (Terrain Sciences Division, Ottawa)	Sept. 1	Strathcona Sound	Bottom Sediment: Pb, Cd, Cu, Ni, Co, Ag, ON, N, OC			
76-0012	DEPT. OF FISHERIES AND OCEANS, WESTERN REGION	May, Aug.	Arctic Bay, Strathcona Sound	Bottom Sediment: Pb, Zn, Cd, Hg, Mn, Cu, Fe Biota: (Seaweeds, sea urchins) Pb, Zn, Cd, As, Se, Hg			The study includes the water quality of Twin Lakes Creek; See 74-0026, 75-0030, 79-0024
76-0021	LGL LTD. (for Norlands Petroleum Ltd.)	July, Aug.	Lancaster Sound	Biota: (Seabirds) As, Cu, Zn, Cd, Cr, V, Hg, Pb, DDE, p,p'-DDT, p,p'-DDD, o,p'-DDT, HCB, PCB			The study was to meet one of the requirements for drilling an offshore exploration well in the Sound
76-0022	ARCTIC BIOLOGICAL STATION, D.O.E.	1976	Barrow Strait	Biota: (Seals) Hg, Me-Hg, Se			See 75-0033

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
77-0014	C.S.S. HUDSON (Chemical Oceanography Division, A.O.L., B.I.O.)	Sept. 8-11	Lancaster Sound	Water Column: CH ₄ , O ₂ , oil, alk., CO ₃ , SiO ₃ , PO ₄ , NO ₃ , NH ₃ POC, POH, PON $\delta^{18}\text{O}$, $\delta^{13}\text{C}$ $\delta^3\text{He}$, He, Ne, $\delta^3\text{H}$ Mn, Fe, Ni, Cu, Cd chl. a Bottom Sediment: oil	T, S, light intensity	primary productivity	(AOL, cruise 77-0024) (AOL, Physical Chemistry Data Listing) (AOL, Nutrient Chemistry Data Listing) (AOL, Organic Chemistry Data Listing) (AOL, Isotope Chemistry Data Listing) (Top et al., 1980) (Campbell and Yeats, 1982) (Irwin et al., 1978) (Levy, 1979)

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
77-0015A	LGL LTD. (for Polar Gas Project)	May 23-27	Brentford Bay	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , chl. <u>a</u> , Σchl.	T, S, ice conditions	Under-ice Flora and Fauna, Phyto- and Zooplankton, Benthos, Marine Fish, Birds and Mammals: 74-0013C, 74-0014A,B, 75-0013	Continuation of the POLAR GAS marine research in 1974 and 1975; See 74-0013C, 74-0014A,B, 75-0013
77-0015B	LGL LTD. (for Polar Gas Project)	Sept. 2	Brentford Bay	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , chl. <u>a</u>	T, S	Phyto- and Zooplankton, Benthos, Marine Fish, Birds and Mammals: 74-0013C, 74-0014 A,B, 75-0013	continuation of the POLAR GAS marine research in 1974 and 1975; See 74-0013C, 74-0014 A,B, 75-0013
77-0016A	LGL LTD. (for Petro-Canada, "The Arctic Pilot Project")	June 9-14	Bridport Inlet	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , O ₂ , chl. <u>a</u>	T, S, ice conditions	Ice Biota, Phyto- and Zooplankton, Benthos and Marine Fish: Species composition and numerical abundance	

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
77-0016B	LGL LTD. (for Petro-Canada, 'The Arctic Pilot Project')	Aug. 5-28	Bridport Inlet	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , O ₂ , chl. a	T, S, currents	Phyto- and Zoo-plankton, Benthos and Marine Fish: Species composition and numerical abundance	
78-0006	FROZEN SEA RESEARCH GROUP, INSTITUTE OF OCEAN SCIENCES, (Sidney, B.C.)	Aug. 22	Bridport Inlet	Water Column: O ₂	CTD, tide gauges		See 79-0014, 80-0009
78-0020	C.S.S. HUDSON (Chemical Oceanography Division, A.O.L., B.I.O.)	Sept. 10, 11	Lancaster Sound	Water Column: O ₂ chl. a, C, N, RNA, DNA, ATP, protein, PO ₄ , NO ₂ , SiO ₃ , NH ₃ , oil, CH ₄	T, S, irradiance	productivity, Secchi	(B.I.O., Chemical Oceanography Division Location File) (Irwin et al., 1980) (oil and methane data were confidential at press time)

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
78-0021	MV GULF STAR (Atlantic Geoscience Centre, DEMR)	Aug. 17- Sept. 21	Lancaster Sound	Water Column: SPM	Bathymetry, sediment grain size and composition	Benthos: Species composition and numerical abundance	Under the Eastern Arctic Marine Environmental Study (EAMES) program, DIAND, See 78-0022
78-0022A	MV THERON MV GULF STAR (LGL Ltd.)	July 28- Aug. 5	Lancaster Sound	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , chl. <u>a</u> , NO ₃	T, S	Plankton and Marine Fish: Species composition, numerical abundance and distribution	Part of the Eastern Arctic Marine Environment Study (EAMES) program, See 78-0021. One station (EM) was sampled for chl. <u>a</u> in 1976 and in the present study, see 76-0008
78-0022B	MV THERON MV GULF STAR (LGL Ltd.)	Aug. 19- Sept. 8	Lancaster Sound	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , chl. <u>a</u> , NO ₃	T, S	Plankton and Marine Fish: Species composition, numerical abundance and distribution	Part of the EAMES study program, see 78-0021. Three stations (CW, EM, NB) were sampled for chl. <u>a</u> in 1976 and in the present study, see 76-0008

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
78-0022C	MV THERON MV GULF STAR (LGL Ltd.)	Sept. 19-22	Lancaster Sound	Water Column: NO ₃ + NO ₂ , PO ₄ , SiO ₃ , chl. <u>a</u> ,	T, S	Plankton and Marine Fish: Species composition, numerical abundance and distribution	Part of the EAMES study program, see 78-0021. Two stations (CW, NB) were sampled for chl. <u>a</u> in 1976 and in the present study, see 76-0008
79-0014	FROZEN SEA RESEARCH GROUP, INSTITUTE OF OCEAN SCIENCES (Sidney, B.C.)	March 13, 20	Bridport Inlet	Water Column: O ₂	CTD, tide gauges, currents		See 78-0006, 80-0009
79-0024	DEPT. OF FISHERIES AND OCEANS, WESTERN REGION	Aug.	Strathcona Sound	Bottom Sediment: Pb, Zn, Cd, As, Hg, Mn, Fe Biota: (Seaweeds, sea urchins, molluscs, sea star, sea cucumber, fish) Pb, Zn, Cd, As, Se, Hg			See 74-0026, 75-0030, 76-0012

DATA SET ID.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
80-0007	C.S.S. HUDSON (Marine Ecology Laboratory, B.I.O.)	Aug. 5-23	Lancaster Sound	Water Column: O ₂ , SiO ₃ , PO ₄ , NO ₃ , SPM	CTD, light, sediment grain size	Primary productivity, seabird survey. Fish: Levels of anti-freeze protein and hepatic mixed function oxidase enzymes Zooplankton: Species composition, numerical abundance, grazing and metabolic rates, digestive enzyme level	
80-0009	FROZEN SEA RESEARCH GROUP, INSTITUTE OF OCEAN SCIENCES (Sidney, B.C.)	April 6-9	Bridport Inlet	Water Column: O ₂	CTD, tide gauges, currents		See 78-0006, 79-0014

DATA SET I.D.	COLLECTING AGENCY, SHIP	DATES OF MEASUREMENT	AREA	CHEMICAL QUANTITIES MEASURED OR SAMPLED	CONCURRENT PHYSICAL MEASUREMENTS	CONCURRENT BIOLOGICAL MEASUREMENTS	REMARKS
80-0018	Medical Services Branch	1980	North Baffin Island, Cornwallis Island	Biota: (polar bear hair) Hg			
81-0007	RESEARCH AND DEVELOPMENT DIVISION, OSS CENTRAL REGION (Burlington, Ontario)	March 31 - April 19	Barrow Strait	Water Column: PO ₄ , NO ₃ + NO ₂ , SiO ₃	CTD, currents, tide gauges	Zooplankton hauls, ice fauna	No chemical values were published; values unreliable due to poor preservation of samples.

8. DATA INVENTORY TABLE 2.

NOTES APPLICABLE TO TABLE 2

The following notes are referred to in the remarks column of Table 2.

NOTE 1.

Delta value

The delta value (in parts per thousand) is defined as:

$$\delta Y = \left(\frac{R_x}{R_{ST}} - 1 \right) \times 1000$$

where Y = ^{13}C , ^{18}O , ^{34}S etc.

R_x = isotopic ratio of unknown (e.g. $^{18}\text{O}/^{16}\text{O}$)

R_{ST} = isotopic ratio of standard

The standard is Chicago PDB for carbon; SMOW (Standard Mean Ocean Water) for oxygen and CDT (Cañon Diablo Troilite) for sulphur.

NOTE 2.

Ratings for dissolved heavy metals and nutrients were downgraded to 0 if values reported for background conditions exceeded the concentrations listed below. These "upper limit values" are based on generally accepted maximum published values for unpolluted coastal areas.

Element	Concentration $\mu\text{mol m}^{-3}$
Mn	180
Fe	180
Zn	76
Cr	19
Cu	79
Pb	5
Cd	9
Ni	85
Co	85
Mo	52
Hg	0.1
As	67
	mmol m^{-3}
Si	70
$\text{NO}_3\text{-N}$	30
$\text{PO}_4\text{-P}$	3.0

A double rating score, (x,0) indicates that some (unspecified) samples may have a rating of 0 while others are of higher data quality.

NOTE 3.

Silicate determined on samples having salinity values less than 27×10^{-3} that have been stored by freezing are prone to large uncertainties (see Section 5.5.3). In the inventory a value of 0 is assigned to those samples for which the above conditions prevail. A double rating of (0, x) indicates some data with a 0 rating (low salinity, stored by freezing) and others with a rating of x.

NOTE 4.

When results are quoted on a WWB and no water content is given, a 4-rating is impossible because of the large variability (perhaps $\pm 50\%$ or more) possible among the samples. Where no water content is given for results when wet weight basis is used, "NOTE 4" is indicated in the remarks column. A maximum rating score of 2 is assigned.

NOTE 5.

Reference materials having certified values for given constituents that are greatly different from values obtained with "unknown" samples may not be good measures of the performance of the analytical method used. As a result, data shown in this inventory were limited to a maximum rating score of 2 when the mean value of the reference material differed from the mean value of the data by an order of magnitude or more. Choice of this criterion is totally subjective.

NOTE 6.

When appropriate reference materials have been used and when these indicate analyte recoveries less than 50% or greater than 150% of the expected (certified) values, a serious problem with the analytical method is suspected. The data are downrated to a maximum score of 1.

NOTE 7.

No estimate of precision or accuracy was made for the actual data in the data set. The report indicates values for accuracy and precision that might be achievable by some under optimum conditions.

NOTE 8.

This note identifies cases where the matrix of the reference material used to validate the method and estimate accuracy is judged to be sufficiently

different from that of the sample so as to make the results obtained for the reference material irrelevant for assessing the quality of the sample data (see additional information, Section 5.1 E). In these cases the maximum rating score is 2. Examples of inappropriate use of reference materials are (1) use of National Bureau of Standards SRM 1577 (Bovine Liver), National Bureau of Standards SRM 1566 (Oyster tissue) or National Bureau of Standards RM 50 (Tuna) to validate seaweed analyses and (2) use of National Bureau of Standards SRM 1571 (Orchard Leaves) to validate seaweed, fish and mammal tissue analyses.

NOTE 9.

All or almost all of the values reported for a data set are less than the detection limit. In such cases, it is impossible to interpret any of the data. Maximum rating is 1 because with additional information some estimation of ranges may be possible. The most probable data rating for these data, however, is zero.

NOTE 10.

This note upgrades the rating score for Bedford Institute of Oceanography nutrient data from 2 to 3. This results from scrutiny of other BIO nutrient data and the authors' familiarity with the procedures, standardizations, analytical details and consistently high performance of the nutrient lab at the Bedford Institute of Oceanography. On the strength of the printed word, only a rating of 2 was given to these data. Other data sets were not investigated to the extent that the BIO nutrient data were investigated because of time and financial limitations. This was totally the judgment of the authors.

NOTE 11.

No mean values are given for pH. This is because pH is a logarithmic function and mean values would be biased to the low end of the pH range and not easily interpreted.

ADDITIONAL COMMENTS ABOUT TABLE 2.

Use of NPG

where NPG (No Position Given) is used in the remarks column it appears only on the first line of the data set but applies to all parameters of the data set unless specifically indicated otherwise.



[illegible]

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT	UNIT	NO. STATIONS	NO. SAMPLES	COLLECTOR	ANALYST	PRECISION	ACCURACY	RANGE	MEAN	MEDIAN	DATA RATING	REMARKS
71 0009														
71 0010	SEA WATER	CHL	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		TIN	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		ALP	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		P	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
72 0011	SEA WATER	CHL	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		TIN	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		ALP	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		P	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
72 0012	ROTTOM SEDIMENT	CHL	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		TIN	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		ALP	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		P	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
72 0013	ROTTOM SEDIMENT	CHL	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		TIN	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		ALP	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100
		P	mg/L	1	1	NO. 100	NO. 100	NO	NO	0.00	0.00	0.00	2	NO. 100

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	METHODS		INFORMATION		MEASURED VALUES			DATA ENTRY	REMARKS
		Qty	Units			Collection	Storage	Analyte	Precision	Accuracy	Range	Mean	Median	
73-0016 (cont'd)		p,p'-DDT	µmol kg ⁻¹	1	1	hoose and line	wrapped in aluminum foil frozen (-20°C)	heptane extraction; ECCG	NS	NS	-	0.034	-	2
		p,p'-DDT	µmol kg ⁻¹	1	1	hoose and line	wrapped in aluminum foil frozen (-20°C)	heptane extraction; ECCG	NS	NS	-	1.77	-	2
		p,p'-DDT	µmol kg ⁻¹	1	1	hoose and line	wrapped in aluminum foil frozen (-20°C)	heptane extraction; ECCG	NS	NS	-	0.028	-	2
		p,p'-DDT	µmol kg ⁻¹	1	1	hoose and line	wrapped in aluminum foil frozen (-20°C)	heptane extraction; ECCG	NS	NS	-	1.42	-	2
73-0017	SEA WATER	Tar	µg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	-	GB < 0.1	-	1
		HEC Tar	µg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	-	GB < 0.1	-	1
		HEC Tar	µg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	-	GB < 0.1	-	1
		HEC Tar	µg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	-	GB < 0.1	-	1
73-0005	SEA WATER	PO ₄	mmol m ⁻³	5	52	Nansen bottles	frozen	Technicon AuA	NS	NS	0.45 - 2.28	1.16	1.24	2
		NO ₂	mmol m ⁻³	5	52	Nansen bottles	frozen	Technicon AuA	NS	NS	0.0 - 0.9	0.30	0.30	2
		NO ₃	mmol m ⁻³	5	52	Nansen bottles	frozen	Technicon AuA	NS	NS	0.0 - 12.9	6.12	8.45	2
		SiO ₃	mmol m ⁻³	5	52	Nansen bottles	frozen	Technicon AuA	NS	NS	0.0 - 26.0	13.48	17.50	2
73-0006	SEA WATER	SiO ₃	mmol m ⁻³	1	9	Fjarie bottles	NS	Strickland and Parsons (1972)	NS	NS	8.0 - 22.0	15.3	19.0	2
		NO ₃	mmol m ⁻³	1	9	Fjarie bottles	NS	Technicon AuA	NS	NS	3.0 - 11.9	7.6	9.6	2
		PO ₄	mmol m ⁻³	1	9	Fjarie bottles	NS	Technicon AuA	NS	NS	0.9 - 2.2	1.5	1.6	2
		Tar	mg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	0.00 - 0.026	0.013	0.013	2
73-0006A	SEA WATER	HEC Tar	mg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	0.00 - 0.019	0.010	0.010	2
		HEC Tar	mg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	0.00 - 0.006	0.003	0.003	2
		HEC Tar	mg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	0.00 - 0.006	0.003	0.003	2
		HEC Tar	mg m ⁻²	2	2	modified Neutron sampler	frozen in glass jars	manual separation; sequential extraction; CCl ₄ + evap/n-hexane + evap/benzene + evap; gravimetry	NS	NS	0.00 - 0.006	0.003	0.003	2
73-0006B	BOTTOM SEDIMENT (surface layer; whole sediment)	CaCO ₃	mol kg ⁻¹	104	104	Powder grab	NS	NS	NS	NS	2.41 - 7.99	4.36	3.96	2

Information obtained solely from unpublished data for stations 73-0006B and 73-0006C.

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS SAMPLED		METHODOLOGY		INFORMATION			MEASURED VALUES			DATA MATING	REMARKS
		Qty	Units	NO.	NO.	Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
73-0010	ICE CORE	chl. a	mg m ⁻²	1	15	15	SCUBA ice core	NS	NS	NS	10.2 - 22.0	15.2	NC	2	data presented as range and mean; NPG
73-0011	SEA WATER	Tar	µg m ⁻²	3	3	0	modified Neutron sampler	frozen in glass jars	NS	NS	-	58 ± 0.1	-	1	HEC + BEC detection limit approx. only; Note 9
		HEC tar	µg m ⁻²	3	3	0	modified Neutron sampler	frozen in glass jars	NS	NS	-	58 ± 0.1	-	1	detection limit approx. only; Note 9
		BEC tar	µg m ⁻²	3	3	0	modified Neutron sampler	frozen in glass jars	NS	NS	-	58 ± 0.1	-	1	detection limit approx. only; Note 9
73-0013A	SEA WATER	Tar	µg m ⁻²	3	3	0	modified Neutron sampler	frozen in glass jars	NS	NS	-	58 ± 0.1	-	1	HEC + BEC detection limit approx. only; Note 9
		HEC tar	µg m ⁻²	3	3	0	modified Neutron sampler	frozen in glass jars	NS	NS	-	58 ± 0.1	-	1	detection limit approx. only; Note 9
		BEC tar	µg m ⁻²	3	3	0	modified Neutron sampler	frozen in glass jars	NS	NS	-	58 ± 0.1	-	1	detection limit approx. only; Note 9
73-0013B	BOTTOM SEDIMENT (whole sediment)	CaCO ₃	mol kg ⁻¹	31	31	31	bottom gravity corer; Van Veen Grab	NS	NS	NS	0.59 - 7.35	3.28	3.83	2	information obtained solely from original measurements
73-0014A	SEA WATER	chl. a	mg m ⁻³	1	3	3	Van Dorn bottles	filtered; frozen	NS	NS	0.07 - 0.11	NC	NC	2	data presented graphically; NPG
		PO ₄	nmol m ⁻³	1	3	3	Van Dorn bottles	filtered; frozen	NS	NS	1.0 - 1.0	NC	NC	2	data presented graphically
		NO ₃	nmol m ⁻³	1	3	3	Van Dorn bottles	filtered; frozen	NS	NS	4.0 - 4.5	NC	NC	2	data presented graphically
		chl. b	mg m ⁻²	1	NS	NS	SIPRE ice corer	melted; filtered; frozen	NS	NS	-	12.35	-	2	data presented as mean values
		chl. b	mg m ⁻²	1	NS	NS	SIPRE ice corer	melted; filtered; frozen	NS	NS	-	2.48	-	2	data presented as mean values
		chl. c	mg m ⁻²	1	NS	NS	SIPRE ice corer	melted; filtered; frozen	NS	NS	-	11.59	-	2	data presented as mean values
		phaeo.	mg m ⁻²	1	NS	NS	SIPRE ice corer	melted; filtered; frozen	NS	NS	-	4.43	-	2	data presented as mean values
		PO ₄	nmol m ⁻³	1	7	7	SIPRE ice corer	melted; filtered; preserved with CHCl ₃ ; frozen	NS	NS	0.091 - 2.1	NC	NC	2	data presented graphically
		NO ₃	nmol m ⁻³	1	7	7	SIPRE ice corer	melted; filtered; preserved with CHCl ₃ ; frozen	NS	NS	0.30 - 1.6	NC	NC	2	data presented graphically

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. SAMPLES W/L	METHODOLOGY		INFORMATION		MEASURED VALUES			DATA BATTING	REMARKS	
		Qty	Units				Collection	Storage	Analyte	Precision	Accuracy	Range	Mean			Median
74-01140	SEA ICE (ice core)	chl. a	mg m ⁻²	11	NS	NS	SIPE ice core	metion (litteral) in plastic bottles after filtration (0.45 µm) stored outside	spectrophotometry (SIPE) (1972)	NS	NS	<1.2 - <2.0	NC	NC	1	data presented graphically; Note 9
		PO ₄	mg m ⁻³	11	NS	NS	SIPE ice core	metion (litteral) in plastic bottles after filtration (0.45 µm) stored outside	Strickland and Parsons (1972)	NS	NS	0.3 - 0.7	NC	NC	2	data presented graphically
74-0115	SEA WATER	O ₂	mol m ⁻³	14	109	109	Kurt Goble water samplers	none; analysis on shipboard	InterOcean System Inc. CSTD unit	NS	NS	0.061 - 0.470	0.246	0.240	2	NPG
		pH	pH units	4	16	16	Kurt Goble water samplers	none; analysis on shipboard	pH meter with combination electrode	NS	NS	7.3 - 8.2	Note 11	7.8	2	
		Zn	nmol m ⁻³	8	31	31	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	APDC extraction; FAAS	NS	NS	0.21 - 0.44	0.32	0.34	0	Note 2
		Pb	nmol m ⁻³	8	31	0	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	APDC extraction; FAAS	NS	NS	-	< 0.02	-	0	Note 2
		As	nmol m ⁻³	8	31	24	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	Fe(OH) ₃ coprecipitation; APDC extraction; colorimetry	NS	NS	<0.01 - 0.04	NC	NC	2	some values expressed as ranges
		Cd	µmol m ⁻³	8	31	0	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	APDC extraction; FAAS	NS	NS	-	< 8.9	-	1	Note 9
		Cu	nmol m ⁻³	8	31	24	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	APDC extraction; FAAS	NS	NS	<0.02 - 0.11	< 0.03	<0.02	2.0	Note 2
		Fe	nmol m ⁻³	8	31	2	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	APDC extraction; FAAS	NS	NS	<0.04 - 0.09	0.04	0.04	1	Note 8 (29 of 31 samples <4.1)
		Ni	nmol m ⁻³	8	31	0	Kurt Goble water samplers	in plastic bottles after filtration (0.45 µm) stored outside	APDC extraction; FAAS	NS	NS	-	< 0.09	-	0	Note 2
		Zn	nmol kg ⁻¹	14	14	14	Shipek sediment dredge	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.06 - 2.42	1.36	1.50	2	DWB
74-0116	BOTTOM (gabbies, organic debris and shells removed)	Pb	nmol kg ⁻¹	14	14	14	Shipek sediment dredge	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.019 - 0.093	0.043	0.038	2	DWB
		As	nmol kg ⁻¹	14	14	14	Shipek sediment dredge	frozen in whirl pak bags	dry ashing; APDC extraction; colorimetry	NS	NS	0.013 - 0.103	0.034	0.033	2	DWB
		Cd	µmol kg ⁻¹	14	14	0	Shipek sediment dredge	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	-	< 3.36	-	1	DWB; Note 9
		Cu	nmol kg ⁻¹	14	14	14	Shipek sediment dredge	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.038 - 0.439	0.319	0.338	2	DWB
		Fe	mol kg ⁻¹	14	14	14	Shipek sediment dredge	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.031 - 0.409	0.439	0.448	2	DWB
		Ni	nmol kg ⁻¹	14	14	14	Shipek sediment dredge	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.044 - 0.465	0.344	0.373	2	DWB

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. SAMPLES W/L	METHODOLOGY		INFORMATION		MEASURED VALUES				DATA DATING	REMARKS
		Qty	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
78-0815 MOTA (cont'd) Mussel Mussel Mussel Mussel Mussel	Zn	mmol kg ⁻¹	1	77	77	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	-1.5%	0.43 - 1.53	0.657	0.628	3	fillet - DWB; accuracy based on bovine liver reference material	
			1	77	77	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	-1.5%	0.73 - 2.34	1.42	1.41	3	fillet - DWB; accuracy based on bovine liver reference material	
			1	77	0	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; GFAAS	NS	-17.6%	<0.015 - 0.022	NC	NC	1	fillet - DWB; d.l. varies with sample weight; accuracy based on bovine liver reference material; Note 9	
			1	77	1	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; GFAAS	NS	-17.6%	-0.014 - <0.020	NC	NC	1	fillet - DWB; d.l. varies with sample weight; accuracy based on bovine liver reference material; Note 9	
			1	14	14	bailed hooks; gillnet	frozen in whirl pak bags	dry ashing; AgEDTC extraction; colorimetry	NS	NS	0.135 - 0.499	0.288	0.282	2	fillet - DWB	
78-0815 MOTA (cont'd) Mussel Mussel Mussel Mussel Mussel	As	mmol kg ⁻¹	1	14	14	bailed hooks; gillnet	frozen in whirl pak bags	dry ashing; AgEDTC extraction; colorimetry	NS	NS	0.190 - 0.865	0.444	0.347	2	fillet - DWB	
			1	77	3	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	<2.8 - <18	NC	NC	1	fillet - DWB; d.l. varies with sample weight; Note 9	
			1	77	71	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	<7.7 - 15%	NC	NC	2	fillet - DWB; d.l. varies with sample weight; Note 9	
			1	77	76	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	-3.42%	<0.020 - 0.156	0.061	0.052	2	fillet - DWB; Note 9; accuracy based on bovine liver reference material	
			1	77	77	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	-3.42%	0.028 - 0.414	0.113	0.083	2	fillet - DWB; Note 9; accuracy based on bovine liver reference material	
78-0815 MOTA (cont'd) Mussel Mussel Mussel Mussel Mussel	Fe	mmol kg ⁻¹	1	77	77	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.473 - 4.210	1.130	0.897	2	fillet - DWB	
			1	77	77	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	0.402 - 5.590	2.08	1.80	2	fillet - DWB	
			1	77	0	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	<0.027 - <0.134	NC	NC	1	fillet - DWB; d.l. varies with sample weight; Note 9	
			1	77	0	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	NS	<0.023 - <0.411	NC	NC	1	fillet - DWB; d.l. varies with sample weight; Note 9	
			1	14	14	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	-1.5%	0.210 - 0.566	0.394	0.322	3	fillet - DWB; accuracy based on bovine liver reference material	
78-0815 MOTA (cont'd) Mussel Mussel Mussel Mussel Mussel	Zn	mmol kg ⁻¹	1	14	14	bailed hooks; gillnet	frozen in whirl pak bags	HNO ₃ digestion; FAAS	NS	-1.5%	0.340 - 2.710	1.82	1.87	3	fillet - DWB; accuracy based on bovine liver reference material	

DATA SET NO.	METAL SAMPLED	MEASUREMENT		NO. STATIONS SAMPLES		NO. NO. SAMPLES	METEOROLOGY		INFORMATION		MEASURED VALUES			DATA RATING	REMARKS	
		Qty	Unit	NO.	NO.		Storage	Collection	Analysis	Precision	Accuracy	Range	Mean			Median
P-0813 (cont'd)	Pb	mixed kg. 1		1	1a	5	gillnet	Frozen in whole job bags	HNC's digestion; GF AAS	NS	+17.4%	+0.010 - -0.019	NC	NC	1	fillet - DWB; d.l. varies with sample weight; accuracy based on borine liver reference material; Note 9
				1	1b	2	gillnet	Frozen in whole job bags	HNC's digestion; GF AAS	NS	+17.4%	+0.017 - -0.182	NC	NC	1	liver - DWB; d.l. varies with sample weight; accuracy based on borine liver reference material; Note 9
				1	1b	3	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	NS	+2.0 - -7.2	NC	NC	1	fillet - DWB; d.l. varies with sample weight; Note 9
				1	1b	4	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	NS	2.4 - -13.6	NC	NC	1	liver - DWB; d.l. varies with sample weight; Note 9
				1	1b	1a	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	-1.62%	0.018 - 0.019	0.030	0.032	3	fillet - DWB; Note 5; accuracy based on borine liver reference material
				1	1b	1a	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	-1.62%	0.48 - 1.48	1.50	1.10	3	liver - DWB; Note 5; accuracy based on borine liver reference material
				1	1b	1a	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	NS	0.288 - 0.614	0.332	0.420	2	fillet - DWB
				1	1b	1a	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	NS	11.44 - 115.93	52.57	50.74	2	liver - DWB
				1	1b	5	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	NS	+0.018 - -0.069	NC	NC	1	fillet - DWB; d.l. varies with sample weight; Note 9
				1	1b	3	gillnet	Frozen in whole job bags	HNC's digestion; FAAS	NS	NS	+0.031 - -0.122	NC	NC	1	liver - DWB; d.l. varies with sample weight; Note 9
(Greenland shark Gadomus nasutus)				1	1a	3	commercial long line fishing equipment	Frozen in whole job bags	HNC's digestion; FAAS	NS	1.5%	-	0.11	-	3	muscle - DWB; accuracy based on borine liver reference material
				1	2	2	commercial long line fishing equipment	Frozen in whole job bags	HNC's digestion; FAAS	NS	1.5%	0.132 - 0.838	0.595	0.595	3	liver + kidney - DWB; accuracy based on borine liver reference material
				1	1	5	commercial long line fishing equipment	Frozen in whole job bags	HNC's digestion; GF AAS	NS	+17.4%	-	+0.014	-	1	muscle - DWB; d.l. varies with sample weight; accuracy based on borine liver reference material; Note 9
				1	2	2	commercial long line fishing equipment	Frozen in whole job bags	HNC's digestion; GF AAS	NS	+17.4%	+0.012 - +0.026	NC	NC	1	liver + kidney - DWB; d.l. varies with sample weight; accuracy based on borine liver reference material; Note 9
				1	1	1	commercial long line fishing equipment	dry ashing; AgITC; colorimetry	NS	NS	-	1.34	-	2	muscle - DWB	
				1	2	2	commercial long line fishing equipment	dry ashing; AgITC; colorimetry	NS	NS	0.410 - 1.310	0.940	0.940	2	liver + kidney - DWB	
				1	1	1	commercial long line fishing equipment	dry ashing; AgITC; colorimetry	NS	NS	-	-	-	2	muscle - DWB	
				1	1	1	commercial long line fishing equipment	dry ashing; AgITC; colorimetry	NS	NS	-	-	-	2	muscle - DWB	
				1	1	1	commercial long line fishing equipment	dry ashing; AgITC; colorimetry	NS	NS	-	-	-	2	muscle - DWB	
				1	1	1	commercial long line fishing equipment	dry ashing; AgITC; colorimetry	NS	NS	-	-	-	2	muscle - DWB	

DATA KEY NO.	MEASUREMENT	QTY	UNITS	NO. STATIONS	NO. SAMPLES	NO. SAMPLING M.L.	METHODOLOGY		INFORMATION			MEASURED VALUES			DATA REMARKS
							Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median	
74-0115 (cont'd)	Cd	mmol kg ⁻¹		1	1	0	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	-	< 0.001	-	1 muscle - DWB; Note 9
	Cu	mmol kg ⁻¹		1	2	2	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.083 - 0.206	0.145	0.145	2 liver + kidney - DWB
	Cu	mmol kg ⁻¹		1	1	1	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	- 3.42%	-	0.035	-	2 muscle - DWB; accuracy based on bovine liver reference material
	Fe	mmol kg ⁻¹		1	2	2	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	- 3.42%	0.080 - 0.194	0.137	0.137	2 liver + kidney - DWB; accuracy based on bovine liver reference material
	Fe	mmol kg ⁻¹		1	1	1	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	-	0.16	-	2 muscle - DWB
	Ni	mmol kg ⁻¹		1	2	2	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.741 - 6.490	3.820	3.820	2 liver + kidney - DWB
	Ni	mmol kg ⁻¹		1	1	1	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	-	0.036	-	2 muscle - DWB
	Zn	mmol kg ⁻¹		1	2	2	commercial long-line fishing equipment	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.020 - 0.046	0.033	0.033	2 liver + kidney - DWB
	Zn	mmol kg ⁻¹		1	6	6	scraper/haul type net dredge	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.761 - 5.910	2.020	1.410	2 DWB
	Pb	mmol kg ⁻¹		1	6	2	scraper/haul type net dredge	frozen in whel	HNO ₃ digestion; GFAAS	NS	NS	< 0.020 - 0.171	0.046	0.020	2 DWB
(see note 9) 74-0116 (cont'd)	As	mmol kg ⁻¹		1	6	6	scraper/haul type net dredge	frozen in whel	dry ashing, AgNOC extraction, colorimetry	NS	NS	0.212 - 0.619	0.183	0.149	2 DWB
	Cd	mmol kg ⁻¹		1	6	6	scraper/haul type net dredge	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0 - 13	7	5	2 DWB
	Cu	mmol kg ⁻¹		1	6	6	scraper/haul type net dredge	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.037 - 0.098	0.083	0.090	2 DWB
	Fe	mmol kg ⁻¹		1	6	6	scraper/haul type net dredge	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	25.8 - 48.9	40.7	45.3	2 DWB
	Ni	mmol kg ⁻¹		1	6	6	scraper/haul type net dredge	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.061 - 0.089	0.073	0.072	2 DWB
	Zn	mmol kg ⁻¹		1	3	3	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.882 - 1.060	0.980	1.001	2 DWB
	Pb	mmol kg ⁻¹		1	3	0	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; GFAAS	NS	NS	< 0.038 - < 0.074	NC	NC	1 DWB; d.l. values with sample weight; Note 9
	As	mmol kg ⁻¹		1	3	3	plankton net; (mesh 28 µm)	frozen in whel	dry ashing, AgNOC extraction, colorimetry	NS	NS	0.325 - 0.273	0.417	0.548	2 DWB
	Cd	mmol kg ⁻¹		1	3	3	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.013 - 0.011	0.025	0.030	2 DWB
	Cu	mmol kg ⁻¹		1	3	3	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	0.713 - 1.000	0.818	0.740	2 DWB
(plant/soil) 74-0117 (cont'd)	Fe	mmol kg ⁻¹		1	3	3	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	4.37 - 8.26	6.91	8.091	2 DWB
	Ni	mmol kg ⁻¹		1	3	2	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	< 0.078 - 0.13	0.099	0.099	2 DWB
	Zn	mmol kg ⁻¹		1	1	1	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; FAAS	NS	NS	-	1.37	-	2 DWB
	Pb	mmol kg ⁻¹		1	1	1	plankton net; (mesh 28 µm)	frozen in whel	HNO ₃ digestion; GFAAS	NS	NS	-	0.30	-	2 DWB

DATA SYNTHESIS NO.	MEDIA SAMPLES	MEASUREMENT		NO. OF STATISTICAL SAMPLES		ANALYSIS	PRECISION	ACCURACY	RANGE	MEASURED VALUES			DATA RATING	REMARKS
		Qty	Units	NO.	NO.	Storage				Mean	Median			
76-0015 (cont'd)		As	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	-	2	DNR	
		Cd	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	0.02	0.02	-	2	DNR	
		Cu	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.13	5.13	-	2	DNR	
		Pb	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.16	5.16	-	2	DNR	
		Se	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	0.11	0.11	-	2	DNR	
76-0023 (cont'd)	SEA WATER (surface)	NO ₃ -N	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		NO ₂ -N	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	0.06	60.31	0.16	2		
		phos.	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	0.02	31.50	0.09	2		
		Tar	total mg/l	2	2	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2	HEC + REC	
		ME ₁ Tar	total mg/l	2	2	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
76-0026 (cont'd)	SEA WATER (surface)	Pb	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		Cd	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		Cu	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		Se	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		NO ₃ -N	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
76-0029 (cont'd)	SEA WATER (surface)	Pb	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		Cd	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		Cu	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		Se	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		
		NO ₃ -N	total mg/l	1	1	freeze-dried, 10% NaCl, glass vials	NS	NS	5.49	0.08	0.08	2		

[illegible]

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS SAMPLED	NO. SAMPLES	METEOROLOGY	MEASURED VALUES					DATA RATING	REMARKS			
		Qty	Unit				Analysis	Precision	Accuracy	Range	Mean			Median		
75-009 (cont'd)	0-100 gms Fractions	10g	gms/10g	15	15	Shore, g-10	NS	NS	NS	1.50	14.96	9.49	2			
	0-2 gms Fractions	2g	gms/2g	22	22	Shore, g-2	NS	NS	NS	9.97	14.40	9.97	2			
	0-100 gms Fractions	10g	gms/10g	22	22	Shore, g-10	NS	NS	NS	0.42	1.33	1.25	2			
	0-100 gms Fractions	10g	gms/10g	22	22	Shore, g-10	NS	NS	NS	0.06	1.90	0.16	2			
75-010	BOTTOM SEDIMENT	10g	gms/10g	18	18	Shore, g-10	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.111	0.048	0.051	2	Information for this data set: This data set is a duplicate of the data set for Station 75-009, but with data from July 1982 and July 1983. The data for the 1982 and 1983 samples are from unpublished reports and are not included in the sediment use (SWB) NPS.
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.222	0.128	0.097	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
	0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2	
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-2 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS	NS	NS	0.000	0.000	0.000	0.000	2		
0-100 gms Fractions	2g	gms/2g	18	18	Shore, g-2	Freeze dried, 100% 100% 100% digestion	NS									

Information for
this table was
obtained partly from
Falls (1982) and
partly from unpub-
lished data for
sediment are (DWB) APC.

Note 9

Note 9

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS SAMPLED		ANALYSIS		MEASURED VALUES			DATE RATING	REMARKS					
		Qty	Units	NO.	NO.	Collection	Storage	Analysis	Precision	Accuracy			Range	Mean	Median		
73-0310	6-180 µm frac (tan)	Co	mmol kg ⁻¹	16	16	16	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	0.19	0.42	0.31	2		
	6-180 µm frac (tan)	Mn	mmol kg ⁻¹	16	16	16	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	3.294	7.280	6.349	2		
	6-2 µm frac (tan)			8	8	8	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	3.276	3.096	6.070	2		
	(white sediment)			8	8	8	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	2.819	6.006	6.186	2		
	6-180 µm frac (tan)	Cr	mmol kg ⁻¹	16	16	16	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	0.15	0.15	0.28	2		
	6-180 µm frac (tan)	Hg	µmol kg ⁻¹	16	16	16	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	0.50	1.00	0.82	2		
	6-2 µm frac (tan)			8	8	8	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	9.97	19.94	12.44	2		
	(white sediment)			8	8	8	Exman drage	Exman in whet sub bags	Freeze-dried, HNO ₃ & H ₂ O ₂ digestion, P.A.A.S.	N/S	N/S	2.49	16.94	6.15	2		
	6-180 µm frac (tan)	Pb	µmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, ashed, filtered, P.A.A.S.	N/S	1.1%	N/C	2.9	N/C	N/C	3	some values expressed as range
	6-2 µm frac (tan)	Zn	mmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, ashed, filtered, P.A.A.S.	N/S	1.1%	N/C	2.82	N/C	N/C	3	some values expressed as range
73-0311A	SEA WATER	Co	µmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, ashed, filtered, P.A.A.S.	N/S	0.0%	N/C	7.7	N/C	N/C	3	some values expressed as range
	SEA WATER	As	mmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, addition of H ₂ O ₂ , (P.A.A.S.)	N/S	27.1%	N/C	0.16	N/C	N/C	3	some values expressed as range
	SEA WATER	Hg	µmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, addition of H ₂ O ₂ , (P.A.A.S.)	N/S	N/S	0.25	0.45	0.37	N/C	2	some values expressed as range
	SEA WATER	Fe	mmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, ashed, filtered, P.A.A.S.	N/S	N/S	N/C	1.7	N/C	N/C	2	some values expressed as range
	SEA WATER	Cr	µmol kg ⁻¹	1	28	28	Exman and Kohler drage	Exman in plastic bags	HNO ₃ digestion, ashed, filtered, P.A.A.S.	N/S	N/S	N/C	0.18	N/C	N/C	3	some values expressed as range
	SEA WATER	pH	pH units	8	N/S	N/S	in situ	N/A	N/A	N/S	N/S	5.8	7.8	Note (1)	N/C	only range given; different methods used for pH; certain values are reference; N/C, reference; N/C,	
	SEA WATER	Cr	mmol kg ⁻¹	8	15	15	N/S	N/S	N/S	N/S	N/S	0.26	0.42	N/C	N/C	2	data presented as range
	SEA WATER	pH	pH units	8	N/S	N/S	N/S	N/S	N/S	N/S	N/S	7.7	8.3	Note (1)	N/C	data presented as range	
	SEA WATER	As	µmol kg ⁻¹	1	N/S	N/S	N/S	N/S	N/S	N/S	N/S	8	15	N/C	N/C	2	data presented as range

[illegible]

DATA SET NO.	MEASUREMENT		NO. STATISTICS SAMPLES			METHODOLOGY		INFORMATION			MEASURED VALUES			DATA BAYING	REMARKS
	Qty	Units	1	2	3	Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
7140318 (Competition, Copepod, whole body, 100% survival, 100% 100% sp.)	As	nmol kg ⁻¹	1	3	3	nylon net, PVC bucket, 5/5 mesh screen	frozen	HNO ₃ digestion, FAAS	NS	-1% - +1%	0.075 - 0.110	0.094	NC	1	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Cd	nmol kg ⁻¹	1	3	3	nylon net, PVC bucket, 5/5 mesh screen	frozen	HNO ₃ digestion, FAAS	NS	NS	0.11 - 0.062	0.031	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Cu	nmol kg ⁻¹	1	3	3	nylon net, PVC bucket, 5/5 mesh screen	frozen	HNO ₃ digestion, FAAS	NS	-3.0%	0.03 - 0.41	0.14	NC	3	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Fe	nmol kg ⁻¹	1	3	3	nylon net, PVC bucket, 5/5 mesh screen	frozen	HNO ₃ digestion, FAAS	NS	-5.0%	0.37 - 1.4	1.1	NC	3	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Zn	nmol kg ⁻¹	1	3	3	nylon net, PVC bucket, 5/5 mesh screen	frozen	HNO ₃ digestion, FAAS	NS	-1.0%	0.44 - 1.40	1.10	NC	3	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	As	nmol kg ⁻¹	5	21	21	non-galvanized steel dredge	frozen in whorl pak bags	HNO ₃ digestion, colorimetry	NS	-13%	0.28 - 0.43	0.34	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
(Purified algae, 100% 100% sp.)	Cd	nmol kg ⁻¹	5	33	33	non-galvanized steel dredge	frozen in whorl pak bags	HNO ₃ digestion, FAAS	NS	NS	6 - 16	9.9	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Cu	nmol kg ⁻¹	5	33	33	non-galvanized steel dredge	frozen in whorl pak bags	HNO ₃ digestion, FAAS	NS	-3% - +3%	0.033 - 0.074	0.047	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Fe	nmol kg ⁻¹	5	33	33	non-galvanized steel dredge	frozen in whorl pak bags	HNO ₃ digestion, FAAS	NS	-3% - +3%	5.04 - 14.5	7.77	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Pb	nmol kg ⁻¹	5	10	10	non-galvanized steel dredge	frozen in whorl pak bags	HNO ₃ digestion, FAAS	NS	-6.7%	2 - 6.8	4	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8
	Zn	nmol kg ⁻¹	5	33	33	non-galvanized steel dredge	frozen in whorl pak bags	HNO ₃ digestion, FAAS	NS	-3% - +3%	0.34 - 2.40	0.90	NC	2	DWB data presented as range and mean; accuracy based on NBS archard material; Note 8

[illegible]

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. ANAL.	METHODOLOGY		INFORMATION		MEASURED VALUES			DATA RATING	REMARKS		
		Qty	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean			Median	
74-0008 (Control)		phen.	mg m ⁻³	7	120	116		Niskin or Nansen bottles	NS	spectrophotometry Strickland and Parsons (1972)	NS	NS (Note 7)	0.00 - 2.44	0.42	0.36	2	
		Tar	mg m ⁻³	7	30	0		Kahl Neutron net (140 µm mesh)	frozen in plastic bottles	benzene extraction; gravimetry	NS	NS	-	< d.l.	-	1	Note 9
74-0009	SEA WATER	SPM	g m ⁻³	8	NS	NS		horizontal alpha bottles	NS	NS	NS	NS	< 1 - 24	NC	NC	2	data presented graphically; NPG
74-0010	SEA WATER	NO ₃	mmol m ⁻³	1	6	6		Niskin bottle	PE bottles; filtered (0.45 µm); frozen	Strickland and Parsons (1972)	NS	NS (Note 7)	6.42 - 9.94	7.96	7.73	2	NPG
		PO ₄	mmol m ⁻³	1	6	6		Niskin bottle	PE bottles; filtered (0.45 µm); frozen	Strickland and Parsons (1972)	NS	NS (Note 7)	0.86 - 1.28	0.99	0.94	2	
		SiO ₃	mmol m ⁻³	1	6	6		Niskin bottle	PE bottles; filtered (0.45 µm); frozen	Strickland and Parsons (1972)	NS	NS (Note 7)	7.09 - 13.89	10.35	10.22	2	
		chl. a	mg m ⁻³	1	6	6		Niskin bottle	PE bottles; filtered (0.45 µm); frozen	spectrophotometry Strickland and Parsons (1972)	NS	NS	0.13 - 0.34	0.24	0.24	2	
		O ₂	mol m ⁻³	5	49	49		Kruken bottles	NS	tightly capped glass bottles	NS	NS	0.237 - 0.394 (18.7) - (-1.8)	0.322	0.318	2	data presented graphically
74-0011B	BOTTOM SEDIMENT ($<180\text{ }\mu\text{m}$ fraction)	Pb	mmol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	0.15 - 0.28	0.20	0.19	2	Information for this data set obtained solely from unpublished original worksheets
		Zn	mmol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	1.29 - 1.79	1.45	1.48	2	
		Cu	mmol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	0.32 - 0.94	0.74	0.76	2	
		Ni	mmol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	0.40 - 0.94	0.80	0.82	2	
		Co	mmol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	0.29 - 0.36	0.31	0.31	2	
		Ag	mmol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	0.011 - 0.014	0.013	0.013	2	
		ON	mol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	0.043 - 0.310	0.16	0.16	2	
		OC	mol kg ⁻¹	9	9	9		Shipek grab	NS	NS	NS	NS	d.l. - 3.3	NC	2.08	2	d.l. not given
74-0012	BOTTOM SEDIMENT ($<180\text{ }\mu\text{m}$ fraction)	Pb	µmol kg ⁻¹	1	1	0		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; MBK extraction; FAAS	NS	NS	-	< 1.2	-	1	Note 9; NPG
		Zn	mmol kg ⁻¹	1	1	1		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; FAAS	NS	NS	-	0.938	-	2	
		Cd	µmol kg ⁻¹	1	1	0		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; FAAS	NS	NS	-	< 2.2	-	1	Note 9
		Hg	µmol kg ⁻¹	1	1	1		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; FAAS	NS	NS	-	0.03	-	2	
		Mn	mmol kg ⁻¹	1	1	1		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; FAAS	NS	NS	-	2.84	-	2	
		Cu	mmol kg ⁻¹	1	1	1		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; FAAS	NS	NS	-	0.529	-	2	
		Fe	mol kg ⁻¹	1	1	1		Elman dredge	frozen in whirl; pak bags	freeze-dried; HNO ₃ /HF/HClO ₄ digestion; FAAS	NS	NS	-	0.355	-	2	
		Pb	µmol kg ⁻¹	2	25	23		Kollwitz dredge and gillnets	frozen in plastic bags	HNO ₃ digestion; freeze-dried; ashed; filtered; benzene extraction; FAAS	NS	+4.4%	NC	8.1	NC	2	some data presented as mean ± SD; accuracy based on NBS archived seawater reference material; Note 8

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS SAMPLED			METHODOLOGY		INFORMATION		MEASURED VALUES			DATA RATING		REMARKS
		Qty	Units	NO.	NO.	NO.	Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
74-0012 (cont'd)	(Sea urchins (whole body analysis) - NBS certificate 74-0012)	Zn	mmol kg ⁻¹	2	25	25	Kolwitt dredge and gillnets	frozen in plastic bags	HNO ₃ digestion; ashed; filtered; FAAS	NS	+4.2%	NC	1.15	NC	3	some data presented as means; DWB; accuracy based on NBS orchard leaves reference material; Note 8
		Cd	µmol kg ⁻¹	2	25	25	Kolwitt dredge and gillnets	frozen in plastic bags	HNO ₃ digestion; ashed; filtered; FAAS	NS	-100%	NS	9.96	NC	1	some data presented as means; DWB; accuracy based on NBS orchard leaves reference material; Note 8
		As	mmol kg ⁻¹	2	25	25	Kolwitt dredge and gillnets	frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; CF AAS	NS	+3.2%	NC	5.37	NC	3	some data presented as means; DWB; accuracy based on NBS orchard leaves reference material; Note 8
		Se	µmol kg ⁻¹	2	25	25	Kolwitt dredge and gillnets	frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; CF AAS	NS	-12.4%	NC	3.2	NC	3	some data presented as means; DWB; accuracy based on NBS orchard leaves reference material; Note 8
		Hg	µmol kg ⁻¹	2	25	25	Kolwitt dredge and gillnets	frozen in plastic bags	addition of V ²⁺ ; HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; CVAAS	NS	NS	NC	5.07	NC	1	some data presented as means; DWB; Note 9
		Pb	µmol kg ⁻¹	2	34	21	Elman and Kolwitt dredge	frozen in plastic bags	HNO ₃ digestion; ashed; filtered; FAAS	NS	NS	NC	6.6	NC	1	some data presented as means; DWB; Note 9
		Zn	mmol kg ⁻¹	2	34	34	Elman and Kolwitt dredge	frozen in plastic bags	HNO ₃ digestion; ashed; filtered; FAAS	NS	NS	NC	5.767	NC	2	some data presented as means; DWB; Note 9
		Cd	µmol kg ⁻¹	2	34	34	Elman and Kolwitt dredge	frozen in plastic bags	HNO ₃ digestion; ashed; filtered; FAAS	NS	NS	NC	10.7	NC	2	some data presented as means; DWB; Note 9
		As	mmol kg ⁻¹	2	34	34	Elman and Kolwitt dredge	frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of V ²⁺ ; NaBH ₄ ; CF AAS	NS	NS	NC	0.047	NC	2	some data presented as means; DWB; Note 9
		Hg	µmol kg ⁻¹	2	34	27	Elman and Kolwitt dredge	frozen in plastic bags	addition of V ²⁺ ; HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; CVAAS	NS	NS	NC	9.1	NC	1	DWB; data presented as means; Note 9
74-0021	(NBSA (Northern Fulmar, Black-legged Kittiwake, Herring Gull, Murre, Dovekie, Black Guillemot)	As	mmol kg ⁻¹	6	50	50	firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	AGDC; extractions; UV spec photometry	NS	NS	0.019 - 0.672	0.171	0.165	2	muscle - DWB; NPC
				6	50	50	firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	AGDC; extractions; UV spec photometry	NS	NS	0.015 - 1.213	0.217	0.109	2	liver - DWB
		Cu	mmol kg ⁻¹	6	50	50	firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.048 - 0.346	0.233	0.233	2	muscle - DWB
				6	50	50	firearms and lead suit	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.137 - 0.343	0.249	0.248	2	liver - DWB

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. SAMPLES g/L	METHODOLOGY		INFORMATION		MEASURED VALUES			DATA BAYING	REMARKS		
		Qty	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean			Median	
76-0021 (cont)	Zn	mmol kg ⁻¹	6	50	50	50	firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.340 -	0.819	0.572	0.575	2	muscle - DWB
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.997 -	2.788	1.458	1.384	2	liver - DWB
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.001 -	0.072	0.015	0.006	2	muscle - DWB
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.005 -	0.417	0.110	0.081	2	liver - DWB
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.003 -	0.115	0.029	0.022	2	muscle - DWB
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.002 -	0.270	0.040	0.020	2	liver - DWB
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	-	-	< d.l.	-	1	muscle - DWB; Note 9; d.l. not specified
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	-	-	< d.l.	-	1	liver - DWB; Note 9; d.l. not specified
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.120 -	3.703	0.800	0.500	2	muscle - DWB (Note 4)
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	H ₂ SO ₄ /HNO ₃ digestion; CVAAS	NS	NS	1.198 -	18.888	4.990	1.700	2	liver - DWB (Note 4)
(total)	Pb	mmol kg ⁻¹	6	50	50	50	firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	HNO ₃ digestion; FAAS	NS	NS	0.013 -	0.370	0.090	0.085	0	log house - DWB; to reduce contamination, perforated aluminum bags were used and tissues were not used in analysis; Pb contamination by authors
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	ECGC; TLC	NS	NS	< d.l.	-	0.039	NC	0.007	1, 2
(total)	DDE	mmol kg ⁻¹	6	50	48	48	firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	ECGC; TLC	NS	NS	-	0.040	NC	0.003	1, 2	LWB, d.l. not specified; correction factors were used and tissues were not used in analysis; Note 9
							firearms and lead shot	wrapped in aluminum foil; frozen; tissues later frozen in glass jars	ECGC; TLC	NS	NS	< d.l.	-	0.040	NC	0.003	1, 2

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. S.D.	METHODOLOGY		INFORMATION			MEASURED VALUES			DATA RATING	REMARKS
		Qty	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
76-0021 (cont'd)		Pd-DDD	mmol kg ⁻¹	6	50	39	forearms and lead shot	wrapped in aluminum foil; frozen tissues later frozen in glass jars	ECGC; TLC	NS	NS	< d.l.	NC	0.007	1, 2	LWB; d.l. not given; standards were used and correction factors applied; will be documented; Note 9
		o,p-DDT	mmol kg ⁻¹	6	50	39	forearms and lead shot	wrapped in aluminum foil; frozen tissues later frozen in glass jars	ECGC; TLC	NS	NS	< d.l.	NC	0.003	1, 2	LWB; d.l. not given; standards were used and correction factors applied; will be documented; Note 9
		HCB	μmol kg ⁻¹	6	50	17	forearms and lead shot	wrapped in aluminum foil; frozen tissues later frozen in glass jars	ECGC; TLC	NS	NS	< d.l.	NC	0.003	1, 2	LWB; d.l. not given; standards were used and correction factors applied; will be documented; Note 9
		PCB	mg kg ⁻¹	6	50	14	forearms and lead shot	wrapped in aluminum foil; frozen tissues later frozen in glass jars	ECGC; TLC	NS	NS	< d.l.	NC	0.002	1, 2	LWB; d.l. not given; standards were used and correction factors applied; will be documented; Note 9
		Hg	μmol kg ⁻¹	2	63	63	hunted	frozen	Armstrong and Linn (1971)	23% - 86%	NS	NS	41.1	NC	2	liver - WWS (Note 9); data presented as mean; NPC
76-0022	NOTA (cont'd)	Me-Hg	μmol kg ⁻¹	1	10	10	hunted	frozen	Armstrong and Linn (1971)	42% - 88%	NS	NS	2.1	NC	2	muscle - WWS (Note 9); data presented as mean
		Se	mmol kg ⁻¹	1	10	10	hunted	frozen	Armstrong and Linn (1971)	48%	NS	NS	4.1	NC	2	liver - WWS (Note 9); data presented as mean
		O ₂	mol m ⁻³	15	163	163	Nitium bottles mounted on rosette	NS	NS	10.9%	NS	0.236 - 0.428	0.331	0.331	3	liver - WWS (Note 9); data presented as mean
		oil	mg m ⁻³	15	157	154	CTD/rosette package with Nitium samplers	Teflon bottles	CCl ₄ extraction followed by PS immediately	NS	NS	< d.l.	NC	0.40	1, 2	d.l. not given; Note 9
		oil	mg m ⁻³	13	13	10	Nevium net	frozen	CCl ₄ extraction; gravimetry	NS	NS	< d.l.	NC	< d.l.	1, 2	d.l. not given; Note 9
77-0010	SEA WATER	oil	mg m ⁻³	2	2	2	S/S screen lowered onto sea surface; recovered water retained on mesh	NS	CCl ₄ extraction; PS	NS	NS	25.2 - 43.5	34.4	34.4	2	liver - WWS (Note 9); data presented as mean
		CH ₄	mmol m ⁻³	15	159	159	glass-lined bottle; sampling	none; analysis on shipboard	CCl ₄ extraction; FID/GC	NS	NS	1.44 - 13.46	3.34	3.05	2	liver - WWS (Note 9); data presented as mean
		alk ₁	equiv m ⁻³	15	159	159	NS	NS	NS	NS	NS	2.0456 - 2.8788	2.2286	2.2265	2	liver - WWS (Note 9); data presented as mean
		CO ₂	equiv m ⁻³	15	159	159	NS	NS	NS	NS	NS	1.8936 - 2.7793	2.1001	2.1187	2	liver - WWS (Note 9); data presented as mean
		SiO ₃	mmol m ⁻³	15	175	175	Nitium bottles mounted on shipboard sampler	none; analysis on shipboard; collection	Technicon AUA	NS	NS (Note 7)	0.94 - 34.96	12.46	13.78	3	Note 10
77-0010	SEA WATER	PO ₄	mmol m ⁻³	15	176	176	Nitium bottles mounted on shipboard sampler	none; analysis on shipboard; collection	Technicon AUA	NS	NS (Note 7)	0.80 - 1.97	1.45	1.52	3	Note 10

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. SAMPLES ANAL.	METHODOLOGY		INFORMATION		MEASURED VALUES				DATA RATING	REMARKS			
		Qty	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median					
77-0018 (Cont'd)		NO ₃	mmol m ⁻³	15	176	146	Niskin bottles on rosette sampler	none, analysis within a few h of collection	Technicon Auto	N5	N5 (N=100.7)	+ d.f.	-	17.26	NC	10.31	3	d.f. not given; Note 10	
		NH ₄	mmol m ⁻³	3	9	9	Niskin bottles on rosette sampler	none, analysis within a few h of collection	wet chemical oxidation; spectrophotometry at 410 nm	N5	N5 (N=100.7)	0.12	-	0.51	0.26	0.17	3	Note 10	
		CHL-4	mg m ⁻³	15	53	53	Niskin bottles on rosette sampler	none, analysis within a few h of collection	PS; Yench and Menzer (1963)	N5	N5	0.10	-	2.92	0.60	0.46	2		
		POC	mg m ⁻³	6	12	12	Niskin bottles	none, analysis on shipboard	N5	N5	N5	N5	7.3	-	191.3	71.7	48.9	2	
		POH	mg m ⁻³	6	12	12	N5	none, analysis on shipboard	N5	N5	N5	N5	2.49	-	16.30	13.83	11.55	2	
		PON	mg m ⁻³	6	12	12	N5	none, analysis on shipboard	N5	N5	N5	N5	0.88	-	7.56	5.16	5.63	2	
		(NO ₂)	μmol	10	95	95	Niskin bottles	tightly capped glass bottles	Egerton and Mayfield (1993)	N5	N5	(-3.02)	-	(0.10)	-1.65	-1.69	2	relative to SMOB	
		13C	‰	1	7	7	Niskin bottles	N5	N5	N5	N5	0.16	-	0.72	0.52	0.51	2	in total dissolved CO ₂	
		34e	‰	5	22	22	Niskin bottles, Geo-Flo samplers	sealed copper tubes; vacuum extraction; 6 months	N5	N5	N5	N5	(-0.3)	-	(18.9)	15.0	13.4	2	standard not specified
		16e	μmol m ⁻³	5	22	22	Niskin bottles, Geo-Flo samplers	sealed copper tubes; vacuum extraction; 6 months	N5	N5	± 0.1%	N5	1.80	-	2.56	1.88	1.87	3	relative to 13C/16C; 1.384 ± 0.4
		16e	μmol m ⁻³	5	21	21	Niskin bottles, Geo-Flo samplers	sealed copper tubes; vacuum extraction; 6 months	N5	N5	± 0.1%	N5	8.01	-	9.09	8.62	8.47	3	
		34	‰	5	29	26	Niskin bottles, Geo-Flo samplers	sealed copper tubes; vacuum extraction; 6 months	N5	N5	± 1% - ± 1%	N5	9.6	-	50.3	17.13	18.20	3	precision varies with concentration
		(total)	μmol m ⁻³	2	10	10	Niskin bottles, Geo-Flo samplers	acidified with Arstar HC 1 to pH - 2; stored in acid-cleaned PE bottles	N5	N5	N5	N5	6.2	-	6.9	5.5	5.4	2	
		(total)	μmol m ⁻³	2	10	10	Niskin bottles, Geo-Flo samplers	acidified with Arstar HC 1 to pH - 2; stored in acid-cleaned PE bottles	N5	N5	N5	N5	23.1	-	62.1	32.0	30.1	2	
		Ni	μmol m ⁻³	2	10	10	Niskin bottles, Geo-Flo samplers	acidified with Arstar HC 1 to pH - 2; stored in acid-cleaned PE bottles	N5	N5	± 1%	N5	3.9	-	5.6	6.7	6.7	3	
		Cu	μmol m ⁻³	2	10	10	Niskin bottles, Geo-Flo samplers	acidified with Arstar HC 1 to pH - 2; stored in acid-cleaned PE bottles	N5	N5	± 1%	N5	3.9	-	5.2	6.7	6.6	3	
	Cd	μmol m ⁻³	2	10	10	Niskin bottles, Geo-Flo samplers	acidified with Arstar HC 1 to pH - 2; stored in acid-cleaned PE bottles	N5	N5	± 1%	N5	0.23	-	0.76	0.52	0.56	3		
	oil	mg kg ⁻¹	9	9	9	Unique grab	freeze-dried; freeze extraction; 6 months	N5	N5	N5	N5	8.75	-	61.23	22.78	22.50	2		
BOTTOM SAMPLES (whole sediment)																			

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS SAMPLED	NO. SAMPLES	NO. VOLS.	METHODOLOGY		INFORMATION		MEASURED VALUES			DATA RATING	REMARKS	
		Qty	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean			Median
77-0013A	SEA WATER	NO ₃ + NO ₂	mmol m ⁻³	1	15	13	Niskin bottles	NS	Technician AuA	NS	NS (Note 7)	7.8	9.5	8.4	2	NPL
		PO ₄	mmol m ⁻³	1	15	15	Niskin bottles	NS	Technician AuA	NS	NS (Note 7)	1.28	1.50	1.39	2	
		SiO ₃	mmol m ⁻³	1	15	15	Niskin bottles	NS	Technician AuA	NS	NS (Note 7)	19.0	22.7	20.6	2	
		chl. a	mg m ⁻³	1	15	14	Niskin bottles	plastic bottles (Note NS)	spectrophotometry; Strickland and Parsons (1972)	NS	NS (Note 7)	-0.01	0.05	0.02	2	
		chl.	mg m ⁻³	1	15	15	Niskin bottles	plastic bottles (Note NS)	spectrophotometry; Strickland and Parsons (1972)	NS	NS	0.02	0.12	0.06	2	
77-0013B	SEA WATER	NO ₃ + NO ₂	mmol m ⁻³	3	8	6	Niskin bottles	NS	Technician AuA	NS	NS (Note 7)	-0.1	7.6	1.8	2	NPL
		PO ₄	mmol m ⁻³	3	8	8	Niskin bottles	NS	Technician AuA	NS	NS (Note 7)	0.40	0.96	0.71	2	
		SiO ₃	mmol m ⁻³	3	8	7	Niskin bottles	NS	Technician AuA	NS	NS (Note 7)	-0.1	7.6	2.7	2	
		chl. a	mg m ⁻³	3	8	8	Niskin bottles	NS	spectrophotometry; Strickland and Parsons (1972)	NS	NS (Note 7)	0.08	0.80	0.22	2	
		NO ₃ + NO ₂	mmol m ⁻³	3	15	15	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technician AuA	NS	NS (Note 7)	0.7	7.6	2.5	2	data presented as range and mean; NPL
77-0014A	SEA WATER	PO ₄	mmol m ⁻³	3	15	15	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technician AuA	NS	NS (Note 7)	0.90	2.00	1.17	2	data presented as range and mean
		SiO ₃	mmol m ⁻³	3	15	15	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technician AuA	NS	NS (Note 7)	6.9	19.0	9.4	2	data presented as range and mean
		O ₂	mol m ⁻³	1	5	5	Niskin and Niskin bottles	none	Heath kit	NS	NS (Note 7)	0.325	0.413	0.376	2	
		chl. a	mg m ⁻³	2	10	10	Niskin bottle	filtered (0.45 µm); MgCl ₂ preserved; frozen	spectrophotometry; Strickland and Parsons (1972)	NS	NS	0.01	0.10	NC	2	data presented graphically
		NO ₃ + NO ₂	mmol m ⁻³	9	46	46	Niskin bottle	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technician AuA	NS	NS (Note 7)	0.0	9.2	0.122	2	data presented as range and mean; NPL
77-0014B	SEA WATER	PO ₄	mmol m ⁻³	9	46	46	Niskin bottle	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technician AuA	NS	NS (Note 7)	0.10	1.40	0.86	2	data presented as range and mean
		SiO ₃	mmol m ⁻³	9	46	46	Niskin bottle	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technician AuA	NS	NS (Note 7)	1.80	27.9	5.36	2	data presented as range and mean
		O ₂	mol m ⁻³	1	9	9	Niskin and Niskin bottles	none	Heath kit	NS	NS (Note 7)	0.219	0.416	0.369	2	
		chl. a	mg m ⁻³	7	28	28	Niskin bottle	filtered (0.45 µm); MgCl ₂ preserved; frozen	spectrophotometry; Strickland and Parsons (1972)	NS	NS	0.08	0.80	NC	2	data presented graphically
		O ₂	mol m ⁻³	1	12	12	NS	bottles	Heath kit	± 0.004	NS	NC	0.433	NC	1	data presented as mean; uncertain per- centage of Heath kit
78-0006	SEA WATER															

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		STATIONS SAMPLED			METHODOLOGY		INFORMATION			MEASURED VALUES			DATA RATING	REMARKS
		Qty	Units	NO.	NO.	NO.	Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
78-0020	SEA WATER	O ₂	mol m ⁻³	4	18	18	SFC; plastic bucket; depth: Niskin bottles	preserved with 1 ml. sodium bicarbonate	modified 0 T	N/S	N/S	0.207 - 0.455	0.308	0.313	2	
		chl. a	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	filtered; frozen on glass fiber in a desiccator (-20°C)	FS	N/S	N/S	0.28 - 5.44	0.29	0.44	2	
		C	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	filtered; freeze-dried; frozen (-20°C)	C/N analyzer	N/S	N/S	27.5 - 102	288	287	2	
		N	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	filtered; freeze-dried; frozen (-20°C)	C/N analyzer	N/S	N/S	48 - 53	51	51	2	
		RNA	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	filtered; frozen (-20°C)	FS	N/S	N/S	22.13 - 22.80	22.15	22.13	2	
		DNA	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	filtered; frozen (-20°C)	FS	N/S	N/S	3.47 - 4.62	4.16	4.38	2	
		ATP	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	three minute extraction in boiling water; frozen (-20°C)	photonometry	N/S	N/S	0.431 - 0.710	0.421	0.452	2	
		protein	mg m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	filtered; frozen (-20°C)	FS	N/S	N/S	135.4 - 198.3	170.8	158.9	2	
		PO ₄	mmol m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	frozen (-20°C)	Technicon Auto	N/S	N/S	0.44 - 0.76	0.70	0.71	3	Note 10
		NO ₃	mmol m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	frozen (-20°C)	Technicon Auto	N/S	N/S	0.40 - 0.46	0.43	0.42	3	Note 10
		SiO ₃	mmol m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	frozen (-20°C)	Technicon Auto	N/S	N/S	2.28 - 2.76	2.46	2.36	3	Note 10
		NH ₃	mmol m ⁻³	1	1	1	SFC; plastic bucket; depth: Niskin bottles	none; analysis was done with photometry at 440 nm	wet chemical oxidation; spectrophotometry at 440 nm	N/S	N/S	0.03 - 0.04	0.04	0.04	3	Note 10
		SPM	g m ⁻³	16	16	16	N/S	N/S	N/S	N/S	N/S	7.0 - 48.3	19.6	19.0	2	NPL
		NO ₃ + NO ₂	mmol m ⁻³	8	72	72	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technicon Auto	N/S	N/S	0.0 - 16.15	N/C	N/C	2	data presented graphically as ranges and mean
78-0021 78-0022A	SEA WATER	NO ₃	mmol m ⁻³	12	32	32	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technicon Auto	N/S	N/S	0.21 - 3.72	0.71	N/C	2	data presented graphically as ranges and mean; different station represented (N/S) for NO ₃
		PO ₄	mmol m ⁻³	17	101	101	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technicon Auto	N/S	N/S	0.50 - 1.44	N/C	N/C	2	data presented graphically as ranges and mean
		SiO ₃	mmol m ⁻³	8	72	72	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	Technicon Auto	N/S	N/S	0.20 - 43.4	N/C	N/C	2	data presented graphically as ranges and mean
		CH ₄	mmol m ⁻³	10	42	42	Niskin bottles	filtered (0.45 µm); preserved with CHCl ₃ ; frozen in PE bottles	FS	N/S	N/S	0.2 - 8.2	N/C	N/C	2	data presented graphically as ranges and mean

DATA SET NO.	MEASUREMENT	NO. STATIONS	NO. SAMPLES	NO. SITES	MEASUREMENT	ANALYSIS	PRECISION	ACCURACY	MEASURED VALUES			REMARKS
									Range	Mean	Median	
78-0026 SEA WATER	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
78-0028 SEA WATER	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
78-0030 SEA WATER	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
78-0032 BOTTOM SAMPLING	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically
	NO ₃ -N	8	78	78	NO ₃ -N	NO ₃ -N	NO ₃ -N	NO ₃ -N	0.0	0.0	0.0	data presented graphically
	NO ₂ -N	8	78	78	NO ₂ -N	NO ₂ -N	NO ₂ -N	NO ₂ -N	0.0	0.0	0.0	data presented graphically

DATA SET NO.	SITES SAMPLED	MEASUREMENT		NO. OF STATIONS SAMPLED			METHODS		INFORMATION			MEASURED VALUES			REMARKS
		Qty	Units	8	67	67	Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median	
79-4829	MUTTA (Seawater) Foca musculosa Alouatta palliata Alouatta palliata Alouatta palliata	Pb	nmol kg ⁻¹	8	67	67	SCUBA	Frozen in plastic bags	HNO ₃ digestion; HClO ₄ digestion; DDDC extraction; FAAS	NS	-2.1% -20.0%	NC	0.031	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Zn	nmol kg ⁻¹	8	67	67	SCUBA	Frozen in plastic bags	HNO ₃ digestion; HClO ₄ digestion; DDDC extraction; FAAS	NS	-2.0% -1.1% -16.2%	NC	1.67	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Cd	pmol kg ⁻¹	8	67	67	SCUBA	Frozen in plastic bags	HNO ₃ digestion; HClO ₄ digestion; DDDC extraction; FAAS	NS	-3.1% 0%	NC	0.42	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		As	nmol kg ⁻¹	8	67	67	SCUBA	Frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /HClO ₄ digestion; addition of NaBH ₄ ; GFAAS	NS	-6.7% -27.1% -10.0%	NC	0.42	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Se	pmol kg ⁻¹	8	67	67	SCUBA	Frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /HClO ₄ digestion; addition of NaBH ₄ ; GFAAS	NS	-10.0% -1.8% -7.8%	NC	1.9	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Hg	pmol kg ⁻¹	8	67	67	SCUBA	Frozen in plastic bags	addition of V ₂ O ₅ ; HNO ₃ /H ₂ SO ₄ /HClO ₄ digestion; CVAAS	NS	NS	NC	0.050	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Pb	nmol kg ⁻¹	5	49	49	SCUBA	Frozen in plastic bags	HNO ₃ digestion; HClO ₄ digestion; DDDC extraction; FAAS	NS	-2.1% -0.2% -28.0%	NC	0.053	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Zn	nmol kg ⁻¹	5	49	49	SCUBA	Frozen in plastic bags	HNO ₃ digestion; HClO ₄ digestion; DDDC extraction; FAAS	NS	-2.0% -1.1% -16.2%	NC	2.19	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Cd	pmol kg ⁻¹	5	49	49	SCUBA	Frozen in plastic bags	HNO ₃ digestion; HClO ₄ digestion; DDDC extraction; FAAS	NS	-3.1% 0%	NC	5.30	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		As	nmol kg ⁻¹	5	49	49	SCUBA	Frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /HClO ₄ digestion; addition of NaBH ₄ ; GFAAS	NS	-6.7% -27.1% -10.0%	NC	0.047	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
	[See section (Whole body analysis) for details of sample collection and analysis]	Se	pmol kg ⁻¹	5	49	49	SCUBA	Frozen in plastic bags	HNO ₃ /H ₂ SO ₄ /HClO ₄ digestion; addition of NaBH ₄ ; GFAAS	NS	-10.0% -1.8% -7.8%	NC	6.3	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8
		Hg	pmol kg ⁻¹	5	49	49	SCUBA	Frozen in plastic bags	addition of V ₂ O ₅ ; HNO ₃ /H ₂ SO ₄ /HClO ₄ digestion; CVAAS	NS	NS	NC	0.15	NC	DWB; data presented as mean values; accuracy based on NBS system, liver and tuna reference materials; Note 8

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATIONS	NO. SAMPLES	NO. SETS	METHODS		INFORMATION		MEASURED VALUES				DATA BATING	REMARKS
		QTY	Units				Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
79-0026 (bottom 5 feet only) (bottom 5 feet only) (bottom 5 feet only) (bottom 5 feet only) (bottom 5 feet only)	Pb	µmol kg ⁻¹	6	6	6	6	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	+2.1% -5.0% -26.0%	N/A	9.46	N/A	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials	
		nmol kg ⁻¹	6	6	6	6	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	+2.0% -1.5% -16.7%	N/A	3.26	N/A	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials	
		nmol kg ⁻¹	6	6	6	6	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	3.1% -5.0%	N/A	0.025	N/A	DIB, data pre-sent as mean values accurate based on NBS on liver reference materials	
		nmol kg ⁻¹	6	6	6	6	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	6.7% -27.5% +10.0%	N/A	0.216	N/A	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials	
		nmol kg ⁻¹	6	6	6	6	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	10.0% -1.8% +7.8%	N/A	0.036	N/A	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials	
	Hg	µmol kg ⁻¹	6	6	6	6	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	N/A	N/A	0.2	N/A	2	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials
(New Year, 1980) (bottom 5 feet only)	Pb	µmol kg ⁻¹	2	2	2	2	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	+2.1% -5.0% -26.0%	N/A	9.79	N/A	0	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials
		nmol kg ⁻¹	2	2	2	2	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	+2.0% -1.5% -16.7%	N/A	0.499	N/A	3	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials
		nmol kg ⁻¹	2	2	2	2	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	3.1% -5.0%	N/A	0.018	N/A	3	DIB, data pre-sent as mean values accurate based on NBS on liver reference materials
		nmol kg ⁻¹	2	2	2	2	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	6.7% -27.5% +10.0%	N/A	0.046	N/A	3	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials
		nmol kg ⁻¹	2	2	2	2	N ₁ (R.A.)	Frogs, dissected in whole (bag)	HNO ₃ digestion, ashd, filtered, P.A.A.N.	N/A	10.0% -1.8% +7.8%	N/A	0.012	N/A	3	DIB, data pre-sent as mean values accurate based on NBS on liver and lung reference materials

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		STATIONS SAMPLED			METHODOLOGY		INFORMATION			MEASURED VALUES			DATA BIASING	REMARKS
		Qty	Units	NO.	NO.	NO.	Collection	Storage	Analysis	Precision	Accuracy	Range	Mean	Median		
79-0209 (cont'd)	Hg	unad kg ⁻¹	2	6	6	6	SCUBA	from, desiccated in whorl bag	addition of V ₂ O ₅ /HNO ₃ /H ₂ SO ₄ digestion; CFAAS	NS	NS	NC	0.40	NC	2	DRE, data presented as mean values
	Pb	unad kg ⁻¹	1	3	3	3	SCUBA	from, desiccated in whorl bag	HNO ₃ digestion; ascorbic acid; DDC extraction; CFAAS	NS	-2.1% 0.0% -28.0%	NC	1.6	NC	3	DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	Zn	mmol kg ⁻¹	1	3	3	3	SCUBA	from, desiccated in whorl bag	HNO ₃ digestion; ascorbic acid; CFAAS	NS	-2.2% -1.1% -16.2%	NC	0.798	NC	3	DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	Cd	mmol kg ⁻¹	1	3	3	3	SCUBA	from, desiccated in whorl bag	HNO ₃ digestion; ascorbic acid; CFAAS	NS	-3.1% 0.0%	NC	0.046	NC	3	DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	As	mmol kg ⁻¹	1	3	3	3	SCUBA	from, desiccated in whorl bag	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; CFAAS	NS	-6.7% -27.1% +10.0%	NC	0.065	NC	3	DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
(Fate of Myxozoon sp. in the digestive tract of the mussel, Mytilus edulis)	Se	mmol kg ⁻¹	1	3	3	3	SCUBA	from, desiccated in whorl bag	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; CFAAS	NS	-10.0% -1.8% -7.8%	NC	0.018	NC	3	DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	Hg	unad kg ⁻¹	1	3	3	3	SCUBA	from, desiccated in whorl bag	addition of V ₂ O ₅ /HNO ₃ /H ₂ SO ₄ digestion; CFAAS	NS	NS	NC	0.2	NC	2	DRE, data presented as mean values
	Pb	unad kg ⁻¹	6	17	17	17	SCUBA	from, desiccated in whorl bag	HNO ₃ digestion; ascorbic acid; DDC extraction; CFAAS	NS	-2.1% 0.0% -28.0%	NC	2.1	NC	3	mean - DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	Zn	mmol kg ⁻¹	6	18	18	18	SCUBA	from, desiccated in whorl bag	HNO ₃ digestion; ascorbic acid; CFAAS	NS	-2.2% -1.1% -16.2%	NC	1.11	NC	3	mean - DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	Cd	unad kg ⁻¹	6	18	18	18	SCUBA	from, desiccated in whorl bag	HNO ₃ digestion; ascorbic acid; CFAAS	NS	-3.1% 0.0%	NC	2.0	NC	3	mean - DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials
	As	mmol kg ⁻¹	6	18	18	18	SCUBA	from, desiccated in whorl bag	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; CFAAS	NS	-6.7% -27.1% +10.0%	NC	0.188	NC	3	mean - DRE, data presented as mean values; accuracies based on NBS system, liver and lung reference materials

DATA SET NO.	MEDIUM SAMPLED	MEASUREMENT		NO. STATION SAMPLES	NO. ANALYSES	METHODOLOGY		INFORMATION		MEASURED VALUES			DATA BIASING	REMARKS		
		Qty	Units			Collection	Storage	Analysis	Precision	Accuracy	Range	Mean			Median	
79-0028 (cont'd)		Sr	mmol kg ⁻¹	6	18	18	SCUBA	Freeze; desiccated in whirligig bags	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; GF AAS	NS	10.0% +1.8% +7.8%	NC	0.029	NC	1	lower - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		Hg	µmol kg ⁻¹	6	18	18	SCUBA	Freeze; desiccated in whirligig bags	addition of V ₂ O ₅ ; HNO ₃ /H ₂ SO ₄ digestion; GF AAS	NS	NS	NC	2.7	NC	2	lower - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		Pb	µmol kg ⁻¹	5	11	11	SCUBA	Freeze; desiccated in whirligig bags	HNO ₃ digestion; addition of NaBH ₄ ; DDC extraction; GF AAS	NS	-2.1% +2.0% +8.0%	NC	2.1	NC	3	live - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		Zn	mmol kg ⁻¹	5	11	11	SCUBA	Freeze; desiccated in whirligig bags	HNO ₃ digestion; addition of NaBH ₄ ; GF AAS	NS	-2.0% +1.9% +16.7%	NC	1.48	NC	3	live - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		Cd	µmol kg ⁻¹	5	11	11	SCUBA	Freeze; desiccated in whirligig bags	HNO ₃ digestion; addition of NaBH ₄ ; GF AAS	NS	-3.1% +2.0%	NC	30.7	NC	3	live - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
80-0027	SEA WATER	As	mmol kg ⁻¹	6	18	18	SCUBA	Freeze; desiccated in whirligig bags	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; GF AAS	NS	-6.7% +27.7% +10.0%	NC	0.217	NC	3	live - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		Sr	mmol kg ⁻¹	6	18	18	SCUBA	Freeze; desiccated in whirligig bags	HNO ₃ /H ₂ SO ₄ /H ₂ O ₂ digestion; addition of NaBH ₄ ; GF AAS	NS	-10.0% +1.8% +7.8%	NC	0.088	NC	3	live - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		Hg	µmol kg ⁻¹	6	17	17	SCUBA	Freeze; desiccated in whirligig bags	addition of V ₂ O ₅ ; HNO ₃ /H ₂ SO ₄ digestion; GF AAS	NS	NS	NC	0.80	NC	2	live - (DB); data presented as mean values; accuracy based on NBS system; live and lute reference materials
		O ₂	mol m ⁻³	9	426	426	NS	NS	NS	NS	0.122 - 0.445	0.119	0.113	2	NS 10	
		SiO ₂	mmol m ⁻³	9	421	421	NS	NS	NS	NS	0.22 - 61.70	16.76	19.06	3	NS 10	
80-0029	SEA WATER	PO ₄	mmol m ⁻³	9	421	421	NS	NS	NS	NS	0.47 - 3.61	1.30	1.36	3	NS 10	
		NO ₃	mmol m ⁻³	9	421	136	NS	NS	NS	NS	4.1 - 20.87	NC	10.06	3	6.1 not given; NS 10	
		O ₂	mol m ⁻³	3	13	13	NS	NS	High-act	NS	0.47 - 0.48	0.47	0.47	1	information retained only; original values; accuracy based on NBS system; live and lute reference materials	

9. DATA INVENTORY TABLE 3.

NOTES APPLICABLE TO TABLE 3

- Note 1. Latitude and longitude approximated from hydrographic chart.
- Note 2. 20 stations were sampled for O₂ and PO₄ determinations in Prince Regent Inlet and Gulf of Boothia.
- Note 3. 13 stations were sampled in Lancaster Sound and Barrow Strait for O₂ determination only.
- Note 4. The reported time is doubtful.
- Note 5. One-ringed seal muscle sample.
- Note 6. Two polar bear brain samples.
- Note 7. One polar bear muscle sample.
- Note 8. The station depth represents the mean depth of Resolute Bay.
- Note 9. The measurements were made either in Resolute Bay or immediately outside the bay.
- Note 10. 1 arctic char sample.
- Note 11. Sampling between April 20 (2314 GMT) and April 21 (0126 GMT).
- Note 12. Sampling between August and September.
- Note 13. 11 stations within Cunningham Inlet were sampled.
- Note 14. 77 sculpins.
- Note 15. 14 arctic char.
- Note 16. 1 greenland shark.
- Note 17. 3 samples of shrimp.
- Note 18. 1 sample of seaweed.
- Note 19. 2 samples of seaweed.
- Note 20. Sampling area is approximately between 84° 30' W and 84° 45' W along southern shore of Strathcona Sound.
- Note 21. Ice core.
- Note 22. Sampling location only represents the general location of Strathcona Sound.

- Note 23. Between September 4 and 29, 1975.
- Note 24. 28 samples of bivalve molluscs and gastropods collected in vicinity of Station 3.
- Note 25. Station depths estimated from bathymetry diagram of Strathcona Sound.
- Note 26. Geographic coordinates represent the approximate area of trawling.
- Note 27. 84 samples of arctic cod.
- Note 28. 2 samples of copepods.
- Note 29. 1 sample of amphipods.
- Note 30. 2 samples of chaetognaths.
- Note 31. 4 samples of furoid algae.
- Note 32. 8 samples of furoid algae and 10 samples of purple sea urchin.
- Note 33. 5 samples of furoid algae.
- Note 34. 10 samples of furoid algae.
- Note 35. 3 samples of purple sea urchin.
- Note 36. 6 samples of furoid algae.
- Note 37. 89 samples of ringed seal muscle.
- Note 38. 88 samples of ringed seal muscle.
- Note 39. Early July, approximately July 3.
- Note 40. At 300 m Knudsen sampler did not close tightly.
- Note 41. 8 samples of seaweeds.
- Note 42. 2 samples of sea urchins.
- Note 43. 13 samples of sea urchins.
- Note 44. 17 samples of seaweeds.
- Note 45. 19 samples of sea urchins.
- Note 46. 5 northern Fulmars.
- Note 47. 10 black-legged Kittiwakes.
- Note 48. 4 thick-billed Murres.

- Note 49. 5 black Guillemots.
- Note 50. 6 thick-billed Murres.
- Note 51. 5 Dovekies.
- Note 52. 5 black Guillemots.
- Note 53. 5 Dovekies.
- Note 54. 5 northern Fulmars.
- Note 55. 27 ringed seals.
- Note 56. 1 bearded seal.
- Note 57. 37 ringed seals.
- Note 58. Between August 17 and September 21, 1978.
- Note 59. A total of 9 stations were sampled at this location.
- Note 60. 25 points within Cape Hay were sampled, spaced 0.83 km apart.
- Note 61. 4 samples of seaweeds.
- Note 62. 5 samples of sea urchins.
- Note 63. 8 samples of molluscs.
- Note 64. 3 samples of sea cucumber.
- Note 65. 2 samples of fish.
- Note 66. 9 samples of seaweeds.
- Note 67. 20 samples of molluscs.
- Note 68. 5 samples of fish.
- Note 69. 6 samples of seaweeds.
- Note 70. 3 samples of sea urchins.
- Note 71. 34 samples of molluscs.
- Note 72. 3 samples of sea stars.
- Note 73. 4 samples of fish.
- Note 74. 21 samples of seaweeds.
- Note 75. 6 samples of molluscs.

- Note 76. 8 samples of seaweeds.
- Note 77. 12 samples of sea urchins.
- Note 78. 13 samples of molluscs.
- Note 79. 8 samples of fish.
- Note 80. 7 samples of seaweeds.
- Note 81. 3 samples of fish.
- Note 82. 11 samples of sea urchins.
- Note 83. 5 samples of molluscs.
- Note 84. 1 sample of sea star.
- Note 85. 11 samples of seaweeds.
- Note 86. 12 samples of fish.
- Note 87. Station depth as estimated in the report.
- Note 88. Coordinates in error.

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
28-0001	122	74° 26.5'	81° 50'	735	1928	Aug.	20			
	124	74° 16.5'	81° 56'	770	1928	Aug.	20	2	700	
	126	74° 03'	82° 02'	650	1928	Aug.	20	4	750	
	128	73° 53'	82° 07.5'	775	1928	Aug.	20	4	620	
50-0002	12	74° 30'	82° 45'	713	1950	Aug.	17	2	775	
	13	74° 36'	92° 40'	173	1950	Aug.	17	6	400	
52-0003	35	74° 37'	94° 42'	120	1952	Aug.	8	5	149	
	36	74° 50'	101° 45'	182	1952	Aug.	9	9	100	
	37	74° 57'	108° 52'	63	1952	Aug.	11	4	150	
	38	74° 56'	105° 05'	100	1952	Aug.	12	3	50	
	39	74° 53'	102° 48'	128	1952	Aug.	12	7	100	
	40	74° 46'	101° 22'	128	1952	Aug.	12	6	100	
	41	74° 38'	96° 32'	256	1952	Aug.	13	6	100	
	12	71° 59'	120° 02'	23	1953	Aug.	14	6	200	
	13	71° 50'	118° 54'	66	1953	Aug.	15	3	15	
	14	71° 55'	119° 32'	97	1953	Aug.	15	4	25	
	15	72° 11'	119° 04'	97	1953	Aug.	15	6	90	
53-0001	17	72° 22'	118° 46'	80	1953	Aug.	16	6	90	
	18	72° 23'	118° 30'	42	1953	Aug.	16	5	40	
	19	73° 12'	115° 30'	53	1953	Aug.	20	5	40	
	20	73° 11'	115° 35'	86	1953	Aug.	20	4	40	
	21	72° 24'	118° 42'	82	1953	Aug.	22	5	80	
	22	72° 25'	119° 00'	49	1953	Aug.	22	4	40	
	23	72° 21'	119° 15'	63	1953	Aug.	22	5	60	
	24	72° 56'	117° 25'	51	1953	Aug.	23	4	40	
	25	72° 35'	118° 25'	55	1953	Aug.	23	4	40	
	26	72° 38'	118° 38'	92	1953	Aug.	24	4	40	
	27	72° 40'	118° 54'	101	1953	Aug.	24	5	85	
	28	72° 23'	118° 42'	74	1953	Aug.	24	5	95	
	29	72° 23'	118° 58'	93	1953	Aug.	25	5	70	
	30	72° 22'	119° 14'	64	1953	Aug.	25	5	90	
	31	71° 58'	120° 13'	75	1953	Aug.	25	4	40	
	32	71° 56'	119° 38'	120	1953	Aug.	25	5	115	
	33	71° 53'	119° 08'	26	1953	Aug.	25	5	15	
	34	71° 38'	119° 10'	21	1953	Aug.	25	2	15	
	35	71° 31'	119° 50'	150	1953	Aug.	25	3	18	
	37	73° 24'	115° 05'	54	1953	Aug.	31	6	145	
	38	73° 21'	115° 03'	86	1953	Aug.	31	4	40	
	39	73° 14'	115° 30'	109	1953	Aug.	31	5	80	
	40	73° 05'	116° 38'	115	1953	Sept.	1	4	40	
	41	73° 02'	116° 34'	150	1953	Sept.	1	6	140	
	42	73° 01'	116° 28'	42	1953	Sept.	1	4	45	

DATA SET NUMBER	STATION NUMBER	LATITUDE	LONGITUDE	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS		
					YEAR	MONTH	DAY					
54-0002	7	73° 27'	115° 03'	45	1954	Aug.	11	0200	6	40		
	8	73° 27'	115° 04'	30	1954	Aug.	11	2000	5	25		
	9	74° 30'	113° 47'	145	1954	Aug.	13	2000	7	130		
	10	74° 24'	113° 50'	362	1954	Aug.	13	2300	11	350		
	11	73° 57'	114° 24'	493	1954	Aug.	14	2000	11	475		
	12	73° 27'	114° 42'	100	1954	Aug.	15	0900	7	89		
	13	73° 56'	115° 50'	430	1954	Aug.	15	1900	10	400		
	14	73° 48'	116° 06'	130	1954	Aug.	15	2300	7	120		
	15	74° 37'	117° 22'	480	1954	Aug.	17	1900	11	450		
	16	74° 25'	117° 09'	493	1954	Aug.	18	1200	11	475		
	17	74° 09'	115° 55'	485	1954	Aug.	18	1900	9	450		
	19	74° 24'	112° 30'	235	1954	Aug.	19	1900	8	220		
	20	74° 59'	108° 50'	48	1954	Aug.	24	2200	3	10		
	21	74° 59'	108° 50'	48	1954	Aug.	25	0500	6	40		
	23	74° 59'	108° 50'	48	1954	Aug.	25	1700	6	30		
	24	74° 51'	109° 21'	55	1954	Aug.	27	0200	6	45		
	25	74° 40'	110° 36'	45	1954	Aug.	27	1900	4	45		
	26	74° 33'	111° 06'	226	1954	Aug.	27	2100	9	250		
	27	74° 31'	111° 30'	98	1954	Aug.	28	0300	5	90		
	28	74° 24'	111° 33'	329	1954	Aug.	28	0500	9	325		
	29	74° 06'	111° 33'	493	1954	Aug.	28	0800	10	490		
	30	73° 47'	111° 33'	285	1954	Aug.	28	1300	9	275		
	31	73° 32'	111° 27'	179	1954	Aug.	28	1600	8	175		
	32	73° 15'	113° 03'	201	1954	Aug.	30	0700	8	199		
	33	73° 33'	113° 07'	170	1954	Aug.	30	1100	7	165		
	34	73° 51'	112° 27'	430	1954	Aug.	30	1400	10	417		
	35	73° 45'	113° 45'	435	1954	Aug.	30	2200	10	388		
	36	73° 25'	114° 13'	38	1954	Aug.	31	0300	6	34		
	37	73° 27'	114° 29'	76	1954	Aug.	31	0800	7	68		
	38	73° 07'	116° 39'	70	1954	Sept.	4	1800	7	70		
	39	73° 14'	116° 18'	70	1954	Sept.	4	2300	7	70		
	41	73° 03'	116° 46'	75	1954	Sept.	5	0300	6	75		
	42	72° 19'	119° 20'	65	1954	Sept.	6	1000	6	60		
	43	72° 22'	118° 55'	80	1954	Sept.	6	1200	7	80		
	45	71° 36'	119° 50'	150	1954	Sept.	6	1800	8	150		
	54-0003	8	74° 35'	121° 27'	243	1954	Aug.	14	1500	11	197	
		9	74° 12'	119° 02'	57	1954	Aug.	15	2300	8	50	
		10	74° 25'	118° 24'	448	1954	Aug.	16	1600	10	400	
		11	74° 16'	117° 33'	351	1954	Aug.	16	2100	11	300	

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
54-0003 (cont'd)	12	74° 18'	118° 51'	342	1954	Aug.	17	12	299	
	13	74° 16'	117° 32'	346	1954	Aug.	18	11	299	
	14	74° 16'	117° 32'	346	1954	Aug.	19	10	196	
	15	73° 55'	116° 23'	210	1954	Aug.	20	10	173	
	16	73° 21'	115° 51'	20	1954	Aug.	23	6	15	
	17	73° 23'	108° 22'	117	1954	Aug.	24	7	105	
	18	73° 23'	109° 39'	132	1954	Aug.	25	9	130	
	19	72° 58'	109° 02'	73	1954	Aug.	25	7	65	
	20	73° 18'	109° 57'	276	1954	Aug.	26	8	200	
	21	73° 32'	110° 51'	192	1954	Aug.	26	8	149	
	22	73° 09'	111° 24'	249	1954	Aug.	26	9	198	
	23	72° 59'	112° 55'	42	1954	Aug.	28	8	39	
	24	72° 46'	118° 07'	44	1954	Sept.	3	8	30	
	25	72° 05'	119° 00'	44	1954	Sept.	5	8	40	
	26	72° 01'	119° 42'	104	1954	Sept.	6	8	90	
	27	71° 58'	119° 08'	50	1954	Sept.	6	8	40	
56-0002	1	74° 43.94'	95° 00.83'	NS	1956	July	18	4	10	Note 1
	1	74° 43.94'	95° 00.83'	NS	1956	July	19	3	4	Note 1
	1	74° 43.94'	95° 00.83'	NS	1956	July	20	4	10	Note 1
	1	74° 43.94'	95° 00.83'	NS	1956	July	21	4	10	Note 1
	1	74° 43.94'	95° 00.83'	NS	1956	July	23	4	8	Note 1
	1	74° 43.94'	95° 00.83'	NS	1956	July	24	4	9	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	July	26	4	10	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	July	27	3	6	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	July	30	5	20	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	July	31	4	15	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	2	5	25	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	3	5	25	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	4	5	20	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	5	4	15	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	7	5	50	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	8	5	20	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	10	2	3	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	11	4	15	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	12	5	20	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	14	3	10	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	15	5	25	Note 1
	2	74° 42.62'	95° 00.99'	NS	1956	Aug.	18	4	15	Note 1

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
56-0002 (cont'd)	2	74° 42.62"	93° 00.99'	NS	1956	Aug.	19	5	25	Note 1
	2	74° 42.62"	93° 00.99'	NS	1956	Aug.	20	5	25	Note 1
	2	74° 42.62"	93° 00.99'	NS	1956	Aug.	23	5	25	Note 1
	2	74° 42.62"	93° 00.99'	NS	1956	Aug.	28	5	25	Note 1
56-0003	Prince Regent Inlet	73° 30'	91° 00'	NS	1956	Sept.	NS	NS	bottom	Note 2
	Gulf of Boothia	71° 30'	93° 00'	NS	1956	Sept.	NS	NS	bottom	Note 2
	Lancaster Sound	74° 00'	86° 00'	NS	1956	Sept.	NS	NS	bottom	Note 3
	Barrow Strait	74° 30'	95° 00'	NS	1956	Sept.	NS	NS	bottom	Note 3
57-0003	35	74° 01'	89° 46'	247	1957	Aug.	20	9	200	
	37	73° 42'	87° 46'	236	1957	Aug.	20	9	200	
	39	73° 42'	89° 19'	457	1957	Aug.	20	11	400	
	41	73° 14'	91° 03'	289	1957	Aug.	21	10	250	
	43	73° 02'	89° 28'	183	1957	Aug.	21	8	150	
	45	72° 18'	90° 16'	73	1957	Aug.	21	5	50	
	47	72° 14'	92° 11'	101	1957	Aug.	21	7	90	
	50	72° 01'	95° 49'	412	1957	Aug.	25	11	312	
	53	71° 48'	96° 00'	439	1957	Aug.	25	11	398	
	55	72° 11'	93° 16'	137	1957	Aug.	26	8	120	
	57	71° 47'	93° 16'	75	1957	Aug.	26	6	75	
	59	72° 28'	90° 07'	104	1957	Aug.	27	7	100	
	61	72° 41'	92° 03'	223	1957	Aug.	27	9	195	
	62	74° 10'	92° 27'	145	1957	Aug.	28	8	140	
	64	74° 30'	92° 08'	183	1957	Aug.	28	8	150	
	66	74° 40'	92° 30'	132	1957	Aug.	28	8	120	
	68	74° 37'	93° 31'	177	1957	Aug.	28	8	145	
	70	74° 22'	93° 26'	165	1957	Aug.	28	8	150	
	71	74° 11'	94° 16'	152	1957	Aug.	29	8	150	
	72	74° 19'	94° 19'	157	1957	Aug.	29	8	150	
	73	74° 28'	94° 22'	165	1957	Aug.	29	8	150	

DATA SET NUMBER	STATION NUMBER	LATITUDE	LONGITUDE		STATION DEPTH (m)	TIME SAMPLED				NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
			*N	*W		YEAR	MONTH	DAY	HOUR GMT			
57-0003 (cont'd)	77	74° 18'	95° 14'	165	1957	Aug.	30	1400	10	149		
	80	73° 43'	96° 20'	247	1957	Aug.	30	2000	9	194		
	82	73° 11'	96° 14'	247	1957	Aug.	31	0100	8	196		
	83	72° 15'	96° 12'	357	1957	Aug.	31	0400	10	324		
	84	70° 56'	96° 54'	85	1957	Sept.	4	0100	6	75		
	85	71° 01'	97° 20'	87	1957	Sept.	4	0200	6	75		
	86	71° 09'	98° 00'	130	1957	Sept.	4	0400	7	100		
	88	71° 46'	92° 23'	124	1957	Sept.	8	0400	7	100		
	90	71° 47'	90° 11'	46	1957	Sept.	8	0900	5	40		
	92	71° 16'	90° 58'	174	1957	Sept.	8	1300	8	150		
	93	71° 10'	91° 47'	220	1957	Sept.	8	1500	9	200		
	96	70° 42'	91° 09'	146	1957	Sept.	8	2100	8	125		
	97	70° 47'	90° 22'	128	1957	Sept.	8	2200	7	100		
	100	70° 23'	89° 40'	128	1957	Sept.	9	0300	7	100		
60-0008	101	70° 16'	90° 26'	146	1957	Sept.	9	0800	8	125		
	103	74° 05'	96° 05'	192	1957	Sept.	13	2100	9	170		
	106	74° 04'	96° 44'	220	1957	Sept.	13	2300	9	200		
	126	74° 29'	90° 00'	274	1957	Sept.	18	0000	10	250		
	128	74° 14'	90° 02'	238	1957	Sept.	18	0300	9	200		
	130	73° 49'	87° 11'	284	1957	Sept.	19	0300	9	230		
	131	74° 01'	87° 09'	457	1957	Sept.	19	0300	9	396		
	132	74° 13'	87° 10'	424	1957	Sept.	19	0700	10	307		
	133	74° 25'	87° 10'	402	1957	Sept.	19	0800	6	355		
	135	74° 02'	85° 52'	521	1957	Sept.	20	1700	11	439		
	137	74° 28'	84° 52'	272	1957	Sept.	20	2000	9	239		
	139	74° 15'	82° 46'	768	1957	Sept.	21	0100	11	704		
	141	73° 48'	82° 52'	530	1957	Sept.	21	0500	10	500		
	142	73° 45'	80° 12'	635	1957	Sept.	23	1500	10	500		
143	74° 02'	80° 15'	801	1957	Sept.	23	2100	10	600			
145	74° 33'	80° 21'	640	1957	Sept.	24	0200	11	500			
60-0008	3	74° 32'	82° 57'	626	1960	NS	NS	NS	1	bottom	Note 1	
	4	74° 30'	82° 40'	746	1960	NS	NS	NS	1	bottom	Note 1	
	7	74° 27.5'	84° 28'	604	1960	NS	NS	NS	1	bottom	Note 1	
	10	74° 29'	83° 30'	752	1960	NS	NS	NS	1	bottom	Note 1	
	12	74° 28'	82° 58'	772	1960	NS	NS	NS	1	bottom	Note 1	
	13	74° 28'	82° 40'	784	1960	NS	NS	NS	1	bottom	Note 1	
	17	74° 24'	84° 45'	576	1960	NS	NS	NS	1	bottom	Note 1	
	19	74° 23.5'	84° 05'	674	1960	NS	NS	NS	1	bottom	Note 1	
21	74° 23'	83° 30'	764	1960	NS	NS	NS	1	bottom	Note 1		

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
60-0008 (Cont'd)	23	74° 23'	82° 57'	776	1960	NS	NS	1	bottom	Note 1
	26	74° 23.5'	82° 10'	766	1960	NS	NS	1	bottom	Note 1
	27	74° 18.5'	85° 05'	580	1960	NS	NS	1	bottom	Note 1
	30	74° 18.5'	84° 05'	634	1960	NS	NS	1	bottom	Note 1
	33	74° 19'	83° 11'	808	1960	NS	NS	1	bottom	Note 1
	36	74° 19'	82° 23'	868	1960	NS	NS	1	bottom	Note 1
	41	74° 14.3'	84° 05'	652	1960	NS	NS	1	bottom	Note 1
	46	74° 14.5'	82° 38'	828	1960	NS	NS	1	bottom	Note 1
	51	74° 09.5'	84° 23'	624	1960	NS	NS	1	bottom	Note 1
	54	74° 10'	83° 25'	732	1960	NS	NS	1	bottom	Note 1
	57	74° 10'	82° 36'	776	1960	NS	NS	1	bottom	Note 1
	60	74° 04.5'	85° 03'	512	1960	NS	NS	1	bottom	Note 1
	63	74° 05'	84° 03'	708	1960	NS	NS	1	bottom	Note 1
	66	74° 05'	83° 06'	732	1960	NS	NS	1	bottom	Note 1
	69	74° 06.5'	82° 19'	774	1960	NS	NS	1	bottom	Note 1
	72	74° 00.5'	84° 41'	520	1960	NS	NS	1	bottom	Note 1
	74	74° 01'	84° 03'	638	1960	NS	NS	1	bottom	Note 1
	78	74° 01.5'	82° 50'	788	1960	NS	NS	1	bottom	Note 1
	82	73° 56'	85° 00'	372	1960	NS	NS	1	bottom	Note 1
	84	73° 56'	84° 21'	432	1960	NS	NS	1	bottom	Note 1
	86	73° 56.5'	83° 38'	416	1960	NS	NS	1	bottom	Note 1
	89	73° 57'	82° 48'	700	1960	NS	NS	1	bottom	Note 1
	91	73° 58'	82° 18'	700	1960	NS	NS	1	bottom	Note 1
	93	73° 51.5'	84° 20'	288	1960	NS	NS	1	bottom	Note 1
	96	73° 52.2'	83° 20'	324	1960	NS	NS	1	bottom	Note 1
	99	73° 53'	82° 30'	344	1960	NS	NS	1	bottom	Note 1
	101	73° 54'	82° 03'	316	1960	NS	NS	1	bottom	Note 1
	104	73° 48'	84° 00'	344	1960	NS	NS	1	bottom	Note 1
	106	73° 48'	83° 20'	380	1960	NS	NS	1	bottom	Note 1
	108	73° 43.5'	84° 40'	328	1960	NS	NS	1	bottom	Note 1
	111	73° 43.5'	83° 34'	456	1960	NS	NS	1	bottom	Note 1
	113	73° 39.5'	84° 20'	684	1960	NS	NS	1	bottom	Note 1
61-0004	78	74° 28.5'	115° 05.0'	485	1961	Aug.	28	12	400	
	80	74° 31.0'	115° 33.0'	483	1961	Aug.	29	13	450	
	82	74° 28.2'	115° 30.8'	474	1961	Aug.	29	7	100	
	83	74° 27.0'	115° 30.0'	491	1961	Aug.	31	13	450	
	85	74° 21.3'	115° 18.2'	494	1961	Sept.	1	13	450	
	86	74° 22.0'	115° 09.0'	488	1961	Sept.	2	13	450	

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
61-0004 (cont'd)	87	74° 25.3'	111° 54.5'	329	1961	Sept.	4	11	2124	300
	90	74° 30.0'	94° 30.0'	155	1961	Sept.	7	8	0400	140
	92	74° 17.0'	94° 30.0'	184	1961	Sept.	7	8	0900	150
	101	74° 19.0'	91° 55.0'	228	1961	Sept.	8	9	0200	200
	102	74° 23.0'	91° 55.0'	311	1961	Sept.	8	10	0418	250
	103	74° 24.5'	87° 10.0'	454	1961	Sept.	9	12	0612	400
	104	74° 12.0'	87° 12.0'	446	1961	Sept.	9	12	0818	400
	105	74° 02.0'	87° 11.0'	485	1961	Sept.	9	13	1012	450
	108	72° 40.0'	91° 26.0'	333	1961	Sept.	11	11	1624	297
	110	72° 34.0'	90° 25.0'	346	1961	Sept.	11	11	2012	300
	111	72° 33.0'	89° 59.0'	79	1961	Sept.	11	6	2148	68
	114	73° 12.5'	90° 08.0'	508	1961	Sept.	12	12	0536	500
	116	73° 17.0'	90° 58.0'	285	1961	Sept.	12	10	0912	250
	123	74° 04.0'	87° 10.0'	419	1961	Sept.	13	12	1512	396
	124	74° 13.0'	87° 10.0'	430	1961	Sept.	13	12	1724	400
	130	73° 48.5'	82° 42.5'	622	1961	Sept.	14	14	1212	600
	131	74° 01.0'	82° 42.5'	686	1961	Sept.	14	15	1530	650
	132	74° 15.0'	82° 44.0'	793	1961	Sept.	14	13	1854	500
	133	74° 28.5'	82° 47.0'	662	1961	Sept.	14	14	2206	587
62-0005 A	1	74° 36.0'	94° 13.0'	55	1962	June	28	9	2006	50
	1	74° 36.0'	94° 13.0'	52	1962	July	6	9	2300	50
	1	74° 36.0'	94° 13.0'	62	1962	July	13	9	1524	50
	1	74° 36.0'	94° 13.0'	60	1962	July	25	9	0236	50
	1	74° 36.0'	94° 13.0'	55	1962	Aug.	7	9	0106	50
	1	74° 36.0'	94° 13.0'	41	1962	Aug.	14	9	1818	40
62-0005 B	2	72° 45.0'	94° 04.0'	34	1962	June	23	8	2300	33
	2	72° 45.0'	94° 04.0'	34	1962	June	30	8	1748	33
	2	72° 45.0'	94° 04.0'	34	1962	July	14	9	2130	32
	2	72° 45.0'	94° 04.0'	46	1962	July	21	9	1848	45
	2	72° 45.0'	94° 04.0'	46	1962	July	29	11	0018	45
	2	72° 45.0'	94° 04.0'	41	1962	Aug.	4	10	1618	40
	2	72° 45.0'	94° 04.0'	31	1962	Aug.	11	9	1724	30
	3	72° 45.3'	94° 06.0'	35	1962	June	27	9	2248	34
	8	72° 45.5'	94° 04.5'	44	1962	July	3	2	1830	43
	32	72° 46.0'	94° 17.0'	62	1962	Aug.	13	1	1930	61

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY	HOUR GMT		
62-0005 C	80	69° 07.0'	105° 10.0'	50	1962	Sept.	7	1200	49	Note 4
62-0006	9	74° 23.0'	92° 00.0'	300	1962	Aug.	7	0406	10	250
	63	71° 56.0'	95° 32.0'	258	1962	Sept.	3	1536	9	200
	69	72° 58.0'	102° 54.0'	348	1962	Sept.	6	2242	11	330
62-0009	11	69° 10'	106° 20'	NS	1962	Aug.	27	NS	1	0
65-0002	1	69° 06.3'	105° 02.2'	39	1965	July	29	NS	8	38
	1	69° 06.3'	105° 02.2'	41	1965	Aug.	5	NS	8	40
	2	69° 06.1'	105° 06.0'	73	1965	Aug.	4	NS	8	70
	4	68° 20.0'	107° 38.0'	13	1965	Aug.	10	NS	4	12
	8	68° 18.2'	109° 15.0'	123	1965	Aug.	23	NS	9	120
	9	69° 10.0'	105° 50.0'	51	1965	Aug.	27	NS	7	50
	10	69° 07.0'	105° 01.0'	43	1965	Aug.	28	NS	7	42
	13	69° 06.3'	105° 04.8'	57	1965	Sept.	5	NS	7	50
	49	74° 08.9'	81° 22.0'	786	1966	Aug.	6	0530	13	755
	50	74° 02.0'	81° 08.0'	748	1966	Aug.	6	0818	13	710
66-0004	55	74° 09.2'	81° 27.4'	784	1966	Aug.	6	2218	12	706
	56	74° 01.9'	81° 08.5'	757	1966	Aug.	7	0048	12	691
	1	69° 09.6'	105° 51.3'	55	1966	Aug.	2	NS	6	50
	2	69° 09.5'	105° 52.7'	56	1966	Aug.	11	NS	6	50
66-0005	2	69° 09.5'	105° 52.7'	51	1966	Aug.	17	NS	6	50
	2	69° 09.5'	105° 52.7'	54	1966	Aug.	24	NS	6	50
	1	69° 09.5'	105° 52.7'	54	1967	Aug.	17	NS	7	49
	1	69° 09.5'	105° 52.7'	51	1967	Aug.	24	NS	7	50
67-0001 A	1	69° 09.5'	105° 52.7'	54	1967	Sept.	1	NS	7	50
	1	69° 09.5'	105° 52.7'	54	1967	Sept.	7	NS	7	50
	1	69° 09.5'	105° 52.7'	54	1967	Sept.	7	NS	7	50
67-0001 B	1	69° 08.7'	105° 53.0'	50	1967	July	2	NS	6	47
	1	69° 08.7'	105° 53.0'	50	1967	July	8	NS	7	45
	3	69° 08.0'	105° 57.0'	31	1967	July	16	NS	6	30
	4	69° 08.6'	105° 53.0'	44	1967	July	29	NS	7	43

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
68-0001	13	73° 40.0'	81° 23.0'	NS	1968	Aug.	24	12	350	
	14	74° 10.0'	85° 59.4'	NS	1968	Aug.	24	13	500	
	15	74° 38.5'	109° 47.5'	NS	1968	Aug.	30	10	175	
	16	74° 18.0'	110° 00.0'	NS	1968	Aug.	31	13	500	
	17	74° 27.0'	113° 48.0'	NS	1968	Aug.	31	12	325	
	18	74° 22.3'	114° 36.0'	NS	1968	Aug.	31	13	450	
70-0010	Lancaster Sound	74° 10'	84° 00'	NS	1970	May	NS	NS		Note 1, 5
71-0009	Spence Bay	69° 30'	93° 30'	NS	1971	Nov.	NS	NS		Note 1, 6
	Spence Bay	69° 30'	93° 30'	NS	1971	Dec.	NS	NS		Note 1, 7
71-0016	Resolute Bay	74° 42'	94° 50'	11.3	1971	Sept.	NS	3	10	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1971	Oct.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1971	Nov.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1971	Dec.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	Jan.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	Feb.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	March	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	April	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	May	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	June	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	July	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	Aug.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	Aug.	NS	1	0	Note 8, 9
	Resolute Bay	74° 42'	94° 50'	11.3	1972	Aug.	NS	1	0	Note 8, 9

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
72-0011	1	74° 21.0'	90° 42.0'	200	1972	Sept.	27	12	200	
	2	74° 44.5'	93° 14.0'	130	1972	Sept.	28	9	130	
	3	74° 45.0'	92° 44.0'	100	1972	Sept.	28	7	100	
	4	74° 46.0'	92° 10.0'	60	1972	Sept.	28	7	60	
	5	74° 06.5'	91° 38.0'	150	1972	Sept.	29	10	150	
	6	74° 21.6'	91° 19.0'	170	1972	Sept.	29	11	170	
	7	74° 35.0'	91° 23.0'	100	1972	Sept.	30	7	100	
	SED-1	74° 30.1'	91° 34.5'	NS	1972	Sept.	NS	1	bottom	
	SED-2	74° 34.0'	91° 34.0'	NS	1972	Sept.	NS	1	bottom	
	SED-3	74° 29.0'	91° 34.0'	NS	1972	Sept.	NS	1	bottom	
	SED-4	74° 29.5'	91° 32.0'	NS	1972	Sept.	NS	1	bottom	
72-0016	SED-5	74° 31.5'	91° 32.0'	NS	1972	Sept.	NS	1	bottom	
	SED-6	74° 21.5'	91° 19.0'	NS	1972	Sept.	NS	1	bottom	
	SED-7	74° 21.0'	90° 42.0'	NS	1972	Sept.	NS	1	bottom	
	SED-8	74° 20.2'	91° 55.0'	NS	1972	Sept.	NS	1	bottom	
	SED-9	74° 15.2'	92° 09.0'	NS	1972	Sept.	NS	1	bottom	
	SED-10	74° 14.2'	91° 58.0'	NS	1972	Sept.	NS	1	bottom	
	SED-11	74° 07.4'	91° 35.0'	NS	1972	Sept.	NS	1	bottom	
	Resolute Bay	74° 42'	94° 50'	NS	1972	NS	NS	NS		Note 1, 7
	12	74° 42'	94° 53'	NS	1972	June	NS	NS		Note 10
72-0017	8	74° 20.0'	88° 30.0'	NS	1972	Aug.	20	1	0	
	9	74° 30.0'	94° 05.0'	NS	1972	Aug.	20	1	0	
73-0005	1	74° 27.8'	90° 57.0'	188	1973	Aug.	6	10	170	
	2	74° 19.5'	90° 53.0'	229	1973	Aug.	6	10	220	
	3	74° 14.8'	91° 00.0'	351	1973	Aug.	6	12	330	
	4	74° 09.2'	90° 58.0'	245	1973	Aug.	6	11	220	
	5	74° 05.0'	90° 58.0'	164	1973	Aug.	6	9	140	
73-0006	BA 1	74° 30'	91° 21'	155	1973	April		9	150	Note 11

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
73-0009 A	18	74° 18.0'	86° 28.0'	NS	1973	Aug.	7	1	0	
	19	74° 37.0'	88° 45.0'	NS	1973	Sept.	19	1	0	
73-0009 B	MB-T20	74° 39.5'	88° 42.7'	NS	1973	NS	NS	1	bottom	Note 12
	MB-Z20	74° 39.6'	88° 31.7'	NS	1973	NS	NS	1	bottom	Note 12
	MB-T23	74° 41.1'	88° 44.2'	NS	1973	NS	NS	1	bottom	Note 12
	MB-W23	74° 41.1'	88° 38.0'	NS	1973	NS	NS	1	bottom	Note 12
	MB-Z23	74° 41.1'	88° 32.5'	NS	1973	NS	NS	1	bottom	Note 12
	MB-R74	74° 41.6'	88° 48.0'	NS	1973	NS	NS	1	bottom	Note 12
	MB-T26	74° 42.7'	88° 44.2'	NS	1973	NS	NS	1	bottom	Note 12
	MB-W26	74° 42.8'	88° 38.0'	NS	1973	NS	NS	1	bottom	Note 12
	MB-T29	74° 43.6'	88° 44.1'	NS	1973	NS	NS	1	bottom	Note 12
	MB-W29	74° 44.3'	88° 38.6'	NS	1973	NS	NS	1	bottom	Note 12
	MB-Z29	74° 44.3'	88° 31.6'	NS	1973	NS	NS	1	bottom	Note 12
	MB-BB30	74° 45.0'	88° 28.1'	NS	1973	NS	NS	1	bottom	Note 12
	MB-W32	74° 45.9'	88° 38.5'	NS	1973	NS	NS	1	bottom	Note 12
	MB-Z46	74° 53.4'	88° 33.3'	NS	1973	NS	NS	1	bottom	Note 12
	MB-S00	74° 26.9'	88° 44.5'	282	1973	Sept.	19	1	bottom	Note 12
	MB-S04	74° 29.1'	88° 44.5'	174	1973	Sept.	19	1	bottom	0345
	MB-S08	74° 31.7'	88° 44.5'	106	1973	Sept.	19	1	bottom	0414
	MB-S12	74° 34.2'	88° 44.5'	146	1973	Sept.	19	1	bottom	0445
	MB-S16	74° 36.8'	88° 44.5'	121	1973	Sept.	19	1	bottom	0511
	MB-O16	74° 36.8'	88° 53.0'	128	1973	Sept.	19	1	bottom	0535
	MB-O12	74° 34.2'	88° 53.0'	119	1973	Sept.	19	1	bottom	0600
	MB-O08	74° 31.7'	88° 53.0'	117	1973	Sept.	19	1	bottom	0620
	MB-O04	74° 29.1'	88° 53.0'	179	1973	Sept.	19	1	bottom	0641
	MB-K04	74° 29.1'	89° 02.5'	198	1973	Sept.	19	1	bottom	0705
	MB-G04	74° 29.1'	89° 12.0'	227	1973	Sept.	19	1	bottom	0735
	MB-G08	74° 31.7'	89° 12.0'	68	1973	Sept.	19	1	bottom	0810
	MB-K08	74° 31.7'	89° 02.5'	117	1973	Sept.	19	1	bottom	0840
	MB-K12	74° 34.2'	89° 02.5'	90	1973	Sept.	19	1	bottom	0915
	MB-K16	74° 36.8'	89° 02.5'	86	1973	Sept.	19	1	bottom	0945
	LS-1	74° 27'	88° 18'	NS	1973	Sept.	19	1	bottom	1015
	LS-2	74° 22'	88° 18'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-3	74° 17'	88° 16'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-4	74° 12'	88° 16'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-5	74° 08'	88° 19'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-6	74° 02.5'	88° 15.5'	NS	1973	NS	NS	1	bottom	Note 1, 12

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
73-0009 B (cont'd)	LS-7	73° 57'	88° 13'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-8	74° 30.5'	89° 21.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-9	74° 26'	89° 25'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-10	74° 20.5'	89° 20'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-11	74° 15'	89° 21.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-12	74° 11'	89° 20'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-13	74° 06'	89° 20'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-14	74° 01'	89° 21'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-15	73° 55'	89° 20.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-16	74° 20'	87° 58'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-17	74° 14.3'	87° 56.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-18	74° 09.5'	87° 56.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-19	74° 05.1'	87° 56'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-20	74° 00'	87° 58'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-21	73° 54.8'	87° 59'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-22	73° 45'	87° 58'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-23	74° 28.5'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-24	74° 19'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-25	74° 24'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-26	74° 19'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-27	74° 14'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-28	74° 09.2'	86° 05.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-29	73° 55.1'	86° 42'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-30	73° 56.5'	87° 29'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-31	73° 57.2'	86° 58'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-32	73° 51.8'	86° 58'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-33	73° 56'	86° 59'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-34	74° 01.5'	86° 58.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-35	74° 04.5'	86° 58.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-36	74° 10.5'	86° 59'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-37	74° 16'	86° 58.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-38	74° 21'	86° 55'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-39	74° 49.6'	87° 36'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-40	73° 54'	87° 36'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-41	73° 59.1'	87° 39'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-42	73° 52.8'	87° 30'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-43	73° 53'	87° 14.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-44	73° 58'	87° 13'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-45	73° 55.3'	87° 25'	NS	1973	NS	NS	1	bottom	Note 1, 12

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
73-0009 B (cont'd)	LS-46	NS	NS	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-47	74° 26'	88° 58'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-48	74° 20.8'	88° 56'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-49	74° 15.2'	88° 55'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-50	74° 10.8'	88° 55'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-51	74° 06'	88° 56'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-52	74° 00.9'	88° 56'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-53	73° 55.3'	88° 52'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-54	74° 03.1'	87° 59'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-55	74° 09.8'	87° 59.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-56	74° 14.4'	87° 59.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-57	74° 19.5'	87° 59.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-58	74° 26.3'	88° 17'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-59	74° 26.3'	89° 49'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-60	74° 25.4'	89° 49'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-61	74° 20'	89° 50'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-62	74° 14.3'	89° 49'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-63	74° 09.4'	89° 50'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-64	74° 04.5'	89° 48'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-65	73° 59'	88° 15'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-66	73° 52'	88° 15'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-67	73° 47.5'	88° 15'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-68	74° 25'	87° 37'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-69	74° 20'	87° 37.5'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-70	74° 14.4'	87° 36'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-71	73° 10'	87° 35'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-72	74° 05.2'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-73	73° 49'	88° 38'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-74	73° 53.5'	88° 36'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-75	73° 58.7'	88° 36'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-76	74° 04'	88° 39'	NS	1973	NS	NS	1	bottom	Note 1, 12
	LS-77									
	LS-78									
73-0010	Resolute Bay	74° 42'	94° 50'	11.3	1973	May	8	1	ice	Note 1, 8
	Resolute Bay	74° 42'	94° 50'	11.3	1973	May	13	1	ice	Note 1, 8
73-0011	2	74° 15.5'	87° 03.5'	NS	1973	Aug.	15	1	0	
	4	74° 20.5'	86° 42.0'	NS	1973	Aug.	18	1	0	
	5	74° 24.5'	82° 30.0'	NS	1973	Aug.	18	1	0	

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
74-0013 A	59	73° 54.0'	82° 42.9'	NS	1974	Aug.	15	1	0	
	99	73° 52.2'	86° 06.9'	NS	1974	Aug.	25	1	0	
	104	73° 31.7'	96° 20.9'	NS	1974	Aug.	30	1	0	
74-0013 B	66	74° 19.1'	91° 08.9'	NS	1974	Aug.	20	1	bottom	
	69	74° 21.2'	91° 33.1'	NS	1974	Aug.	20	1	bottom	
	79	74° 12.1'	89° 18.5'	NS	1974	Aug.	21	1	bottom	
	81	73° 54.0'	88° 33.3'	NS	1974	Aug.	22	1	bottom	
	82	73° 53.7'	88° 27.9'	NS	1974	Aug.	22	1	bottom	
	86	74° 11.3'	89° 31.0'	NS	1974	Aug.	22	1	bottom	
	87	74° 09.9'	89° 25.8'	NS	1974	Aug.	23	1	bottom	
	89	74° 26.9'	86° 20.9'	NS	1974	Aug.	23	1	bottom	
	96	74° 45.5'	83° 16.0'	NS	1974	Aug.	24	1	bottom	
	97	74° 27.6'	83° 30.5'	NS	1974	Aug.	24	1	bottom	
	101	74° 19.2'	92° 37.9'	NS	1974	Aug.	26	1	bottom	
	103	74° 28.2'	92° 36.0'	NS	1974	Aug.	27	1	bottom	
	104	74° 37.8'	92° 30.4'	NS	1974	Aug.	27	1	bottom	
	112	74° 25.5'	94° 17.7'	NS	1974	Aug.	28	1	bottom	
	114	74° 15.8'	95° 55.1'	NS	1974	Aug.	28	1	bottom	
	115	74° 27.3'	97° 43.8'	NS	1974	Aug.	28	1	bottom	
	118	74° 24.1'	97° 45.0'	NS	1974	Aug.	28	1	bottom	
	120	74° 20.9'	97° 44.6'	NS	1974	Aug.	28	1	bottom	
	122	74° 17.8'	97° 43.9'	NS	1974	Aug.	28	1	bottom	
	123	74° 15.2'	97° 43.9'	NS	1974	Aug.	28	1	bottom	
	125	74° 11.9'	97° 43.9'	NS	1974	Aug.	28	1	bottom	
	126	74° 08.7'	97° 45.0'	NS	1974	Aug.	29	1	bottom	
	127	74° 09.5'	97° 14.8'	NS	1974	Aug.	29	1	bottom	
	130	74° 09.2'	96° 57.8'	NS	1974	Aug.	29	1	bottom	
	134	74° 08.4'	96° 53.9'	NS	1974	Aug.	29	1	bottom	
	136	73° 50.0'	96° 03.3'	NS	1974	Aug.	30	1	bottom	
	138	73° 32.1'	96° 18.4'	NS	1974	Aug.	30	1	bottom	
	146	73° 33.5'	96° 52.5'	NS	1974	Aug.	30	1	bottom	
	148	73° 33.6'	96° 06.3'	NS	1974	Aug.	30	1	bottom	
	151	73° 31.5'	96° 03.0'	NS	1974	Aug.	30	1	bottom	
	153	73° 29.8'	95° 47.5'	NS	1974	Aug.	31	1	bottom	
74-0014 A	1	74° 40'	97° 30'	NS	1974	May	7	3	30	Note 1
	1	74° 40'	97° 30'	NS	1974	May	7	7	1.4	Note 1, 21

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
74-0014 B	Cunningham Inlet	74° 10'	93° 40'	NS	1974	Aug.	14-27	1	0	Note 13
74-0015	1	72° 57'	84° 34'	55	1974	Aug.	8	6	bottom	Note 1
	1-A	72° 56.5'	84° 33.5'	NS	1974	Aug.	7	2	bottom	Note 1
	2	72° 56.5'	84° 25'	89	1974	Aug.	9	7	bottom	Note 1
	3	72° 59'	84° 23.5'	54	1974	Aug.	10	6	bottom	Note 1
	4	73° 01.5'	84° 22'	86	1974	Aug.	10	7	bottom	Note 1
	5	72° 58.5'	84° 37.5'	84	1974	Aug.	11	6	bottom	Note 1
	6	73° 00'	84° 37.5'	165	1974	Aug.	18	8	bottom	Note 1
	6-A	73° 01'	84° 37.3'	187	1974	Aug.	9	9	bottom	Note 1
	7	73° 03'	84° 37'	243	1974	Aug.	9	10	bottom	Note 1
	8	73° 06'	84° 36.5'	175	1974	Aug.	9	7	bottom	Note 1
	9	73° 03.5'	84° 43'	247	1974	Aug.	18	9	bottom	Note 1
	9-A	73° 02'	84° 42.8'	293	1974	Aug.	18	11	bottom	Note 1
	10	73° 03'	84° 50.5'	304	1974	Aug.	10	11	bottom	Note 1
	11	73° 10.5'	84° 59'	140	1974	Aug.	20	1	bottom	Note 1
	12	73° 11.5'	84° 58'	200	1974	Aug.	20	1	bottom	Note 1
	13	73° 15'	84° 54'	211	1974	Aug.	19	10	bottom	Note 1
	14	73° 16.5'	84° 53'	147	1974	Aug.	10	7	bottom	Note 1
	Strathcona Sound	NS	NS	NS	1974	Aug.	NS			Note 14, 20
	Strathcona Sound	NS	NS	NS	1974	Aug.	NS			Note 15, 20
	Strathcona Sound	NS	NS	NS	1974	Aug.	NS			Note 16, 20
	Strathcona Sound	NS	NS	NS	1974	Aug.	NS			Note 17, 20
	Shore-1	72° 56'	84° 26'	NS	1974	Aug.	NS			Note 1, 18
	Shore-2	72° 57'	84° 31'	NS	1974	Aug.	NS			Note 1, 18
	Shore-3	72° 58'	84° 33'	NS	1974	Aug.	NS			Note 1, 19
	Shore-4	72° 58'	84° 37'	NS	1974	Aug.	NS			Note 1, 19
74-0023	1	74° 47'	97° 47'	NS	1974	May	NS	7	1.4	Note 21

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
74-0024	22	74° 31.0'	93° 00.0'	NS	1974	Aug.	31	1	0	
	23	73° 47.0'	82° 23.0'	NS	1974	Sept.	15	1	0	
74-0026	Strathcona Sound	73° 07'	84° 45'	NS	1974	NS	NS	1	5	Note 22
	3	72° 56.5'	84° 33.5'	NS	1974	Sept.	NS	1	bottom	Note 1
75-0013	4	72° 46'	93° 29.8'	7	1975	Aug.	2	1	0	Note 1
	8	72° 45.3'	93° 30'	16	1975	Aug.	2	2	10	Note 1
	7	72° 44.9'	93° 30.4'	88	1975	Aug.	2	3	50	Note 1
	14	72° 46.3'	93° 24.5'	3	1975	Aug.	3	2	2	Note 1
	60	73° 59.9'	94° 18.5'	1	1975	Aug.	19	1	0	Note 1
	62	73° 58.8'	94° 18.5'	25	1975	Aug.	19	3	10	Note 1
	69	73° 58.4'	94° 18.5'	46	1975	Aug.	25	5	40	Note 1
	2	74° 55.0'	97° 39'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	4	74° 51.7'	97° 33'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	5	74° 49.0'	97° 33'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
75-0029	6	74° 46.7'	97° 33'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	7	74° 45.2'	97° 31'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	8	74° 44.0'	97° 31'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	9	74° 42.9'	97° 28'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	10	74° 25.2'	98° 43'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	11	74° 27.1'	98° 28'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	12	74° 13.9'	99° 01'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	13	74° 43.0'	98° 04'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	14	74° 44.1'	97° 58'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	15	74° 46.0'	97° 58'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	18	74° 48.5'	98° 12'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	19	74° 48.8'	98° 15'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	20	74° 49.1'	98° 15'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	21	74° 49.9'	98° 19'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	22	74° 50.2'	98° 24'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	23	74° 53.0'	98° 25'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	26	74° 55.8'	98° 42'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	27	74° 54.0'	98° 46'	NS	1975	Sept.	NS	1	bottom	Note 1, 23
	28	74° 52.7'	98° 48'	NS	1975	Sept.	NS	1	bottom	Note 1, 23

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
75-0029 (cont'd)	29	74° 51.1'	98° 53'	NS	1975	Sept.				Note 1, 23
	31	74° 48.0'	98° 59'	NS	1975	Sept.			bottom	Note 1, 23
	32	74° 47.0'	99° 00'	NS	1975	Sept.			bottom	Note 1, 23
	33	74° 45.6'	99° 04'	NS	1975	Sept.			bottom	Note 1, 23
	36	74° 52.1'	99° 55'	NS	1975	Sept.			bottom	Note 1, 23
	37	74° 52.5'	99° 35'	NS	1975	Sept.			bottom	Note 1, 23
	38	74° 52.2'	99° 08'	NS	1975	Sept.			bottom	Note 1, 23
	39	74° 52.0'	98° 48'	NS	1975	Sept.			bottom	Note 1, 23
	41	74° 51.8'	98° 11'	NS	1975	Sept.			bottom	Note 1, 23
	45	74° 57.5'	97° 55'	NS	1975	Sept.			bottom	Note 1, 23
	46	74° 55.3'	97° 55'	NS	1975	Sept.			bottom	Note 1, 23
	47	74° 52.0'	97° 56'	NS	1975	Sept.			bottom	Note 1, 23
	48	74° 49.0'	97° 56'	NS	1975	Sept.			bottom	Note 1, 23
	49	74° 46.0'	97° 54'	NS	1975	Sept.			bottom	Note 1, 23
	50	74° 43.0'	97° 55'	NS	1975	Sept.			bottom	Note 1, 23
	51	74° 40.0'	97° 55'	NS	1975	Sept.			bottom	Note 1, 23
	52	74° 37.0'	97° 56'	NS	1975	Sept.			bottom	Note 1, 23
	53	74° 34.0'	97° 56'	NS	1975	Sept.			bottom	Note 1, 23
	54	74° 33.5'	98° 05'	NS	1975	Sept.			bottom	Note 1, 23
	55	74° 34.0'	98° 17'	NS	1975	Sept.			bottom	Note 1, 23
	56	74° 36.5'	98° 18'	NS	1975	Sept.			bottom	Note 1, 23
	57	74° 40.0'	98° 18'	NS	1975	Sept.			bottom	Note 1, 23
	58	74° 42.5'	98° 18'	NS	1975	Sept.			bottom	Note 1, 23
	59	74° 42.5'	98° 28'	NS	1975	Sept.			bottom	Note 1, 23
	61	74° 40.0'	98° 38'	NS	1975	Sept.			bottom	Note 1, 23
	62	74° 37.0'	98° 38'	NS	1975	Sept.			bottom	Note 1, 23
	63	74° 34.6'	98° 37'	NS	1975	Sept.			bottom	Note 1, 23
	64	74° 39.3'	98° 59'	NS	1975	Sept.			bottom	Note 1, 23
	65	74° 42.0'	98° 59.5'	NS	1975	Sept.			bottom	Note 1, 23
	66	74° 45.0'	98° 59.5'	NS	1975	Sept.			bottom	Note 1, 23
	67	74° 48.4'	98° 57'	NS	1975	Sept.			bottom	Note 1, 23
	68	74° 50.0'	99° 00'	NS	1975	Sept.			bottom	Note 1, 23
	69	74° 53.0'	99° 15'	NS	1975	Sept.			bottom	Note 1, 23
	70	74° 54.0'	99° 20'	NS	1975	Sept.			bottom	Note 1, 23
	71	74° 51.0'	99° 19'	NS	1975	Sept.			bottom	Note 1, 23
	72	74° 47.5'	99° 22'	NS	1975	Sept.			bottom	Note 1, 23
	73	74° 46.5'	99° 21'	NS	1975	Sept.			bottom	Note 1, 23
	75	73° 03.8'	84° 22'	NS	1975	Sept.			bottom	Note 1, 23
	76	73° 04.2'	84° 21'	NS	1975	Sept.			bottom	Note 1, 23
	77	73° 04.7'	84° 20'	NS	1975	Sept.			bottom	Note 1, 23

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
75-0029 (cont'd)	78	73° 05.3'	84° 19'	NS	1975	Sept.		1	bottom	Note 1, 23
	79	73° 05.5'	84° 18'	NS	1975	Sept.		1	bottom	Note 1, 23
	80	73° 04.4'	84° 29'	NS	1975	Sept.		1	bottom	Note 1, 23
	81	73° 05.1'	84° 28.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	82	73° 05.5'	84° 28.0'	NS	1975	Sept.		1	bottom	Note 1, 23
	83	73° 06.0'	84° 27.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	84	73° 06.5'	84° 27.0'	NS	1975	Sept.		1	bottom	Note 1, 23
	85	73° 06.2'	84° 34'	NS	1975	Sept.		1	bottom	Note 1, 23
	86	73° 05.1'	84° 41'	NS	1975	Sept.		1	bottom	Note 1, 23
	87	73° 05.5'	84° 40.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	88	73° 06.0'	84° 39.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	89	73° 06.5'	84° 39'	NS	1975	Sept.		1	bottom	Note 1, 23
	90	73° 07.0'	84° 38.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	91	73° 07.4'	84° 38'	NS	1975	Sept.		1	bottom	Note 1, 23
	92	73° 07.0'	84° 43'	NS	1975	Sept.		1	bottom	Note 1, 23
	93	73° 06.6'	84° 48'	NS	1975	Sept.		1	bottom	Note 1, 23
	94	74° 06.4'	84° 52.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	95	73° 06.8'	84° 52'	NS	1975	Sept.		1	bottom	Note 1, 23
	96	73° 07.2'	84° 50.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	97	73° 07.5'	84° 49.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	98	73° 07.8'	84° 48.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	99	73° 08.7'	84° 47'	NS	1975	Sept.		1	bottom	Note 1, 23
	100	73° 09.5'	84° 55'	NS	1975	Sept.		1	bottom	Note 1, 23
	101	73° 09.0'	84° 57'	NS	1975	Sept.		1	bottom	Note 1, 23
	102	73° 09.3'	85° 08'	NS	1975	Sept.		1	bottom	Note 1, 23
	103	73° 10.1'	85° 07.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	104	73° 11.1'	85° 07'	NS	1975	Sept.		1	bottom	Note 1, 23
	105	73° 12.4'	85° 06'	NS	1975	Sept.		1	bottom	Note 1, 23
	106	73° 03.6'	84° 09'	NS	1975	Sept.		1	bottom	Note 1, 23
	107	73° 03.1'	84° 03'	NS	1975	Sept.		1	bottom	Note 1, 23
	108	73° 03.2'	83° 56.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	109	73° 02.3'	83° 50'	NS	1975	Sept.		1	bottom	Note 1, 23
	110	73° 01.0'	83° 45'	NS	1975	Sept.		1	bottom	Note 1, 23
	111	73° 00.0'	83° 39'	NS	1975	Sept.		1	bottom	Note 1, 23
	113	74° 56.5'	99° 00'	NS	1975	Sept.		1	bottom	Note 1, 23
	114	74° 56.5'	98° 54'	NS	1975	Sept.		1	bottom	Note 1, 23
	115	74° 54.5'	98° 54'	NS	1975	Sept.		1	bottom	Note 1, 23
	116	74° 52.0'	98° 54.5'	NS	1975	Sept.		1	bottom	Note 1, 23
	117	74° 52.2'	99° 00'	NS	1975	Sept.		1	bottom	Note 1, 23
	118	74° 54.5'	99° 00'	NS	1975	Sept.		1	Bottom	Note 1, 23

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
75-0029 (cont'd)	120	74° 50.0'	99° 44'	NS	1975	Sept.		1	bottom	Note 1, 23
	122	74° 55.0'	99° 44'	NS	1975	Sept.		1	bottom	Note 1, 23
	126	74° 51.0'	98° 38'	NS	1975	Sept.		1	bottom	Note 1, 23
	127	74° 56.0'	98° 22'	NS	1975	Sept.		1	bottom	Note 1, 23
75-0030	1	72° 57'	84° 31.8'	NS	1975	Aug.	NS	1	bottom	Note 1
	2	72° 58'	84° 33'	NS	1975	Aug.	NS	1	bottom	Note 1
	3	72° 58'	84° 33.8'	NS	1975	Aug.	NS	1	bottom	Note 1
	4	72° 57.3'	84° 34.1'	NS	1975	Aug.	NS	1	bottom	Note 1
	5	72° 57'	84° 30'	NS	1975	Aug.	NS	1	bottom	Note 1
	6	72° 59'	84° 28.9'	NS	1975	Aug.	NS	1	bottom	Note 1
	7	73° 00.5'	84° 28'	NS	1975	Aug.	NS	1	bottom	Note 1
	8	73° 03'	84° 26.8'	NS	1975	Aug.	NS	1	bottom	Note 1
	9	72° 58'	84° 31.2'	NS	1975	Aug.	NS	1	bottom	Note 1
	10	73° 00'	84° 33.5'	NS	1975	Aug.	NS	1	bottom	Note 1
	11	73° 06'	84° 36'	NS	1975	Aug.	NS	1	bottom	Note 1
	12	73° 01'	84° 37'	NS	1975	Aug.	NS	1	bottom	Note 1
	13	73° 02.5'	84° 39.5'	NS	1975	Aug.	NS	1	bottom	Note 1
	14	73° 02.5'	84° 45'	NS	1975	Aug.	NS	1	bottom	Note 1
75-0031 A	1	72° 57'	84° 31.8'	NS	1975	Sept.	NS	1	bottom	Note 1
	2	72° 58'	84° 33'	NS	1975	Sept.	NS	1	bottom	Note 1
	4	72° 57.3'	84° 34.1'	NS	1975	Sept.	NS	1	bottom	Note 1
	5	72° 57'	84° 30'	NS	1975	Sept.	NS	1	bottom	Note 1
	7	73° 00.5'	84° 28'	NS	1975	Sept.	NS	1	bottom	Note 1
	8	73° 03'	84° 26.8'	NS	NS	Sept.	NS	1	bottom	Note 1
	Biota-3	72° 56.5'	84° 33.5'	NS	1975	Aug.	NS	1	bottom	Note 24
	Hole	73° 07.5'	84° 51.5'	300	1975	Mar.	NS	NS	NS	Note 1, 25
75-0031 B	In-sill	73° 17.5'	84° 57.0'	220	1975	Mar.	NS	NS	NS	Note 1, 25
	Out-sill	73° 19'	85° 00'	280	1975	Mar.	NS	NS	NS	Note 1, 25
	Hole	73° 07.5'	84° 51.5'	300	1975	June	NS	11	275	Note 1
	Hole	73° 07.5'	84° 51.5'	300	1975	Aug.	NS	6	275	Note 1
	In-sill	73° 17.5'	84° 57.0'	220	1975	Aug.	NS	8	185	Note 1
	Out-sill	73° 19'	85° 00'	280	1975	Aug.	NS	10	275	Note 1
	Strathcona Sound	73° 11'	85° 10'	NS	1975	Aug.	NS		120-300	Note 26, 27
	Strathcona Sound	73° 11'	85° 10'	NS	1975	Aug.	NS		285-290	Note 26, 28
	Strathcona Sound	73° 11'	85° 10'	NS	1975	Aug.	NS			
	Strathcona Sound	73° 11'	85° 10'	NS	1975	Aug.	NS			

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
75-0031 B (cont'd)	Strathcona Sound	73° 11'	85° 10'	NS	1975	Aug.	NS	NS	175-180	Note 26, 29
	Strathcona Sound	73° 11'	85° 10'	NS	1975	Aug.	NS	NS	285-290	Note 26, 30
	0.1	72° 55'	84° 26.5'	NS	1975	Aug.	NS	NS	1-2	Note 1, 31
	1	72° 57'	84° 33.5'	NS	1975	Aug.	NS	NS	1-2	Note 1, 32
	2	72° 58'	84° 38'	NS	1975	Aug.	NS	NS	1-2	Note 1, 33
	7	73° 00'	84° 45.5'	NS	1975	Aug.	NS	NS	1-2	Note 1, 34
	15	73° 18'	84° 52'	NS	1975	Aug.	NS	NS	1-2	Note 1, 35
	18	73° 11.5'	85° 03'	NS	1975	Aug.	NS	NS	1-2	Note 1, 36
	Aston Bay	73° 40'	95° 00'	NS	1975	NS	NS	NS		Note 1, 37
	Aston Bay	73° 40'	95° 00'	NS	1975	NS	NS	NS		Note 1, 38
75-0033										
76-0007	2	69° 54'	84° 16'	164	1976	May	21	1457	10	160
76-0008	EM	74° 06'	81° 30'	786	1976	July	22	NS	5	150
	EM	74° 06'	81° 30'	786	1976	Aug.	5	NS	6	300
	EM	74° 06'	81° 30'	786	1976	Aug.	19	NS	16	700
	EM	74° 06'	81° 30'	786	1976	Aug.	29	NS	6	300
	EM	74° 06'	81° 30'	786	1976	Sept.	11	NS	300	300
	CS	74° 32'	80° 20'	668	1976	July	23	NS	150	150
	CS	74° 32'	80° 20'	668	1976	Aug.	4	NS	6	300
	CSA	74° 25'	80° 18'	722	1976	Aug.	16	NS	16	675
	CS	74° 32'	80° 20'	668	1976	Aug.	28	NS	6	300
	CS	74° 32'	80° 20'	668	1976	Sept.	8	NS	6	300
	CW	74° 27'	82° 03'	686	1976	July	24	NS	5	150
	CW	74° 27'	82° 03'	686	1976	Aug.	3	NS	6	300
	CW	74° 27'	82° 03'	686	1976	Aug.	17	NS	16	650
	CW	74° 27'	82° 03'	686	1976	Aug.	27	NS	6	300
	CW	74° 27'	82° 03'	686	1976	Sept.	7	NS	6	300
	NB	73° 43'	81° 02'	503	1976	July	26	NS	5	150
	NB	73° 43'	81° 02'	503	1976	Aug.	7	NS	6	300
	NB	73° 43'	81° 02'	503	1976	Aug.	20-21	NS	14	300
	NB	73° 43'	81° 02'	503	1976	Aug.	31	NS	6	370
	NB	73° 43'	81° 02'	503	1976	Sept.	12	NS	6	300

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
76-0008 (cont'd)	MM	74° 07'	82° 37'	741	1976	July	27	NS	5	150
	MM	74° 07'	82° 37'	741	1976	Aug.	6	NS	6	300
	MM	74° 07'	82° 37'	741	1976	Aug.	18	NS	16	650
	MM	74° 07'	82° 37'	741	1976	Aug.	29	NS	6	300
	MM	74° 07'	82° 37'	741	1976	Sept.	11	NS	6	300
	MM	74° 07'	82° 37'	741	1976	Sept.	28	NS	5	150
	WM	74° 12'	87° 57'	430	1976	July	8	NS	6	300
	WM	74° 12'	87° 57'	430	1976	Aug.	22	NS	7	400
	WM	74° 12'	87° 57'	430	1976	Aug.	1	NS	6	300
	WM	74° 12'	87° 57'	430	1976	Sept.	13	NS	6	300
76-0009	Cunningham Inlet	74° 07'	93° 50'	NS	1976	July		NS	NS	Note 1, 39
	X	74° 08.5'	93° 53'	NS	1976	July	21	NS	NS	Note 1
	X	74° 08.5'	93° 53'	NS	1976	July	26	NS	4	Note 1
	Y	74° 08.3'	93° 52.5'	NS	1976	July	21	NS	NS	Note 1
	Y	74° 08.3'	93° 52.5'	NS	1976	July	26	NS	3	Note 1
	Z	74° 08.1'	93° 52'	NS	1976	July	21	NS	NS	Note 1
	Z	74° 08.1'	93° 52'	NS	1976	July	26	NS	2	Note 1
	Delta	74° 03'	93° 47.5'	NS	1976	July	30	NS	2	Note 1
	1.1 km north of Delta	74° 05.6'	93° 47.5'	NS	1976	July	30	NS	30	Note 1
	Head	74° 05.4'	93° 47.5'	NS	1976	Aug.	29	NS	NS	Note 1
76-0010	Entrance	74° 08.2'	93° 52.1'	NS	1976	Aug.	29	NS	NS	Note 1
	I	74° 40.5'	95° 10'	100	1976	July	5	NS	4	Note 1
	I	74° 23.8'	82° 03.5'	664	1976	Aug.	16	1622	15	Note 40
	3	74° 07.0'	81° 58.0'	730	1976	Aug.	16	2125	12	700
	4	73° 49.7'	81° 59.2'	660	1976	Aug.	17	0343	11	620
	7	74° 30.0'	90° 58.0'	183	1976	Aug.	18	0118	7	160
	9	74° 16.5'	90° 39.5'	223	1976	Aug.	18	0600	7	200
	34	73° 04.4'	84° 21.0'	NS	1976	Sept.	1	0255	1	bottom
	35	73° 03.9'	84° 32.3'	NS	1976	Sept.	1	0638	1	bottom
	36	73° 08.9'	84° 56.9'	NS	1976	Sept.	1	0740	1	bottom

DATA SET NUMBER	STATION NUMBER	LATITUDE	LONGITUDE	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
76-0011 B (cont'd)	37	73° 12.6'	85° 09.2'	NS	1976	Sept.	1	1	bottom	
	38	73° 11.6'	85° 10.3'	NS	1976	Sept.	1	1	bottom	
	39	73° 10.4'	85° 02.8'	NS	1976	Sept.	1	1	bottom	
	40	73° 07.5'	84° 51.5'	NS	1976	Sept.	1	1	bottom	
	41	72° 59.0'	83° 39.6'	NS	1976	Sept.	1	1	bottom	
	42	73° 01.7'	83° 48.1'	NS	1976	Sept.	1	1	bottom	
76-0012	Arctic Bay	72° 50'	85° 10'	NS	1976	Aug.	NS	1	bottom	Note 1
	Arctic Bay	72° 50'	85° 10'	NS	1976	Aug.	NS	NS	NS	Note 1, 41
	Arctic Bay	72° 50'	85° 10'	NS	1976	Aug.	NS	NS	NS	Note 1, 42
	3	72° 56.5'	84° 33.5'	NS	1976	May	NS	NS	NS	Note 1, 43
	3	72° 56.5'	84° 33.5'	NS	1976	Aug.	NS	NS	NS	Note 1, 44
	3	72° 56.5'	84° 33.5'	NS	1976	Aug.	NS	NS	NS	Note 1, 45
76-0021	1	74° 12'	87° 56'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 46
	1	74° 12'	87° 56'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 47
	1	74° 12'	87° 56'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 48
	2	74° 30.5'	82° 30'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 49
	3	74° 27'	82° 03'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 50
	4	74° 06.5'	82° 36'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 51
76-0022	1	74° 06.5'	82° 36'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 52
	4	74° 06.5'	82° 36'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 53
	5	74° 06'	81° 30'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 54
	6	73° 45'	80° 57'	NS	1976	July-Aug.	NS	NS	NS	Note 1, 55
	Barrow Strait	74° 30'	95° 00'	NS	1976	NS	NS	NS	NS	Note 1, 56
	Barrow Strait	74° 30'	95° 00'	NS	1976	NS	NS	NS	NS	Note 1, 57
77-0014	Arctic Bay	72° 50'	84° 10'	NS	1976	NS	NS	NS	NS	
	30	73° 47.1'	81° 57.7'	483	1977	Sept.	8	18	bottom	
	31	73° 55.0'	82° 00.0'	655	1977	Sept.	8	17	bottom	
	32	74° 09.4'	81° 58.9'	785	1977	Sept.	8	28	764	
	33	74° 36.6'	94° 20.4'	106	1977	Sept.	9	8	101	
	34	74° 27.7'	94° 16.2'	155	1977	Sept.	9	11	bottom	
76-0022	35	74° 20.0'	94° 15.8'	174	1977	Sept.	10	18	bottom	
	36	74° 10.7'	94° 14.6'	168	1977	Sept.	10	21	bottom	
	37	73° 58.7'	90° 22.9'	110	1977	Sept.	10	8	106	
	38	74° 07.1'	90° 22.6'	214	1977	Sept.	10	13	bottom	

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
77-0014 (cont'd)	39	74° 15.3'	90° 25.9'	232	1977	Sept.	10	14	1700	
	40	74° 24.1'	90° 22.2'	232	1977	Sept.	10	21	2000	bottom
	41	74° 33.0'	90° 22.8'	176	1977	Sept.	10	12	2300	bottom
	42	74° 10.9'	85° 58.5'	520	1977	Sept.	11	35	0600	bottom
	43	74° 18.0'	81° 57.7'	732	1977	Sept.	11	20	1550	bottom
	44	74° 26.3'	81° 58.3'	750	1977	Sept.	11	17	2300	bottom
77-0015 A	3	71° 50.5'	94° 29.0'	NS	1977	May	23	5	NS	Note 1
	3	71° 50.5'	94° 29.0'	NS	1977	May	25	5	NS	Note 1
	3	71° 50.5'	94° 29.0'	NS	1977	May	27	5	NS	Note 1
77-0015 B	137	71° 46.2'	94° 36'	NS	1977	Sept.	2	1	NS	Note 1
	138	71° 46.2'	94° 35.5'	NS	1977	Sept.	2	2	NS	Note 1
	139	71° 46.2'	94° 33'	NS	1977	Sept.	2	5	NS	Note 1
77-0016 A	3	75° 02.1'	108° 45'	NS	1977	June	9	5	NS	Note 1
	2	75° 01.3'	108° 44.5'	NS	1977	June	13	5	NS	Note 1
	6	75° 01.4'	108° 46.5'	NS	1977	June	14	4	NS	Note 1
77-0016 B	1	75° 01.9'	108° 35.8'	NS	1977	Aug.	5	3	NS	Note 1
	6	75° 01.4'	108° 36.5'	NS	1977	Aug.	5	4	NS	Note 1
	2	75° 01.3'	108° 44.5'	NS	1977	Aug.	6	5	NS	Note 1
	4	75° 00'	108° 54'	NS	1977	Aug.	7	5	NS	Note 1
	7	75° 04.6'	108° 49'	NS	1977	Aug.	8	4	NS	Note 1
	8	75° 04.1'	108° 44'	NS	1977	Aug.	9	3	NS	Note 1
	5	74° 58.7'	108° 48.4'	NS	1977	Aug.	11	5	NS	Note 1
	2	75° 01.3'	108° 44.5'	NS	1977	Aug.	27	5	NS	Note 1
	5	74° 58.7'	108° 48.4'	NS	1977	Aug.	28	3	NS	Note 1
	T0.0	75° 04.5'	108° 48.5'	NS	1977	Aug.	15	1	NS	Note 1
	T0.5	75° 04'	108° 49'	NS	1977	Aug.	15	1	NS	Note 1
	T1.0	75° 03.5'	108° 49.5'	NS	1977	Aug.	15	1	NS	Note 1
	T1.5	75° 03.1'	108° 49.9'	NS	1977	Aug.	15	1	NS	Note 1
	T2.0	75° 02.5'	108° 50.1'	NS	1977	Aug.	15	1	NS	Note 1
	T2.5	75° 02.1	108° 50.5'	NS	1977	Aug.	15	1	NS	Note 1

DATA SET NUMBER	STATION NUMBER	LATITUDE "N	LONGITUDE "W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
77-0016 B (cont'd)	T3.0	73° 01.6'	108° 51'	NS	1977	Aug.	13	1	0	Note 1
	T3.5	73° 01'	108° 52'	NS	1977	Aug.	13	1	0	Note 1
	T4.0	73° 00'	108° 53'	NS	1977	Aug.	13	1	0	Note 1
	T4.5	74° 59.5'	108° 53.5'	NS	1977	Aug.	13	1	0	Note 1
	9	73° 04.6'	108° 49.9'	NS	1977	Aug.	26	9	140	Note 1
78-0006	Bridport Inlet	73° 03.2'	108° 52.5'	NS	1978	Aug.	22	6	110	Note 1
78-0020	79	73° 47.2'	79° 59.3'	768	1978	Sept.	10	18	750	
	80	74° 01.7'	80° 00.9'	786	1978	Sept.	10	15	700	
	81	74° 17.0'	80° 02.4'	693	1978	Sept.	11	15	670	
	82	74° 31.9'	80° 00.6'	556	1978	Sept.	11	13	550	
78-0021	1	74° 36.5'	81° 33'	NS	1978		NS	1	0	Note 1, 58
	2	74° 30'	81° 32.5'	NS	1978		NS	1	0	Note 1, 58
	3	74° 30'	81° 29.5'	NS	1978		NS	1	0	Note 1, 58
	4	74° 29.2'	81° 28'	NS	1978		NS	1	0	Note 1, 58
	5	74° 33.9'	81° 11.5'	NS	1978		NS	1	0	Note 1, 58
	6	74° 33.4'	81° 13'	NS	1978		NS	1	0	Note 1, 58
	7	74° 32.9'	81° 12.5'	NS	1978		NS	1	0	Note 1, 58
	8	74° 32.5'	81° 11.5'	NS	1978		NS	1	0	Note 1, 58
	9	74° 31.4'	81° 12.5'	NS	1978		NS	1	0	Note 1, 58
	10	74° 30.9'	81° 10'	NS	1978		NS	1	0	Note 1, 58
	11	74° 30.8'	81° 08'	NS	1978		NS	1	0	Note 1, 58
	12	74° 33.7'	81° 02'	NS	1978		NS	1	0	Note 1, 58
	13	74° 33.9'	80° 59'	NS	1978		NS	1	0	Note 1, 58
	14	74° 32.8'	80° 51'	NS	1978		NS	1	0	Note 1, 58
	15	74° 33.5'	80° 48'	NS	1978		NS	1	0	Note 1, 58
	16	74° 33'	80° 48'	NS	1978		NS	1	0	Note 1, 58
78-0022 A	C-1	73° 48.2'	80° 11'	825	1978	July	31	6	50	
	C-2	73° 48.2'	80° 11'	825	1978	Aug.	5	9	700	
	C-3	73° 59.1'	80° 15'	786	1978	Aug.	5	15	700	
	C-4	74° 09.8'	80° 18'	786	1978	Aug.	4	14	600	
	C-5	74° 19.7'	80° 22'	702	1978	Aug.	4	14	600	
		74° 29.8'	80° 26'	650	1978	Aug.	4	14	600	

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
78-0022 A (cont'd)	B-6	74° 21.8'	82° 00'	673	1978	July	28	6	50	Note 59
	EM	74° 06.0'	81° 30'	750	1978	July	28	6	50	
	PG-1	74° 48'	79° 40'	NS	1978	Aug.	3	3	25	Note 60
	PG-2	74° 48'	79° 39'	NS	1978	Aug.	3	3	25	
	PG-3	74° 48'	79° 39'	NS	1978	Aug.	3	3	25	
	PG-4	74° 45'	80° 10'	NS	1978	Aug.	3	3	25	
	PG-5	74° 45'	80° 09'	NS	1978	Aug.	3	3	25	
	PG-6	74° 45'	80° 08'	NS	1978	Aug.	3	3	25	
	PG-7	74° 37'	80° 15'	NS	1978	Aug.	3	2	25	
	PG-8	74° 37'	80° 14'	NS	1978	Aug.	3	2	10	
	PG-9	74° 37'	80° 13'	NS	1978	Aug.	3	3	25	
	CS-1	74° 35'	80° 25'	12	1978	Aug.	4	2	10	
78-0022 B	CS-2	74° 34.5'	80° 25'	16	1978	Aug.	4	2	10	Note 60
	CS-3	74° 34'	80° 25'	87	1978	Aug.	4	3	25	
	C-1	73° 48.2'	80° 11'	825	1978	Sept.	6	6	50	
	C-2	73° 59.1'	80° 15'	786	1978	Sept.	6	6	50	
	C-3	74° 09.8'	80° 18'	786	1978	Sept.	6	6	50	
	C-4	74° 19.7'	80° 22'	702	1978	Sept.	6	6	50	
	C-5	74° 29.8'	80° 26'	650	1978	Sept.	6	6	50	
	CW	74° 27.0'	82° 03'	768	1978	Sept.	21	6	50	
	CW	74° 27.0'	82° 03'	768	1978	Sept.	7	6	50	
	EM	74° 06.0'	81° 30'	750	1978	Sept.	7	6	50	
78-0022 C	NB	73° 43.0'	81° 02'	547	1978	Aug.	19	6	50	Note 60
	NB	73° 43.0'	81° 02'	547	1978	Aug.	19	6	50	
	PH-1	74° 33.9'	81° 17'	20	1978	Sept.	8	6	50	
	PH-2	74° 33.3'	81° 16'	72	1978	Aug.	28	5	15	
	PH-3	74° 31.5'	81° 02'	46	1978	Aug.	28	7	50	
	Cape Hay	73° 45'	80° 00'	NS	1978	Aug.	29	8	50	
					1978	Sept.	3-4	1	0	
	C-1	73° 48.2'	80° 11'	825	1978	Sept.	20	6	50	
	C-2	73° 59.1'	80° 15'	786	1978	Sept.	20	6	50	
	C-3	74° 09.8'	80° 18'	786	1978	Sept.	20	6	50	
78-0022 D	C-4	74° 19.7'	80° 22'	702	1978	Sept.	20	6	50	Note 60
	C-5	74° 29.8'	80° 26'	650	1978	Sept.	20	6	50	
	CW	74° 27.0'	82° 03'	768	1978	Sept.	19	6	50	
	NB	73° 43.0'	81° 02'	547	1978	Sept.	22	6	50	
					1978	Sept.	22	6	50	


DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
79-0014	1-79	75° 0.8'	108° 48.0'	141	1979	March	13	6	110	Note 1, 61
	2-79	74° 58.8'	108° 53.0'	87	1979	March	20	5	75	Note 1, 62
79-0024	1	73° 08.5'	84° 58.5'	NS	1979	Aug.	NS	1	bottom	Note 1, 63
	1	73° 08.5'	84° 58.5'	NS	1979	Aug.	NS			Note 1, 64
	1	73° 08.5'	84° 58.5'	NS	1979	Aug.	NS			Note 1, 65
	1	73° 08.5'	84° 58.5'	NS	1979	Aug.	NS			Note 1, 66
	1	73° 08.5'	84° 58.5'	NS	1979	Aug.	NS			Note 1, 67
	2	72° 59'	84° 41.5'	NS	1979	Aug.	NS	1	bottom	Note 1, 68
	2	72° 59'	84° 41.5'	NS	1979	Aug.	NS			Note 1, 69
	2	72° 59'	84° 41.5'	NS	1979	Aug.	NS			Note 1, 70
	2	72° 59'	84° 41.5'	NS	1979	Aug.	NS			Note 1, 71
	3	72° 56.5'	84° 33.5'	NS	1979	Aug.	NS	1	bottom	Note 1, 72
	3	72° 56.5'	84° 33.5'	NS	1979	Aug.	NS			Note 1, 73
	3	72° 56.5'	84° 33.5'	NS	1979	Aug.	NS			Note 1, 74
	3	72° 56.5'	84° 33.5'	NS	1979	Aug.	NS			Note 1, 75
	4	72° 55'	84° 24.5'	NS	1979	Aug.	NS	1	bottom	Note 1, 76
	4	72° 55'	84° 24.5'	NS	1979	Aug.	NS			Note 1, 77
	4	72° 55.5'	84° 03.5'	NS	1979	Aug.	NS	1	bottom	Note 1, 78
	5	72° 55.5'	84° 03.5'	NS	1979	Aug.	NS			Note 1, 79
	5	72° 55.5'	84° 03.5'	NS	1979	Aug.	NS			Note 1, 80
	5	72° 55.5'	84° 03.5'	NS	1979	Aug.	NS			Note 1, 81
	6	72° 58'	84° 19.5'	NS	1979	Aug.	NS	1	bottom	Note 1, 82
	6	72° 58'	84° 19.5'	NS	1979	Aug.	NS			Note 1, 83
	7	73° 09'	84° 40'	NS	1979	Aug.	NS			Note 1, 84
	7	73° 09'	84° 40'	NS	1979	Aug.	NS			
	7	73° 09'	84° 40'	NS	1979	Aug.	NS			
	7	73° 09'	84° 40'	NS	1979	Aug.	NS			


DATA SET NUMBER	STATION NUMBER	LATITUDE "N	LONGITUDE "W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS
					YEAR	MONTH	DAY			
79-0024 (cont'd)	8	73° 16'	84° 51'	NS	1979	Aug.	NS	1	bottom	Note 1, 85
	8	73° 16'	84° 51'	NS	1979	Aug.	NS			Note 1, 86
	8	73° 16'	84° 51'	NS	1979	Aug.	NS			
80-0007	50	73° 44.7'	81° 46.5'	240	1980	Aug.	5	8	230	
	51	73° 47.0'	81° 45.7'	570	1980	Aug.	5	8	570	
	52	73° 52.6'	81° 47.5'	622	1980	Aug.	5	8	622	
	53	73° 58.0'	81° 46.0'	650	1980	Aug.	5	8	650	
	54	74° 06.0'	81° 46.0'	713	1980	Aug.	5	8	400	
	55	74° 11.0'	81° 46.5'	732	1980	Aug.	5	8	725	
	56	74° 19.7'	81° 46.5'	680	1980	Aug.	6	5	600	
	57	74° 24.3'	81° 46.5'	660	1980	Aug.	6	8	710	
	58	74° 27.0'	81° 45.0'	720	1980	Aug.	6	8	710	
	60	73° 53.6'	86° 00.0'	238	1980	Aug.	6	8	235	
	61	74° 01.8'	86° 00.0'	420	1980	Aug.	7	8	420	
	62	74° 06.9'	86° 00.0'	510	1980	Aug.	7	8	500	
	63	74° 11.3'	86° 00.0'	515	1980	Aug.	7	8	515	
	64	74° 16.3'	86° 00.0'	530	1980	Aug.	7	8	530	
	65	74° 21.0'	86° 00.0'	550	1980	Aug.	7	8	550	
	66	74° 28.0'	85° 56.5'	219	1980	Aug.	6	8	210	
	67	74° 26.3'	86° 00.3'	503	1980	Aug.	6	8	500	
	68	74° 31.0'	89° 45.0'	215	1980	Aug.	8	8	215	
	69	74° 23.9'	89° 44.3'	300	1980	Aug.	8	8	295	
	70	74° 20.8'	89° 45.0'	280	1980	Aug.	8	8	280	
	71	74° 17.0'	89° 47.0'	270	1980	Aug.	8	8	210	
	72	74° 11.0'	89° 43.0'	220	1980	Aug.	8	8	200	
	73	74° 06.0'	89° 44.0'	200	1980	Aug.	8	8	200	
	74	74° 01.0'	89° 44.0'	235	1980	Aug.	8	8	235	
	75	73° 55.4'	89° 43.7'	285	1980	Aug.	8	8	280	
	81	73° 48.3'	81° 05.5'	620	1980	Aug.	12	8	620	
	82	74° 37.9'	93° 39.0'	95	1980	Aug.	13	8	85	
	83	74° 33.0'	93° 39.0'	135	1980	Aug.	13	8	135	
	84	74° 28.0'	93° 39.0'	160	1980	Aug.	13	8	160	
	85	74° 22.6'	93° 39.0'	160	1980	Aug.	13	8	160	
	86	74° 18.0'	93° 39.7'	160	1980	Aug.	13	8	160	
	87	74° 12.8'	93° 31.0'	140	1980	Aug.	14	8	140	

DATA SET NUMBER	STATION NUMBER	LATITUDE °N	LONGITUDE °W	STATION DEPTH (m)	TIME SAMPLED			NUMBER POINTS	MAXIMUM DEPTH SAMPLED (m)	REMARKS		
					YEAR	MONTH	DAY					
80-0007 (cont'd)	88	74° 40.2'	93° 26.0'	85	1980	Aug.	14	0728	8	70		
	89	74° 40.8'	93° 14.0'	135	1980	Aug.	14	0814	8	125		
	90	74° 41.0'	92° 55.0'	190	1980	Aug.	14	0906	8	140		
	91	74° 36.0'	92° 50.5'	145	1980	Aug.	14	1005	8	140		
	92	74° 29.2'	92° 06.8'	190	1980	Aug.	14	1212	8	185		
	93	74° 41.0'	91° 51.0'	NS	1980	Aug.	14	1528	8	95		
	100	73° 59.5'	89° 42.5'	250	1980	Aug.	15	0619	8	250		
	101	73° 53.6'	89° 44.0'	295	1980	Aug.	15	0726	8	295		
	102	73° 48.4'	89° 40.5'	310	1980	Aug.	23	0815	8	310		
	103	73° 37.6'	89° 44.0'	330	1980	Aug.	15	1022	8	330		
	104	73° 32.6'	89° 47.0'	365	1980	Aug.	15	1358	8	365		
	105	73° 25.8'	89° 44.0'	400	1980	Aug.	15	1530	8	435		
	106	73° 25.8'	89° 44.0'	400	1980	Aug.	15	1808	8	435		
	107	73° 21.1'	89° 43.0'	430	1980	Aug.	15	2014	8	420		
	108	73° 29.8'	88° 31.0'	128	1980	Aug.	15	2356	8	125		
80-0009	109	73° 32.7'	88° 47.3'	310	1980	Aug.	16	0045	8	310		
	110	73° 36.0'	89° 03.0'	384	1980	Aug.	16	0218	8	375		
	111	73° 39.0'	89° 18.0'	420	1980	Aug.	16	0322	8	420		
	112	73° 42.2'	89° 34.0'	365	1980	Aug.	16	0429	8	365		
	113	73° 45.1'	89° 50.5'	310	1980	Aug.	16	0532	8	310		
	114	73° 48.2'	90° 05.5'	295	1980	Aug.	16	0640	8	285		
	115	73° 49.0'	90° 11.0'	200	1980	Aug.	16	0736	8	200		
	80-1	75° 02.93'	108° 48.67'	NS	1980	April	6	NS	3	30	Note 87 Note 88	
	80-2	75° 00.85'	108° 48.50'	135-140	1980	April	7	NS	6	100		
	80-4	74° 58.13'	108° 53.03'	63.5	1980	April	9	0200	4	50		
	80-0018	2	72° 00.0'	96° 00.0'		1980	NS	NS	NS	7		
		4	74° 00.0'	83° 00.0'		1980	NS	NS	NS	27		
	81-0007	81046	74° 13.0'	93° 47.8'	162.6	1981	Mar.	31	2036	NS	NS	
		81084	74° 41.6'	96° 00.0'	241.0	1981	Apr.	1	1623	NS	NS	
		81081	73° 41.6'	96° 49.0'	225.5	1981	Apr.	1	1843	NS	NS	
81016		74° 12.3'	97° 49.3'	174.5	1981	Apr.	19	1650	NS	NS		


10. MAPS

- 10.1 Geographical occurrence of chemical oceanographic data on a cumulative bi-monthly basis.







The locations of stations reporting chemical oceanographic data, all years.




The locations of stations reporting chemical oceanographic data collected during January - February, all years.




The locations of stations reporting chemical oceanographic data collected during March - April, all years.




The locations of stations reporting chemical oceanographic data collected during May - June, all years.




The locations of stations reporting chemical oceanographic data collected during July - August, all years.




The locations of stations reporting chemical oceanographic data collected during September - October, all years.





The locations of stations reporting chemical oceanographic data collected during November - December, all years.





- 10.2 Geographical occurrence of chemical
oceanographic data on a yearly basis.




































































































11. INDICES

**11.1 Alphabetical Listing of Data Set
Occurrence by Geographic Area**

11.2 Index of References by Data Set Number



11.2 Index of References by Data Set Number
(Secondary references are indented.)

- 28-0001 Riis-Carstensen, E., 1936. The Godthaab Expedition 1928. The hydrographic work and material. Meddeleser om Gronland, Bd. 78, Nr. 3, 101 pp. + plates.

Kiilerich, A.B., 1939. The Godthaab Expedition 1928. A theoretical treatment of the hydrographical observation material. Meddeleser om Gronland, Bd. 78, Nr. 5, 146 pp. + plates.

- 50-0002 NODC No. 31ED50375

- 52-0003 NODC No. 31E050368

- 53-0001 NODC No. 31BI50430

- 54-0002 NODC No. 31BI50456

- 54-0003 NODC No. 31NW50457

- 56-0002 Apollonio, S., 1957. Plankton productivity studies in Allen Bay, Cornwallis Island, N.W.T., 1956. Woods Hole Oceanographic Institution and Osborn Zoological Laboratory, Yale University, 8 pp. + tables.

Apollonio, S., (no date). Phytoplankton production in the Canadian Arctic. Unpubl. manuscript. The Arctic Institution of North America, 38 pp. + figures.

Apollonio, S., 1976. Primary production in the Canadian Arctic Archipelago, 1961-1963. Unpubl. manuscript. Bigelow Laboratory for Ocean Sciences Contrib. No. 76016 - 46 pp.

- 56-0003 Collin, A.E., 1958. An oceanographic study of Prince Regent Inlet, the Gulf of Boothia and adjacent waters. Atlantic Oceanographic Group, Fisheries and Research Board of Canada, Manuscript Report No. 8, 43 pp. + figures.

Collin, A.E., 1962. The waters of the Canadian Arctic Archipelago. Bedford Institute of Oceanography (unpublished manuscript), 9 pp.

Collin, A.E., 1962. The waters of the Canadian Arctic Archipelago. Proceedings of the Arctic Basin Symposium, pp. 128-136.

Collin, A.E., 1962. The oceanography of Lancaster Sound. Ph.D. Thesis. McGill University, 204 pp.

- 57-0003 Collin, A.E., 1962. The oceanography of Lancaster Sound. Ph.D. Thesis. McGill University, 204 pp.

NODC No. 18LA50573

Collin, A.E., 1962. The waters of the Canadian Arctic Archipelago. Bedford Institute of Oceanography (unpublished manuscript), 9 pp.

Collin, A.E., 1962. The waters of the Canadian Arctic Archipelago. Proceedings of the Arctic Basin Symposium, pp. 128-136.

- 60-0008** Buckley, D.E., 1963. Recent marine sediments of Lancaster Sound, District of Franklin. M.Sc. Thesis. University of Western Ontario, London, Canada, 112 pp.
- Buckley, D.E., 1971. Recent marine sediments of Lancaster Sound, District of Franklin. Maritime Sediments, Vol. 7(3): 66-117.
- 61-0004** MEDS No. 181061341, NODC No. 18LA50128; CODC (1966) No. 12.
- 62-0005A** CODC (1963) No. 4 C.R.N. 363; NODC No. 18-50003
- 62-0005B** CODC (1963) No. 4 C.R.N. 366; NODC No. 18-50008
- 62-0005C** CODC (1963) No. 4 C.R.N. 367; NODC No. 18-50009
- 62-0006** NODC No. 18MC50142.
- 62-0009** Bursa, A.S., 1968. Starch in the oceans. J. Fish. Res. Bd. Canada, 25(6): pp. 1269-1284.
- 65-0002** MEDS No. 180465001; NODC No. 18SL50278.
- 66-0004** NODC No. 31ED50807
- 66-0005** MEDS No. 180466001; NODC No. 18SL50280.
- 67-0001A** MEDS No. 180467001; NODC No. 18SL50117
- 67-0001B** MEDS No. 180467003; NODC No. 18IC50115
- 68-0001** Herlinveaux, R.H., 1970. Ice pack 8/68 - oceanographic and biological observations. Fisheries Research Board of Canada, Technical Report No. 159, 60 pp.
- NODC No. 18LA51061.
- 70-0010** Bowes, G.W. and C.J. Jonkel, 1975. Presence and distribution of polychlorinated biphenyls (PCB) in arctic and subarctic marine food chains. J. Fish. Res. Bd. Canada, 32: 2111-2123.
- 71-0009** See 70-0010
- 71-0016** Welch, H.E. and J. Kalff, 1975. Marine metabolism at Resolute Bay, N.W.T. Proceedings of the Circumpolar Conference on Northern Ecology; Sept. 15-18, 1975, II 69-75.

- 72-0011** Herlinveaux, R.H. and S.E.G. Wilson, 1974. Oceanographic observations in the Canadian Arctic, September, 1972. Marine Sciences Directorate, Pacific Region (unpublished manuscript), 11 pp.
- 72-0016** See **70-0010**
- 72-0017** Levy, E.M. and J.D. Moffat, 1975. Floating petroleum residues in the North Atlantic (1971-1974). Bedford Institute of Oceanography, 56 pp.
- Levy, E.M. and A. Walton, 1976. High seas oil pollution: particulate petroleum residues in the North Atlantic, J. Fish. Res. Board Canada, 33: 2781-2791.
- 73-0005** NODC No. 32SL58365.
- Codispoti, L.A. and T.G. Owens, 1975. Nutrient transports through Lancaster Sound in relation to the Arctic Ocean's reactive silicate budget and the outflow of Bering Strait waters. Department of Oceanography, University of Washington, Contribution No. 783.
- Codispoti, L.A., 1979. Arctic Ocean processes in relation to the dissolved silicon content of the Atlantic. Marine Science Communications, 5(6), pp. 361-381.
- 73-0006** Herlinveaux, R.H., D.B. Fissel and S.E.G. Wilson, 1973. Oceanographic Observations in Barrow Strait and Wellington Channel, N.W.T. Institute of Ocean Sciences, Sidney, B.C., 170 pp.
- 73-0009A** See **72-0017**
- 73-0009B** Bornhold, B., 1973. Unpublished data on Maxwell Bay and Lancaster Sound. Pacific Geoscience Centre, Sidney, B.C.
- Lewis, C.F.M., J.R. Horsman and D.I. Ross. Geological, geophysical and hydrographic studies in Lancaster Sound and Maxwell Bay, District of Franklin. Geological Survey of Canada, Paper 74-1A: 251-252.
- 73-0010** See **70-0010**
- 73-0011** See **72-0017**
- 74-0013A** See **72-0017**
- 74-0013B** Bornhold, B., 1974. Unpublished sediment data. Pacific Geoscience Centre, Sidney, B.C.
- Atlantic Geoscience Centre, 1974. Cruise Report, C.S.S. Hudson September 26 to November 1, 1974. Bedford Institute of Oceanography, 7 pp. + appendices.

- 74-0014** Thomson, D., S. Woods and J. Acreman, 1975. Marine ecology survey in the central portion of the Canadian Arctic Islands, 1974. Marine Sciences Centre, McGill University. Report prepared for Polar Gas Project, 77 pp.
- 74-0015** B.C. Research, 1975. Baseline survey of the marine environment at Strathcona Sound, N.W.T. Report to Strathcona Mineral Services, Project 1552, 84 pp. + appendices.
- Bohn, A. and B.W. Fallis, 1978. Metal concentrations (As, Cd, Cu, Pb and Zn) in shorthorn sculpins, Myoxocephalus scorpius (linnaeus), and Arctic char, Salvelinus alpinus (linnaeus), from the vicinity of Strathcona Sound, N.W.T. Water Research, Vol. 12, No. 9, 1978, pp. 659-663.
- 74-0023** Dunbar, M.J. and J.C. Acreman, 1980. Standing crops and species composition of diatoms on sea ice from Robeson Channel to the Gulf of St. Lawrence. Marine Sciences Centre, McGill University, Ophelia, 19(1): 61-72.
- 74-0024** See 72-0017
- 74-0026** Fallis, B.W., 1982. Trace metals in sediments and biota from Strathcona Sound, N.W.T.; Nanisivik marine monitoring programme, 1974-1979. Canadian Technical Report of Fisheries and Aquatic Sciences No. 1082, 34 pp.
- 75-0013** Sekerak, A.D., D. Thomson, H. Bain and J. Acreman, 1976. Summer surveys of the marine environment of Creswell Bay, Somerset Island and Assistance Bay, Cornwallis Island, N.W.T. 1975. LGL Ltd., Report prepared for Polar Gas Project, 215 pp.
- 75-0029** Bornhold, B., 1975. Unpublished sediment data on Viscount Melville Sound and Strathcona Sound. Pacific Geoscience Centre, Sidney, B.C.
- Bornhold, B., 1975. Marine surficial geology: central and eastern Arctic. Project 730031, Geological Survey of Canada, Paper 76-1A: 29-31.
- 75-0030** Fallis, B., 1975. Unpublished sediment data for Strathcona Sound. Freshwater Institute, Winnipeg, Manitoba.
- See 74-0026
- 75-0031A** Walker, E.R., 1977. Aspects of oceanography in the archipelago. Institute of Ocean Sciences, Sidney, B.C., 186 pp.
- 75-0031B** Bohn, A., 1979. Trace metals in fucoid algae and purple sea urchins near a high Arctic lead/zinc ore deposit. Marine Pollution Bulletin, 10: 325-327.
- See 75-0031 A

- 75-0031B** Bohn, A. and R.O. McElroy, 1976. Trace metals (As, Cd, Cu, Fe and Zn) in arctic cod, Boreogadus saida, and selected zooplankton from Strathcona Sound, northern Baffin Island. J. Fish. Res. Board Canada, 33: 2836-2840.
- 75-0033** Smith, T.G. and F.H.J. Armstrong. Mercury and selenium in ringed and bearded seal tissues from Arctic Canada. Arctic 31(2): 75-84.
- 76-0007** Sadler, H.E., H.V. Serson and R.K. Chow, 1979. The oceanography of Fury and Hecla Strait. Defence Research Establishment Pacific, Victoria, B.C., Technician Memorandum 79-11, 61 pp. + appendices.
- 76-0008** Sekerak, A.D., R.A. Buchanan, W.B. Griffiths and M.G. Foy, 1976. Biological oceanographic studies in Lancaster Sound, 1976. LGL Ltd., Report prepared for Norlands Petroleum Ltd., 169 pp. + appendices.
- 76-0009** Morison, S.R. and R.B. Taylor, 1978. Physical characteristics and seasonal changes in an arctic estuarine environment. Geol. Surv. Can., Paper 78-1B: 101-106.
- 76-0010** Bain, H., D. Thomson, M. Foy and W. Griffiths, 1977. Marine ecology of fast-ice-edges in Wellington Channel and Resolute Passage, N.W.T. LGL Ltd. Report prepared for Polar Gas Project, 215 pp. + appendices.
- 76-0011A** Chemical Oceanography Division, Cruise report no. 76-023, July 23 to August 19, 1976. Bedford Institute of Oceanography.
- Tan, F.C. and P.M. Strain, 1980. The distribution of sea ice meltwater in the Eastern Canadian Arctic. Journal of Geophysical Research, 85(C4): 1925-1932.
- 76-0011B** Bedford Institute of Oceanography, Cruise report no. 76-025.
- Bornhold, B., 1976. Unpublished sediment data on Strathcona Sound. Pacific Geoscience Centre, Sidney, B.C.
- 76-0012** See **74-0026**
- 76-0021** Renewable Resources Consulting Services Ltd., 1977. Analyses of heavy metals and chlorinated hydrocarbon contamination of five sea birds from Lancaster Sound. 55 pp.
- 76-0022** See **75-0033**
- 77-0014** Chemical Oceanography Division, Isotope Chemistry Data Listing, Nutrient Chemistry Data Listing, Organic Chemistry Data Listing, Physical Chemistry Data Listing, Cruise 77-024. Atlantic Oceanographic Laboratory, Bedford Institute of Oceanography.
- Levy, E.M., 1979. Concentration of petroleum residues in the waters and sediments of Baffin Bay and the Eastern Canadian Arctic - 1977. Atlantic Oceanographic Laboratory, Bedford Institute of Oceanography, 34 pp.

Irwin, B., M. Hodgson, P. Dickie and T. Platt, 1978. Phytoplankton productivity experiments in Baffin Bay and adjacent inlets 22 August to 18 September, 1977. Marine Ecology Laboratory, Bedford Institute of Oceanography, Fisheries and Marine Services Data Report No. 82, 52 pp.

Atlantic Oceanographic Laboratory, Cruise Report No. 77-024 22 August to 17 September, 1977. Bedford Institute of Oceanography.

Campbell, J.A. and P.A. Yeats, 1982. The distribution of manganese, iron, nickel, copper and cadmium in the waters of Baffin Bay and the Canadian Archipelago. *Oceanologica Acta*, 5(2): 161-168.

Top, Z., W.B. Clarke, W.C. Eismount and E.P. Jones, 1980. Radiogenic helium in Baffin Bay bottom water. *Journal of Marine Research*, 38(3): 435-452.

Jones, E.P. and A.R. Coote, 1980. Nutrient distributions in the Canadian Archipelago: indicators of summer water mass and flow characteristics. *Canadian Journal of Fisheries and Aquatic Sciences*, 37(4): 589-599.

Campbell, J.A. and D.H. Loring, 1980. Baseline levels of heavy metals in the waters and sediments of Baffin Bay. *Marine Pollution Bulletin*, Vol. 11, pp. 257-261.

See 76-0011 A (Tan and Strain)

- 77-0015** Thomson, D., W.E. Cross, H. Bain and L. Patterson, 1978. Aspects of the spring and summer marine environment of Brentford Bay, Boothia Peninsula, N.W.T. LGL Ltd. Report prepared for Polar Gas Project, 203 pp.
- 77-0016** Buchanan, R.A., W.E. Cross and D.H. Thomson, 1977. Survey of the marine environment of Bridport Inlet, Melville Island. LGL Ltd. Report prepared for Petro-Canada, 265 pp.
- 78-0006** Lake, R.A., 1978. An oceanographic study of Bridport Inlet, Melville Island, N.W.T. - Part I, August 1978. Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C., 12 pp.
- 78-0020** Chemical Oceanography Division. Location file cruise 78-026, 1978. Bedford Institute of Oceanography.
- Irwin, B., W.G. Harrison, C.L. Gallegos and T. Platt, 1980. Phytoplankton productivity experiments and nutrient measurements in the Labrador Sea, Davis Strait, Baffin Bay and Lancaster Sound from 26 August to 19 September 1978. *Canadian Data Report of Fisheries and Aquatic Sciences* No. 213.
- Levy, E.M., C.C. Cunningham, C.D.W. Conrad and J.D. Moffat, 1977. The determination of dissolved oxygen in sea water. Bedford Institute of Oceanography, Report series/BI-R-77-9, 17 pp.

- 78-0021** McLaren, P., W.B. Barrie, J.M. Sempels, R.A. Sieffert, R.B. Taylor and D. Thomson, 1981. Coastal and environmental data from eastern Lancaster Sound and northeastern Baffin Island, N.W.T. Bedford Institute of Oceanography, Data Series/BI-D-81-1, 283 pp.
- 78-0022** Sekerak, A.D., R.A. Buchanan, M.G. Foy, H. Bain, G.L. Walder and H.E. Stallard, 1979. Studies of plankton in northwest Baffin Bay and adjacent waters July - October, 1978. LGL Ltd., Executive Summary, 412 pp.
- 79-0014** Lake, R.A., 1979. An oceanographic study of Bridport Inlet, Melville Island, N.W.T. - Part II, March 1979. Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C., 288 pp.
- Lake, R.A., 1979. An oceanographic study of Bridport Inlet, Melville Island, N.W.T. - Part II, March to August, 1979. Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C. (unpublished manuscript), 59 pp.
- 79-0024** See **74-0026**
- 80-0007** Marine Ecology Laboratory. Location File cruise 80-027, 1982. Chemical Oceanography Division, Bedford Institute of oceanography.
- Bedford Institute of Oceanography. Biological oceanography report on C.S.S. Hudson. Cruise 80-027, July 24 - August 29, 1980.
- 80-0009** Lake, R., D. Richards and R. Cooke, 1980. Bridport Inlet (unpublished O₂ data), Frozen Sea Research Group, Institute of Ocean Sciences, Sidney, B.C.
- 80-0018** Eaton, R.D.P. and J.P. Farrant, 1982. The polar bear as a biological indicator of the environmental mercury burden. Arctic, 35(3): 422-425.
- 81-0007** Brooks, D.J., 1981. Arctic oceanographic survey, Barrow Strait. 1981 Field Report, Canadian Centre for Inland Waters (unpublished manuscript), 66 pp.

VOLUMES AVAILABLE

1. Beaufort Sea: Physical Oceanography - Temperature, Salinity, Currents and Water Levels.
2. Beaufort Sea: Chemical Oceanography.
3. Northwest Passage: Physical Oceanography - Temperature, Salinity, Currents and Water Levels.
4. Northwest Passage: Chemical Oceanography.

OTHER RELATED PUBLICATIONS

Thomas, D.J., R.W. Macdonald and A.B. Cornford, 1983. A comparison of natural and petroleum resource development related fluxes of metals, hydrocarbons and nitrogen, phosphorus and silicon to the Beaufort Sea. Can. Tech. Rep. of Hydrogr. Ocean Sci. 30: 43 pp.

